



BIROn - Birkbeck Institutional Research Online

Forrester, Gillian and Rodriguez, A. (2015) Slip of the tongue: implications for evolution and language development. *Cognition* 141 , pp. 103-111. ISSN 0010-0277.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/16689/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Slip of the tongue:

Implications for evolution and language development

Gillian S. Forrester^{a*} and Alina Rodriguez^{b,c}

^aDepartment of Psychology, Faculty of Science and Technology, University of
Westminster, 115 New Cavendish Street, London, England, W1W 6UW

^bDepartment of Epidemiology and Biostatistics, School of Public Health, Imperial
College London, Faculty of Medicine, St Mary's Campus, Norfolk Place, Paddington,
London, W2 1PG, UK

^cDepartment of Psychology, Mid Sweden University, Östersund, Sweden

Author contact details:

*Gillian S. Forrester: email: g.forrester@westminster.ac.uk

Alina Rodriguez: a.rodriguez@imperial.ac.uk

*Correspondence concerning this article should be addressed to:

Dr. Gillian S. Forrester

Department of Psychology, Faculty of Science and Technology,
115 New Cavendish Street, University of Westminster, London, W1B 2UH

Email: g.forrester@westminster.ac.uk

26 **Abstract**

27 A prevailing theory regarding the evolution of language implicates a gestural stage
28 prior to the emergence of speech. In support of a transition of human language from a
29 gestural to a vocal system, articulation of the hands and the tongue are underpinned
30 by overlapping regions dominant within the left hemisphere. Behavioral studies
31 demonstrate that human adults perform sympathetic mouth actions in imitative
32 synchrony with manual actions. Additionally, right-handedness for precision manual
33 actions in children has been correlated with the typical development of language,
34 while a lack of hand bias has been associated with psychopathology. It therefore
35 stands to reason that sympathetic mouth actions during fine precision motor action of
36 the hands may be lateralized. We employed a fine-grained behavioral coding
37 paradigm to provide the first investigation of tongue protrusions in typically
38 developing 4-year old children during cognitive tasks that required varying degrees of
39 manual action: precision motor action, gross motor action and no motor actions. The
40 rate of tongue protrusions was influenced by the motor requirements of the task and
41 tongue protrusions were significantly right-biased for *only* precision manual motor
42 action ($p < .001$). From an evolutionary perspective, tongue protrusions can drive new
43 investigations of how an early human communication system transitioned from hand
44 to mouth. From a developmental perspective, the present study may serve to reveal
45 patterns of tongue protrusions during the motor development of typically developing
46 children. Further research may contribute to our understanding of cerebral
47 lateralization of cognitive function.

48

49 **Keywords:** tongue, language, cerebral lateralization, typically developing children

50

51 **1. Introduction**

52 The tongue is one of the largest muscles in the human body, controlled by the
53 hypoglossal nerve (twelfth cranial nerve). Following brain injury, tongue protrusions
54 can be used as a diagnostic tool to determine the anatomical level of damage (Riggs,
55 1984). Patients are asked to stick their tongue out straight. Damage to tongue muscles
56 or the hypoglossal nerve can result in tongue weakness, causing the tongue to deviate
57 towards the weak side (ipsilateral). Conversely, lesions originating from the motor
58 cortex will cause contralateral tongue weakness. Such anatomical organization
59 suggests contralateral hemispheric motor control of articulatory left and right tongue
60 actions. Although the primary role of the tongue is for mastication, swallowing and
61 gustation, a secondary, but critical role of the tongue is phonetic articulation.

62 Moreover, the tongue also becomes active in nonverbal synchrony with manual motor
63 tasks. For example, have you ever found yourself performing a manual task and
64 notice that your tongue is pressed between your lips with the tip protruding from the
65 mouth? This behavior is commonly observed in young children (Mason & Proffit,
66 1974) and may be noticeable in adults when pursuing high precision manual dexterity
67 that requires focused attention, like threading a needle (Givens, 2002). To date, the
68 origin of this motor action and the basis of its functionality, have gone unexplored.

69
70 To date, the literature concerning tongue protrusions concentrates on involuntary
71 tongue protrusion, also called ‘tongue thrust’, ‘reverse swallow’ or ‘immature
72 swallow’. Tongue thrust has been mainly associated with psychopathology and is
73 considered to be an orofacial muscular imbalance whereby the tongue “protrudes
74 through the anterior incisors during swallowing, speech, and while the tongue is at
75 rest” (Council on Children with Disabilities, 2006). Tongue thrust has been

76 documented in patients with Dystonia (Schneider, Aggarwa, Dupont, Tisch,
77 Limousin, Quinn & Bhatia, 2006), Down's syndrome (Limbrock, Fischer-Brandies &
78 A Valle, 1991), Rett syndrome (Einspieler, Kerr & Prechtel, 2008), Tourette's syndrome
79 (Strassing, Hugo & Mueller, 2004), Angelman syndrome (Williams et al., 2006) and
80 in children with non-organic failure to thrive (Mathisen, Skuse, Wolke & Reilly,
81 1989). However, tongue thrust has also been reported in 67-95% of typically
82 developing children aged 5-8 years. It is thought that for most children, it will
83 extinguish by the age of six, as a typical swallowing motor action is developed
84 (Mason & Proffit, 1974). In contrast, involuntary tongue thrust relating to reflexive
85 swallowing actions may differ in function and neural origin from the tongue
86 protrusions produced by typically developing individuals during tasks of high
87 concentration.

88

89 Theories regarding the evolutionary and developmental basis of tongue protrusions
90 during tasks of concentration range from: motor overflow during attentional processes
91 (e.g. Waber, Mann & Merola, 1985), to the physical rejection of the bottle or breast to
92 by infants to indicate satiation (e.g. Morris, 1978). While the former has not been
93 formally investigated, in the latter scenario, it has been hypothesized that the tongue
94 protrusion action is retained throughout development as a symbol of rejection,
95 implying: 'back off' or 'leave me in peace' (e.g. Ingram, 1990). Anecdotal evidence
96 of such an interpretation can be found in Western culture where tongue protrusions
97 have become a popular symbol utilized by celebrities to ward off unwanted public
98 attention. However, if a protruded tongue results from an involuntary, innate behavior
99 to indicate satiation, one should find evidence of this symbolic defiance gesture across
100 cultures. While there is a paucity of empirical data to consider, contrary to the above

101 hypothesis, in Tibet, the protrusion of the tongue is considered to be a greeting
102 (Tsering, 2007).
103
104 A more compelling theory regarding the origins of nonverbal mouth actions (not
105 specific to protrusions) is rooted in the evolution and development of language
106 processes. It has been hypothesized that human speech evolved from a
107 communication system based on hand gestures (Armstrong, Stokoe & Wilcox, 1995),
108 supported by the properties of a ‘mirror’ neuron system (Rizzolatti & Arbib, 1998).
109 This system serves both the production and perception of actions, potentially making
110 a critical contribution to the emergence and development of motor skills for willed
111 communication (Gallese, Fadiga, Fogassi & Rizzolatti, 1996).
112
113 Behavioral evidence from chimpanzee and human studies supports such a synergy.
114 For example, chimpanzees generated sympathetic mouth movements significantly
115 more often during tasks requiring fine motor manipulation compared with tasks
116 requiring gross motor actions (Waters & Fouts, 2002). In humans, Gentilucci,
117 Benuzzi, Gangitano & Grimaldi (2001) demonstrated that the pronunciation of a
118 syllable could be selectively disrupted when producing a simultaneous grasping action
119 with the hand aimed at target objects of a non-congruent size of the mouth
120 vocalization. The finding suggests that the fine motor articulation required for
121 grasping is processed similarly by both hand and mouth in humans, thus they tend to
122 complement each other. In fact, so tightly are the two motor systems entwined that
123 when either gesture or speech is disrupted the other becomes delayed (Chu &
124 Hagoort, 2014).
125

126 Neuroimaging findings indicate close links between brain regions related to speech
127 production and those controlling movement of the hands and arms (Erhard, Kato,
128 Strupp, Andersen, Adriany, Strick & Ugurbil, 1996; Rizzolatti & Arbib, 1998;
129 Rizzolatti & Craighero, 2004). Specifically, Broca's area is activated when imitating
130 hand movements and preparing grasps (Iacoboni, Woods & Mazziotta, 1998) in
131 addition to actual or internal speech (Hinke, Hu, Stillman, Kim, Merkle, Salmi &
132 Ugurbil, 2003), supporting the notion of a common neural substrate for hand and
133 mouth articulation. Thus, in modern humans, there exists an association between
134 speech and gesture that transcends the speaker to communicate, whereby vocalization
135 and the synchronous arm movements appear intertwined in the mutual cognitive
136 activity of language and remain linked throughout the lifespan (Iverson & Thelen,
137 1999).

138

139 In humans, the observation of grasp alone can activate preparation of the same motor
140 act (Fadiga, Fogassi, Pavesi & Rizzolatti, 1995). These findings are reminiscent of the
141 observed and actual grasping behaviors discovered in monkey (Rizzolatti, Camarda,
142 Fogassi, Gentilucci, Luppino & Matelli, 1988), underpinned by a mirror neuron
143 system. Broca's region in humans and the analogous neural region in the monkey
144 brain (F5) may act as a supramodal processor for planned, structured action sequences
145 represented by both the hands and the mouth (e.g. Petersson & Hagoort, 2012;
146 Pulvermüller & Fadiga, 2010). This sort of system would support perception-action
147 coupling and may have catalyzed the emergence of syntactic processes found in
148 modern human language (e.g. Forrester, Leavens, Quaresmini & Vallortigara, 2011;
149 Forrester, Quaresmini, Leavens, Spiezio & Vallortigara, 2012; Tabiowo & Forrester,
150 2013). Such a processor, dominant within the left hemisphere may have also given

151 rise to human population-level right-handedness (Annett, 2002), for efficiency in
152 carrying out sequences of structured motor actions (e.g. Forrester, Quaresmini,
153 Leavens, Mareschal & Thomas, 2013).

154

155 Modern humans demonstrate population-level right-handedness for both object
156 manipulation and gesture (Marchant, McGrew & Eibl-Eibesfeldt, 1995). Recent
157 studies of child handedness indicate that right-handedness is correlated with typical
158 language development (Kastner-Koller & Keimann, 2007) and that consistent hand
159 dominance in early infancy (6-14 months) is associated with subsequent advanced
160 language skills (18-24 months) (Nelson, Campbell & Michel, 2014). Moreover, a lack
161 of hand dominance (e.g. mixed-handed, ambi-preference) may indicate disruption to
162 the cerebral lateralization of language function (e.g. Crow, Crow, Done & Leask,
163 1998; Delcato, 1966; Orton, 1937; Rodriguez, Kaakinen, Moilanen, Taanila,
164 McGough, Loo & Järvelin, 2010; Yeo, Gangestad & Thoma, 2007; Yeo, Gangestad,
165 Thoma, Shaw & Repa, 1997). Thus, strength of handedness has been proposed to be a
166 useful behavioral marker of children at risk for dysfunction of subsequent language
167 processes long before language develops (e.g. Forrester, Pegler, Thomas &
168 Mareschal, 2014). Although it has never been systematically investigated, one may
169 hypothesize that tongue protrusions produced during manual actions may comprise a
170 lateralized component, consistent with a left hemisphere dominant neural generator.

171

172 The present study sought to investigate the frequency and laterality of tongue
173 protrusions in order to provide the first empirical dataset reflecting tongue protrusions
174 in typically developing four year-old children. Tongue protrusions were assessed
175 during six tasks of high concentration requiring either: fine motor object

176 manipulation, gross motor object manipulation or no object manipulation. Based on
177 the limited existing evidence we hypothesized increasing frequency of tongue
178 protrusions during tasks requiring prehension and additionally considered a left
179 hemisphere (right side) bias in the direction of protrusion. Findings are discussed in
180 light of both developmental and evolutionary theories.

181

182 **2. Material and Methods**

183

184 *2.1. Participants*

185 Fourteen typically developing male (n = 8) and female (n = 6) children (age range:
186 53-56 months; mean age = 54.21 months) were randomly sampled from a previously
187 recorded cohort of 150 children during their participation in a neuropsychological
188 battery of cognitive tasks (see Rodriguez & Waldenström, 2008). Rationale for the
189 age range was predicated by a previous report of tongue thrust identified in 67-95% of
190 typically developing children aged 5-8 years, but tending to extinguish by the age of
191 six (Mason & Proffit, 1974). Importantly, participants were considered to have
192 reached an age by which any concerns with delayed language development would
193 have been identified. Children participating in this study were reported to have no
194 symptoms of language dysfunction. All children were right-handed as deemed by
195 maternal and self-reports. All children came from two-parent homes with an average
196 disposable monthly income of 25000 Swedish Crowns, which corresponds to Swedish
197 national average representing 5th-8th income deciles (Swedish Statistical Central
198 Bureau).

199

200 All behavior was digitally recorded in the home of the individual participants with the
201 participant's mother close by. The procedures for this study involving human
202 participants were in accordance with ethical standards of the responsible committee
203 on human experimentation (institutional and national) and with the spirit of the
204 Helsinki Declaration of 1975, as revised in 2000.

205

206 *2.2. Data Collection*

207 Tongue protrusion behaviors were observed during a subset of the neuropsychological
208 test battery of assessed tasks (Small World, Board Game, Lock and Key, Knock and
209 Tap, Picture Block, Story Recall). This set of challenging tasks were part of a battery
210 of tests conducted to assess cognitive, behavioral, and emotional development (see
211 Rodriguez & Waldenström, 2008). The Small World and Board Game tasks were
212 performed with the child's mother and were designed to assess the mother-child
213 relationship during free-play (Small World) and structured-play (Board Game). All
214 other tasks were performed with a female experimenter. All tasks were conducted on
215 a table surface in the home of the child. All tasks except one (Story Recall) required
216 an element of object manipulation (fine motor or gross motor action) as defined by the
217 instructions. For the purposes of the present study, we were interested in the duration
218 of the task for each individual, the motor requirement of the task and the frequency
219 and laterality of spontaneous tongue protrusions produced by the child. The tasks
220 were as follows:

221

222 *Fine Motor Action*

223

224 *Small World:* subjects were provided with a small amount of small world play toys
225 such as miniature dolls, porcelain tea set, and furniture packed into a miniature
226 suitcase. Subjects were observed during independent play and/or interaction with the
227 mother for five minutes. All objects were small and some objects had small moving
228 parts, requiring fine coordinated manipulation.

229

230 *Board Game:* A challenging board game was presented to both child and mother.
231 Turn taking was required and a roll of the die determined a destination based on a
232 combination of a color and a picture. If the picture was present in the column of the
233 given color, a small playing chip was placed on this space on their own board. The
234 object of the game was to complete a full row or column before the other player and
235 thus varied in time across participants. The collection of cards and the movement of
236 playing chips across the spaces of the board required fine motor coordination.

237

238 *Lock and Key:* Subjects were provided with a 4 locked metal padlocks, ranging in
239 shape and size, and a set of five keys on a single ring. Each key opened one lock. The
240 process for opening a lock was demonstrated by the experimenter. The child was
241 given five minutes to open all the locks. This task required fine motor coordination to
242 manipulate both keys and locks.

243

244 *Gross Motor Action*

245

246 *Knock and Tap:* This task was taken from the NEPSY neuropsychological test battery
247 (Kemp, Kirk & Korkman, 2001; Korkman, Kirk & Kemp, 2000) to tap attention and
248 effortful control in four-year-olds. The experimenter engaged the child in the manual

249 motor sequence task. The experimenter sat opposite the child with hands laid flat on
250 the table. The child was asked to mirror the position. The child indicated which hand
251 s/he used most often. The experimenter explained that whenever she knocked (closed
252 fist) on the table, the child was to tap (opened palm down, e.g. slap) on the table. In
253 contrast, whenever the experimenter tapped (opened palm down) on the table the
254 child was to knock. Several practice trials were given to make sure that the child
255 understood the task instructions. Fifteen test trials followed. This task required gross
256 motor movements, and did not require any object manipulation. This task required
257 inhibition of the prepotent action, i.e. imitation of the experimenter's hand movement
258 and was not timed.

259

260 *Picture Block:* The experimenter presented the child a small, 2D square picture of a
261 bear with a ball. The experimenter and child talked about the distinctive features of
262 the picture. The child was then presented with nine approximately 2 inch square
263 blocks. Each block portrayed a small segment, i.e. 1/9th of the 2D picture on the top
264 surface. The cubes were presented in mixed order, but all correct picture segments
265 were always facing up and the child's task was to place the nine blocks to copy the
266 2D picture. Five minutes were allotted to this task. This task required the spatial
267 rotation of blocks into position in accordance with the defined picture.

268

269 *No Motor Action*

270

271 *Story Recall:* The experimenter read the Narrative Memory story from NEPSY (47,
272 48) suitable for four-year-olds. The story comprised of a complex plot involving
273 several characters and events. Children were asked to listen to the story and then were

274 asked to recall information under free and cued-recall conditions. This task did not
275 require any fine or gross manual motor actions and was not timed.

276

277 *2.3. Data Coding*

278 Videos were viewed on Windows Movie Media Player providing a viewing resolution
279 of 30 frames per second. Tongue protrusions were coded based on the following
280 criteria. A tongue protrusion was defined as any visible protrusion of the tongue from
281 or within the mouth. Although the duration of protrusions was not calculated, the start
282 of a protrusion was identified by a visible distortion of the cheek or lip, or by the
283 visible appearance of the tongue through the lips. Only the starting point of the
284 protrusion was considered. While some children performed tongue sweeps, beginning
285 with a protrusion and sweeping to the left or right, there were too few of these events
286 to be considered for further analysis. Viewing video footage of 30 frames per second
287 allowed for fine resolution coding of these events. Under these criteria, tongue
288 protrusions could be internal or external. However, internal protrusions required clear
289 visual distortion of the cheek or lips for identification. Tongue protrusions were
290 identified for lateral position i.e. directed the tip towards the left or the right of the
291 individual. When a lateral position was unclear (e.g. central), a protrusion was only
292 considered for tests of frequency and rate, but not for tests of laterality. It is possible
293 that central protrusions were lateralized, but not to an identifiable extent by the coder.
294 Any instance where one side of the mouth was otherwise engaged was not considered
295 for the final coded data. For example, if the subject was chewing something on the
296 left side of their mouth (e.g. their sleeve, a toy) and protruded their tongue to the right,
297 this was excluded from the coded data set. Tongue protrusions occurred as events
298 rather than bouts (e.g. quick successive repetitions of the same action) and were

299 analyzed accordingly. All subject footage was observed for as long as it took to reach
300 the end of all tasks, which was on average 50 minutes (+/- 10 minutes).

301

302 *2.4. Data Analysis*

303 Analyses of variance and appropriate post-hoc tests were used to assess frequencies,
304 rates and lateral biases of group-level tongue protrusions. Laterality Index scores (LI)
305 were calculated using the formula $[LI = (R-L)/(R+L)]$, with R and L being the
306 frequency counts for right and left navigational path frequency counts. LI values vary
307 on a continuum between -1.0 and +1.0, where the sign indicates the direction of
308 tongue protrusion preference. When R=L, then LI is zero, i.e. no lateral bias. Positive
309 values reflect a right protrusion while negative values reflect a left preference. The
310 absolute value depicts the strength of protrusions. In order to assess differences in the
311 frequencies of tongue protrusions across tasks, rates were calculated. Rates were
312 equal to the frequency of tongue protrusions for a given task for a specific individual
313 divided by the duration in minutes to complete the task. All statistical tests were two-
314 tailed ($\alpha < .05$).

315

316 **3. Results**

317 Raw frequencies of tongue protrusions for each individual by task are presented in
318 Table 1. Tongue protrusions frequencies are divided into left, right and central
319 directions. For ANOVA tests, where sphericity was not assumed, Greenhouse-Geisser
320 correction was used. Non-parametric Wilcoxon signed-rank tests were used for all
321 post-hoc analyses.

322

323 - Insert Table 1 -

324

325 *3.1. General Description of Tongue Protrusions*

326 Across participants, the frequency of tongue protrusions ranged between 16-49, ($M =$
327 30; $SD = 9.89$). On average, the group elicited significantly more detectable external
328 (frequencies: $M = 16.79$, $SE = 1.62$; proportions: $M = 0.562$, $SE = 0.027$) versus
329 internal tongue protrusions (frequencies: $M = 13.21$, $SE = 1.395$; proportions: $M =$
330 0.438, $SE = 0.027$) collapsed across all tasks (frequencies: $t(13) = 2.417$, $P = 0.031$;
331 proportions: $t(13) = 2.314$, $P = 0.038$). A 1-way ANOVA indicated no significant
332 difference in the frequency of tongue protrusions across tasks: small world ($M = 5.23$,
333 $SE = 3.07$); Board Game ($M = 5.50$, $SE = 2.07$); Lock and Key ($M = 4.29$, $SE = 3.34$);
334 Knock and Tap ($M = 4.14$, $SE = 3.44$); Picture Block ($M = 5.50$, $SE = 3.39$); Story
335 Recall ($M = 5.29$, $SE = 4.75$) [$F(5, 65) = 5.812$, $p = 0.277$]. However, as tasks varied
336 in duration or time to completion (see Table 2), thus rates of tongue protrusions per
337 minute (rate = (seconds to complete task/ # of tongue protrusions)/60) were also
338 calculated to equalize the weighting that each task contributed to the dataset (see
339 Table 3).

340

341 - Insert Table 2 -

342 - Insert Table 3 -

343

344 A 1-way ANOVA indicated a significant difference in rates across tasks [Small World
345 ($M = 0.90$, $SE \pm 0.15$); Board Game ($M = 0.76$, $SE \pm 0.11$); Lock and Key ($M = 0.68$,
346 $SE \pm 0.14$); Knock and Tap ($M = 1.84$, $SE \pm 0.37$); Picture Block ($M = 1.27$, $SE \pm$
347 0.25); Story Recall ($M = 0.77$, $SE \pm 0.17$) [$F(2.72, 35.41) = 4.52$, $p = 0.011$].

348 Additionally, a 1-way ANOVA revealed a significant difference in task motor

349 requirement (fine motor, gross motor and no motor) [$F(2, 26) = 6.67, p = 0.005$] (see
350 Figure 1).

351

352 - Insert Figure 1 -

353

354 Post-hoc analyses revealed that tongue protrusion rates for tasks requiring gross
355 motor actions ($M = 1.55, SE \pm 0.23$) elicited a significantly greater rate of tongue
356 protrusions than tasks requiring fine motor action ($M = 0.78, SE \pm 0.08$) ($Z = -3.42$;
357 $p = .001$), or no motor action ($M = 0.77, SE \pm 0.17$), ($Z = -2.27$; $p = .023$).

358

359 *3.2. Lateralized Tongue Protrusions*

360 Frequency of left and right tongue protrusions revealed that participants demonstrated
361 a significant bias for right tongue protrusions (frequencies: $M = 10.79, SE \pm 1.82$)
362 versus left tongue protrusions (frequencies: $M = 5.57, SE \pm 0.78$) collapsed across all
363 tasks ($Z = -2.76$; $p = .006$). (see Figure 2).

364

365 - Insert Figure 2 -

366

367 Further analyses of lateral tongue protrusion biases were conducted employing LI
368 scores. LI scores ensure equal weighting of participant contribution to the analysis
369 (see Table 4).

370

371 - Insert Table 4-

372

373 A 1-way ANOVA of laterality index scores of tongue protrusions was calculated by
374 motor condition (fine motor, gross motor and no motor), resulting in a significant
375 difference for mean LI scores across motor conditions [$F(2, 26) = 12.36, p < 0.001$]
376 (see Figure 3).

377

378 - Insert Figure 3 -

379

380 Post-hoc analyses by motor condition showed that fine motor condition ($M = 0.63, SE$
381 ± 0.11) elicited significantly more right-biased tongue protrusions compared with the
382 gross motor condition ($M = -0.08, SE \pm 0.15$) ($Z = -2.91; p = .003$) and the no motor
383 condition ($M = -0.22, SE \pm 0.17$) ($Z = -2.80; p = .005$). Additionally, mean LI scores
384 by task were as follows: Small World = .46, Board Game = .71, Lock and Key = .52,
385 Knock and Tap = .30, Picture Block = -.28, Story Recall, -.22.

386

387 **4. Discussion**

388

389 *4.1. Rates of Tongue Protrusions*

390 The findings from this investigation demonstrated that tongue protrusions commonly
391 occur in typically developing 4-year old children. Although the literature is sparse, the
392 result is consistent with an earlier report of the incidence of tongue thrust in typically
393 developing children aged 5-8 years (Mason & Proffit, 1974). In the present study,
394 fourteen participants exhibited tongue protrusions while engaging in a range of
395 cognitive tasks requiring fine motor action, gross motor action, or no motor action.
396 There were significantly more visible external tongue protrusions overall, where the
397 tongue breached the lips, compared with internal tongue protrusions, where the

398 tongue created a bulge in the cheek or lips but was not externally visible. However,
399 this result could be due to the fact that internal tongue protrusions may not always be
400 visually detectable and our findings represent a subset of all tongue protrusions.

401

402 Tasks of fine and gross manual motor action elicited tongue protrusions. This finding
403 supports the theory that hand and mouth actions sympathize with one another as a
404 result of a single system of communication that is independent of modality (McNeill,
405 1992). The motor coupling is believed to occur due to shared neural resources for
406 hand actions (Iacoboni, Woods & Mazziotta, 1998) and actual or internal speech
407 (Hinke, Hu, Stillman, Kim, Merkle, Salmi & Ugurbil, 2003) and is further supported
408 by behavioral evidence demonstrating selective disruption of speech syllables when
409 the hands are required to perform non-congruent articulations (Gentilucci et al.,
410 2001). However, tongue protrusions were also reported during the Story Recall task
411 that had no manual motor requirement. This additional finding supports the position
412 that the hands need not be active to elicit tongue protrusions. It is possible that tongue
413 protrusions will be elicited if a task involves active language processing as required
414 by the Story Recall task.

415

416 The rates of tongue protrusions differed significantly across tasks. Rates were
417 calculated to account for the varying task durations and time to completion per
418 participant. While all tasks elicited tongue protrusions in most children, gross motor
419 tasks elicited significantly more tongue protrusions than fine motor and no motor
420 tasks. This finding is in not inconsistent with our hypothesis, predicting more frequent
421 tongue protrusions in tasks of requiring prehension. However, this finding is in
422 contrast to non-human primate research reporting that chimpanzees generated

423 sympathetic mouth actions at a significantly higher frequency during tasks of fine
424 motor manipulation compared with tasks requiring gross motor manual actions
425 (Waters & Fouts, 2002). However, Waters & Fouts (2002) considered mouth actions
426 that were not specific to tongue protrusions. It is possible that the gross motor tasks in
427 the present study required a greater rate of grasping-type hand actions in comparison
428 to the fine motor tasks. Additionally, we consider that the gross motor tasks were both
429 tasks of significant difficulty. The Knock and Tap and Picture Block tasks were both
430 effortful tasks, requiring inhibition of prepotent responses and spatial manipulations,
431 respectively. Future studies may consider how grasping rate and task difficulty
432 influences tongue protrusions in typically developing children.

433

434 The tasks included in the gross motor condition included the Knock and Tap task and
435 the Picture Block task. The Picture Block task did not elicit significantly greater
436 tongue protrusion rate than other tasks (aside from the Board Game task). The Knock
437 and Tap task, however, did elicit significantly more tongue protrusions than all fine
438 motor and no motor tasks. It is possible that the opening and closing of the hand
439 required by the fifteen trials was sufficient to elicit complementary and sympathetic
440 tongue protrusions. Alternatively, we consider the structure of the Knock and Tap
441 task. This task possessed structured rules, rapid turn-taking and hand gesturing
442 performed with only the dominant right hand. Participants were asked to respond with
443 the opposite hand position as the experimenter. The task measures effortful control
444 and the ability to inhibit behavioral impulsivity of the prepotent response (i.e.
445 imitation of the experimenter's hand position) and may have also required an element
446 of symbolic representation. This process may involve internal speech rehearsal of the
447 task rules to actively control hand movements. One interpretation of the finding is that

448 the Knock and Tap task required foundational components of the communication
449 system, engaging both symbolic hand gestures and the internal rehearsal of the verbal
450 instructions. The task elements may even resemble proto language processes both in
451 turn-taking sequences and symbolic representation of manual gestures. While
452 structured sequences are known to be a distinctive component of language (e.g.
453 Hauser, Chomsky & Fitch, 2002), it has been suggested that they also appear in
454 nonlinguistic domains such as object manipulation and gesture (for a review see,
455 Tettamanti, 2003). The rule-based motor activity required by the Knock and Tap task
456 may be likened to sequences of behavioral units, possessing the properties of an
457 action-based proto-syntax prior to the emergence of speech (Corballis, 2009). One
458 hypothesis is that sympathetic tongue protrusions increased with tasks demand for
459 rule-based structured sequences of action and the comprehension and production of
460 symbolic hand gestures (e.g. Gentilucci et al., 2001). Based on evolutionary theory,
461 goal directed sequences of actions are foundational components of human
462 communication driven by left hemisphere dominant processes that can manifest as
463 lateralized motor action (MacNeilage Rogers & Vallortigara, 2009).

464

465 *4.2. Laterality of Tongue Protrusions*

466 A significant group-level right side bias was revealed for the frequency of tongue
467 protrusions. The motor-level analyses demonstrated that fine motor tasks revealed
468 right-biased tongue protrusions. Laterality was next explored using laterality index
469 (LI) scores across fine motor, gross motor and no motor task groups. Unlike tests of
470 frequency, LI scores ensured equal weighting of each task to the analysis. The fine
471 motor action condition revealed significantly right-lateralized tongue protrusions
472 compared with the gross action condition and the no motor action condition.

473 Additionally, all three tasks revealed mean LI scores consistent with a strong right
474 bias (e.g. Oldfield, 1971).
475
476 We considered that all fine motor tasks required precision grasp and was likely to be
477 conducted by the dominant right hand and left hemisphere. The Small World task
478 included a variety of small dollhouse toys and dolls with manipulable limbs. The
479 Board Game task required moving a token across a board and the manipulation of
480 small flat discs that required precision grasp to collect. The Key and Lock task
481 required bimanual coordinated action (e.g. McGrew & Marchant, 1997) to open pad
482 locks. One hand (non-dominant) held a lock in a power grip while the other hand
483 (dominant) used a precision grasp to manipulate a key. One interpretation of this
484 finding is that fine motor tasks precipitate use of the dominant hand because it is more
485 dexterous in for operations involving sequences of fine manipulation. Studies of
486 cerebral lateralization implicate the left hemisphere and the right hand dominant for
487 such processes in the majority of the population (e.g. MacNeilage et al. 2009). We
488 propose that the dominant hand elicited lateralized sympathetic tongue action driven
489 by the support of common left hemisphere dominant neural system for the motor
490 structures that underpin communication processes (McNeill, 1992).
491
492 Gross motor tasks did not reveal a lateral tongue protrusion bias. Although the Knock
493 and Tap task did not require precision grip, it did demonstrated a weak right biased LI
494 score, possibly due to the fact that it required the use of the dominant hand. The
495 Picture Block task conversely, demonstrated a weak left biased LI score. A potential
496 reason this task did not reveal a lateral bias may have been because it did not require a
497 dominant hand. Blocks were easily slid across the surface of the table and did not

498 require turning, as the correct pictures were already oriented face-up for the
499 participant. Studies of primate manual laterality have found that gross motor actions
500 (e.g. reaching) can often fail to exhibit a significant hand preference as actions lack
501 the precision motor skill required for grasping (for a review see: Hopkins, 2006).
502

503 The present study offers the first investigation of tongue protrusions during cognitive
504 tasks requiring varying degrees of motor precision. We report on spontaneous tongue
505 protrusions in a population of typically developing children and suggest that tongue
506 protrusions are commonly exhibited by typically developing right-handed children.
507 Tongue protrusions were detected both internally and externally to the mouth
508 suggesting that this behavior may not cease in adulthood, but conscious awareness of
509 one's physical actions may cause tongue actions to become less detectable in order to
510 conform with social norms. Our findings support an intrinsic connection between
511 actions of the mouth and hands that is consistent with behavioral studies indicating
512 that vocalizations are accompanied by spontaneous and synchronous rhythmic hand
513 movements, visible from early infancy (e.g. Masataka, 2001). Our findings suggest
514 that hand and tongue actions possess a reciprocal relationship such that when
515 structured sequences of hand actions are performed they are accompanied by
516 spontaneous and synchronous tongue action. The detection of lateralized tongue
517 protrusions is consistent with a left hemisphere dominant unified communication
518 system involving both the hands and the mouth (McNeill 1992) and additionally is
519 consistent with a gestural origin of language position (Armstrong, Stokoes & Wilcox,
520 1995; Corballis, 2002). To further explore the evolution of speech and gesture, future
521 research may consider whether tongue protrusions increases in rate, strength of
522 laterality and temporal synchrony during manual motor tasks that possess

523 foundational structured components of communication (e.g. hierarchical sequences of
524 actions). Due to the overlapping neural resources underpinning hand and mouth motor
525 capabilities, the derivation of motor action patterns provides a novel method to draw
526 inference about the evolution of different cognitive abilities.

527

528 **ACKNOWLEDGMENTS**

529 We thank R.-P. Grossett for assistance with video data coding and Angela Fernholm,
530 Pia Risholm-Mothander, and Karin Brocki for data collection. We thank Ulla
531 Waldenström for the launch of the original cohort (Women’s Experience of
532 Childbirth) from which this dataset is derived. This data collection was funded by a
533 grant from The Swedish Research Council to Alina Rodriguez (principal
534 investigator). Work on this study was partially funded by VINNMER grant
535 (Rodriguez).

536

537 **References**

538 Annett, M. (2002). *Handedness and Brain Asymmetry*. The Right Shift Theory,
539 Sussex: Psychology Press.

540

541 Armstrong, D.F., Stokoe, W.C., & Wilcox, S.E. (1995). *Gesture and the Nature of*
542 *Language*. Cambridge, MA: Cambridge University Press.

543

544 Byrne, R.W. & Byrne, J.M. (1991). Hand preferences in the skilled gathering tasks of
545 mountain gorillas (*Gorilla gorilla berengei*). *Cortex*, **27**, 521-536.

546

547 Chu, M. & Hagoort, P. (2014). Synchronization of speech and gesture: Evidence for
548 interaction in action. *Journal of Experimental Psychology: General*, **143**(4), 1726-
549 1741. <http://dx.doi.org/10.1037/a0036281>.
550

551 Corballis, M.C. (2002). *From hand to mouth: The origins of language*. Princeton, NJ:
552 Princeton University Press.
553

554 Corballis, M.C. (2009). The evolution of language. *Annals of the New York Academy*
555 *of Sciences*, **1156**, 19–43. DOI: 10.1111/j.1749-6632.2009.04423.x
556

557 Council on children with disabilities; section on developmental behavioral pediatrics;
558 Bright Futures Steering Committee; Medical Home Initiatives For Children With
559 Special Needs Project Advisory Committee (2006). Identifying infants and young
560 children with developmental disorders in the medical home: an algorithm for
561 developmental surveillance and screening. *Pediatrics*, **118**, 405-20. DOI:
562 10.1542/peds.2006-1231
563

564 Crow, T.J., Crow, L.R., Done, D.J., & Leask, S. (1998). Relative hand skill predicts
565 academic ability: global deficits at the point of hemispheric indecision.
566 *Neuropsychology*, **36**, 1275–1282. [http://dx.doi.org/10.1016/S0028-3932\(98\)00039-6](http://dx.doi.org/10.1016/S0028-3932(98)00039-6)
567

568 Delcato, C.H. (1966). *Neurological organization and reading*. Springfield, Illinois:
569

570 Erhard, P., Kato, T., Strupp, J.P., Andersen, P., Adriany, G., Strick, P.L., & Ugurbil,
571 K. (1996). Functional mapping of motor in and near Broca's area. *Neuroimage*, **3**,
572 S367. DOI: 10.1016/S1053-8119(96)80369-7
573

574 Einspieler, C., Kerr, A.M., & Prechtl, H.F.R. (2005). Is the early development of girls
575 with Rett disorder really normal? *Pediatric Research*, **57**, 696-700.
576 DOI:10.1203/01.PDR.0000155945.94249.0A
577

578 Fadiga, L., Fogassi, L., Pavesi, G., & Rizzolatti, G. (1995). Motor facilitation during
579 action observation—a magnetic stimulation study. *Journal of Neurophysiology*, **73**,
580 2608–2611.
581

582 Forrester, G.S., Leavens, D.A., Quaresmini, C., & Vallortigara G (2011). Target
583 animacy influences gorilla handedness. *Animal Cognition*, **14**, 903–907.
584 DOI 10.1007/s10071-011-0413-6
585

586 Forrester, G.S., Pegler, R., Thomas, M.S.C., & Mareschal, D. (2014). Handedness as
587 a marker of cerebral lateralization in children with and without autism. *Behavioural*
588 *Brain Research*, **15**, 14-21. DOI: 10.1016/j.bbr.2014.03.040
589

590 Forrester, G.S., Quaresmini, C., Leavens, D.A., Mareschal, D., & Thomas, M.S.C.
591 (2013). Human handedness: An inherited evolutionary trait. *Behavioural Brain*
592 *Research*, **237**, 200–206. DOI: 10.1016/j.bbr.2012.09.037
593

594 Forrester, G.S., Quaresmini, C., Leavens, D.S., Spiezio, C., & Vallortigara, G. (2012).
595 Target animacy influences chimpanzee handedness. *Animal Cognition*, **15**, 1121–
596 1127. DOI: 10.1007/s10071-012-0536-4
597

598 Gallese, V., Fadiga, L., Fogassi, L., & Rizzolatti, G. (1996). Action recognition in the
599 premotor cortex. *Brain*, **119**, 593–609. DOI.org/10.1093/brain/119.2.593
600

601 Gentilucci, M., Benuzzi, F., Gangitano, M., & Grimaldi, S. (2001). Grasp with hand
602 and mouth: a kinematic study on healthy subjects. *Journal of Neurophysiology*, **86**,
603 1685–1699.
604

605 Givens, D.V. (2002). *The nonverbal dictionary of gestures, signs & body language*
606 *clues: From Adams-apple jump to zygomatic smile*. Spokane, Washington: Center for
607 Nonverbal Studies Press.
608

609 Hauser, M.D., Chomsky, N., & Fitch, W.T. (2002). The faculty of language: what is
610 it, who has it, and how did it evolve? *Science*, **298**, 1569-1579.
611 DOI:10.1126/science.298.5598.1569
612

613 Hinke, R.M., Hu, X., Stillman, A.E., Kim, S.-G., Merkle, H., Salmi, R., & Ugurbil,
614 K.J. (2003). Functional magnetic resonance imaging of Broca's area during internal
615 speech. *Neurophysiology*, **90**, 3304-16. DOI: 10.1097/00001756-199306000-00018
616

617 Hopkins, W.D. (2006). Comparative and familial analysis of handedness in great
618 apes. *Psychological Bulletin*, **132**, 538-559. DOI:10.1037/0033-2909.132.4.538

619

620 Iacoboni, M., Woods, R.P., & Mazziotta, J.C. (1998). Bimodal (auditory and visual)
621 left frontoparietal circuitry for sensorimotor integration and sensorimotor learning.

622 *Brain*, **121**, 2135– 2143. DOI: <http://dx.doi.org/10.1093/brain/121.11.2135>

623

624 Ingram, J. (1990). *The Science of Everyday Life*. Markham, Ontario: Penguin Books

625 Canada.

626

627 Iverson, J.M., & Thelen, E. (1999). Hand, mouth, and brain: The dynamic emergence
628 of speech and gesture. *Journal of Consciousness Studies*, **6**, 19-40.

629

630 Kastner-Koller, U., Deimann, P., & Bruckner, J. (2007). Assessing handedness in pre-
631 schoolers: Construction and initial validation of a hand preference test for 4-6-year-
632 olds. *Psychology Science*, **49**, 239-254.

633

634 Kemp, S.L., Kirk, U., & Korkman, M. (2001). *Essentials of NEPSY Assessment*. John
635 Wily & Sons.

636

637 Korkman, M., Kirk, U., & Kemp, S. (2000). *NEPSY. Administrasjonsanvisningar*.
638 Stockholm: Psykologiförlaget AB.

639

640 Limbrock, G.J., Fischer-Brandies, H., & Avalle, C. (1991). Castillo-Morales'
641 Orofacial Therapy: Treatment of 67 children with Down Syndrome. *Developmental*
642 *Medicine & Child Neurology*, **33**, 296–303. DOI: 10.1111/j.1469-

643 8749.1991.tb14880.x

644

645 McGrew, W.C., & Marchant, L.F. (1997). On the other hand: current issues in and
646 meta analysis of the behavioral laterality of hand function in nonhuman primates.

647 *Yearbook of Physical Anthropology*, **40**: 201-232. DOI: 10.1002/(SICI)1096-
648 8644(1997)25+<201::AID-AJPA8>3.0.CO;2-6

649

650 McNeill, D. (1992). *Hand and mind: What gestures reveal about thought*. Chicago:
651 University of Chicago Press.

652

653 MacNeilage, P.F., Rogers, L.J., & Vallortigara, G. (2009). Origins of the left and right
654 brain. *Scientific American*, **301**, 60–67. DOI:10.1038/ scientificamerican0709-60

655

656 Marchant, L.F., McGrew, W.C., & Eibl-Eibesfeldt, I (1995). Is human handedness
657 universal? Ethological analyses from three traditional cultures. *Ethology*, **101**, 239–
658 258. DOI: 10.1111/j.1439-0310.1995.tb00362.x

659

660 Masataka, N. (2001). Why early linguistic milestones are delayed in children with
661 Williams syndrome: Late onset of hand banging as a possible rate-limiting constraint
662 on the emergence of canonical babbling. *Developmental Science*, **4**, 158–164.

663 DOI: 10.1111/1467-7687.00161

664

665 Mason, R.M., & Proffit, W.R. (1974). The tongue thrust controversy: Background and
666 recommendations. *Journal of Speech and Hearing Disorders*, **39**, 115-132.

667 DOI: 10.1044/jshd.3902.115

668

669 Mathisen, B., Skuse, D., Wolke, D., & Reilly, S. (1989). Oral-motor dysfunction and
670 failure to thrive among inner-city infants. *Developmental Medicine and Child*
671 *Neurology*, **31**, 293-302. DOI: 10.1111/j.1469-8749.1989.tb03998.x
672

673 Morris, D. (1978). *Manwatching: A Field Guide to Human Behaviour*. St Albans,
674 Hertfordshire, UK: Triad/Panther Books.
675

676 Nelson, E.L., Campbell, J.M., & Michel, G.F. (2014). Early handedness in infancy
677 predicts language ability in toddlers. *Developmental Psychology*, **50**, 809-14. DOI:
678 10.1037/a0033803
679

680 Oldfield, R.C. (1971). The assessment and analysis of handedness: the Edinburgh
681 inventory. *Neuropsychologia*, **9**, 97-113.
682

683 Orton, S.T. (1937). *Reading, writing, and speech problems in children*. New York:
684 Norton: Charles C Thomas Pub Ltd.
685

686 Petersson, K.M., Folia, V., & Hagoort, P. (2012). What artificial grammar learning
687 reveals about the neurobiology of syntax. *Brain and Language*, **120**, 83-95. DOI:
688 10.1016/j.bandl.2010.08.003
689

690 Pulvermüller, F., & Fadiga, L. (2010). Active perception: sensorimotor circuits as a
691 cortical basis for language. *Nature Reviews Neuroscience*, **11**, 351-360. DOI:
692 10.1038/nrn2811
693

694 Riggs, J.E. (1984). Distinguishing between extrinsic and intrinsic tongue muscle
695 weakness in unilateral hypoglossal palsy. *Neurology*, **34**, 1367–68. DOI:
696 10.1212/WNL.34.10.1367
697
698 Rizzolatti, G., & Arbib, M.A. (1998). Language within our grasp. *Trends in*
699 *Neurosciences*, **21**, 188–194. DOI: [http://dx.doi.org/10.1016/S0166-2236\(98\)01260-0](http://dx.doi.org/10.1016/S0166-2236(98)01260-0)
700
701 Rizzolatti, G., & Craighero, L. (2004). The mirror-neuron system. *Annual Review of*
702 *Neuroscience*, **27**, 169-92. DOI: 10.1146/annurev.neuro.27.070203.144230
703
704 Rizzolatti, G., Camarda, R., Fogassi, L., Gentilucci, M., Luppino, G., & Matelli, M.
705 (1988). Functional organization of inferior area 6 in the macaque monkey. II. Area F5
706 and the control of distal movements. *Experimental Brain Research*, **71**, 491–507.
707 DOI: 10.1007/BF00248741
708
709 Rodriguez, A., & Waldenström, U. (2008). Fetal origins of child non-right-
710 handedness and mental health. *The Journal of Child Psychology and Psychiatry*,
711 49(9), 967-976. DOI: 10.1111/j.1469-7610.2008.01923.x
712
713 Rodriguez, A., Kaakinen, M., Moilanen, I., Taanila, J., McGough, J.J., Loo, S., &
714 Järvelin, M.-R. (2010). Mixed-handedness is linked to mental health problems in
715 children and adolescents. *Pediatrics*, **125**, 340-49. DOI: 10.1542/peds.2009-1165
716
717 Schneider, S.A., Aggarwa, A., Bhatt. M., Dupont, E., Tisch, S., Limousin, P., Lee, P.,
718 Quinn, N., & Bhatia, K.P. (2006). Severe tongue protrusion dystonia: clinical

719 syndromes and possible treatment. *Neurology*, **26**, 940-3. DOI:
720 10.1212/01.wnl.0000237446.06971
721
722 Strassnig, M., Hugo, R., & Müller, N. (2004). Electroconvulsive therapy in a patient
723 with Tourette's Syndrome and co-morbid obsessive compulsive disorder. *The World*
724 *Journal of Biological Psychiatry*, **5**, 164-166.
725
726 Tabiowo, E., & Forrester, G.S. (2013). Structured bimanual actions and hand transfers
727 reveal population-level right-handedness in captive gorillas. *Animal Behaviour*, **86**,
728 1049-1057. DOI: 10.1016/j.anbehav.2013.09.010
729
730 Tettamanti, M. (2003). *Language acquisition and processing: hierarchically*
731 *organized cognitive processes*. Ph.D. thesis. University of Zurich.
732
733 Tsering, B.K. (2008). Tibetan culture in the 21st century. *Tibetreport*,
734 <http://tibetreport.wordpress.com/2008/12/23/tibetan-culture-in-the-21st-century/>
735
736 Waber, D.P., Mann, M.B., & Merola, J. (1985). Motor overflow and attentional
737 processes in normal school-age children. *Developmental Medicine & Child Neurology*
738 **27**, 491–497. DOI: 10.1111/j.1469-8749.1985.tb04573.x
739
740 Waters, G.S., & Fouts, R.S. (2002). Sympathetic mouth movements accompanying
741 fine motor movements in chimpanzees (*Pan troglodytes*) with implications toward the
742 evolution of language. *Neurological Research*, **24**, 174-80.
743 DOI: <http://dx.doi.org/10.1179/016164102101199585>

744

745 Williams, C.A. et al. (2006). Angelman syndrome 2005: updated consensus for

746 diagnostic criteria. *American Journal of Medical Genetics*, **1;140A**, 413–418.

747 DOI: 10.1002/ajmg.a.31074

748

749 Yeo, R.A., Gangestad, S.W., Thoma, R., Shaw, P., & Repa K (1997). Developmental

750 instability and cerebral lateralization. *Neuropsychology*, **11**, 552-61.

751 <http://dx.doi.org/10.1037/0894-4105.11.4.552>

752

753 Yeo, R.A., Gangestad, S.W., & Thoma, R.J. (2007). Developmental instability and

754 individual variation in brain development: Implications for the origin of

755 neurodevelopmental disorders. *Current Directions in Psychological Science*, **16**, 245-

756 249. DOI: 10.1111/j.1467-8721.2007.00513.x

757

758 Figure Legends

759

760 Figure 1.

761

762 Figure 1. Mean rates of tongue protrusions across motor conditions.

763

764 Figure 2.

765

766 Figure 2. Right and left tongue protrusions collapsed across all tasks.

767

768 Figure 3.

769

770 Figure 3. Tongue protrusion mean laterality index scores across motor conditions.

771

Highlights

- Tongue and hand articulations are controlled by left hemisphere biased brain regions
- Tongue protrusions in children were right lateralized for only precision manual tasks
- The rate of tongue protrusions was influenced by both motor and language syntax
- Tongue protrusions provide a new method to study language evolution and development

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Table 1. Left, right and central tongue protrusion frequencies by task and motor condition.

P	Fine Motor									Gross Motor						No Motor		
	SW (L)	SW (R)	SW (C)	BG (L)	BG (R)	BG (C)	LK (L)	LK (R)	LK (C)	KT (L)	KT (R)	KT (C)	BL (L)	BL (R)	BL (C)	SR (L)	SR (R)	SR (C)
1	3	3	0	1	6	3	2	6	2	0	4	1	4	3	2	0	2	0
2	0	0	2	0	1	4	0	3	1	0	0	0	5	2	3	0	0	0
3	0	4	2	0	5	3	0	2	2	0	5	5	2	6	2	2	0	1
4	0	1	2	0	1	4	0	3	2	0	0	2	0	0	1	0	0	5
5	0	2	5	0	0	4	0	3	3	0	0	1	2	0	1	1	0	0
6	1	2	2	0	2	4	0	1	1	0	0	2	0	0	0	5	2	8
7	3	1	4	0	2	2	2	2	0	0	2	3	1	0	3	1	9	3
8	0	1	3	2	0	2	0	0	4	0	0	3	3	0	1	1	0	5
9	0	1	3	1	4	1	1	0	0	3	1	0	1	4	2	3	4	2
10	1	4	4	1	5	2	0	4	7	1	2	8	5	0	4	0	0	1
11	3	4	5	0	2	3	0	0	1	0	0	2	3	1	4	1	0	3
12	0	0	0	0	2	2	0	0	0	1	2	3	0	4	2	0	0	0
13	0	2	1	0	1	1	0	1	0	0	0	0	2	2	0	2	1	6
14	2	1	2	0	4	2	1	4	2	0	3	4	2	0	0	3	2	1
M	0.93	1.86	2.50	0.36	2.50	2.64	0.43	2.07	1.79	0.36	1.36	2.43	2.14	1.57	1.79	1.36	1.43	2.50
SD	1.27	1.41	1.61	0.63	1.95	1.08	0.76	1.86	1.93	0.84	1.69	2.21	1.70	1.99	1.37	1.50	2.50	2.59

P = participant, SW = Small World, BG = Board Game, LK = Lock and Key, KT = Knock and Tap, PB = Picture Block, SR = Story Recall; (l) = left, (r) = right, (c) = central, M = mean, SD = standard deviation

Table 2. Time to complete task in seconds.

P	SW	BG	LK	KT	PB	SR
1	380	540	410	97	335	354
2	355	531	423	105	174	338
3	319	699	383	125	356	330
4	360	552	393	116	412	333
5	359	422	240	73	224	365
6	342	471	400	131	420	444
7	401	565	376	151	250	442
8	545	863	415	133	334	407
9	334	344	421	86	406	460
10	335	346	411	206	229	334
11	336	180	423	123	209	391
12	318	456	424	207	398	367
13	331	472	391	124	224	400
14	290	418	384	140	160	377
M	357.50	489.93	392.43	129.79	295.07	381.57
SD	60.53	163.20	46.88	38.69	94.39	44.05

P = participant, SW = Small World, BG = Board Game, LK = Lock and Key, KT = Knock and Tap, PB = Picture Block, SR = Story Recall, M = mean, SD = standard deviation

Table 3. The rate of tongue protrusions by motor condition and task

P	Fine Motor			Gross Motor		No Motor
	SW	BG	LK	KT	PB	SR
1	0.95	1.11	1.46	3.09	1.61	0.34
2	0.34	0.56	0.57	0.00	3.45	0.00
3	1.13	0.69	0.63	4.80	1.69	0.55
4	0.50	0.54	0.76	1.03	0.15	0.90
5	1.17	0.57	1.50	0.82	0.80	0.16
6	0.88	0.76	0.30	0.92	0.00	2.03
7	1.20	0.42	0.64	1.99	0.96	1.76
8	0.44	0.28	0.58	1.35	0.72	0.88
9	0.72	1.05	0.14	2.79	1.03	1.17
10	1.61	1.39	1.61	3.20	2.36	0.18
11	2.14	1.67	0.14	0.98	2.30	0.61
12	0.00	0.53	0.00	1.74	0.90	0.00
13	0.54	0.25	0.15	0.00	1.07	1.35
14	1.03	0.86	1.09	3.00	0.75	0.95
M	0.90	0.76	0.68	1.84	1.27	0.80
SD	0.14	0.11	0.14	0.37	0.25	0.16

P=participant, SW = Small World, BG = Board Game, LK = Lock and Key, KT = Knock and Tap, PB = Picture Block, SR = Story Recall, M = mean, SD = standard deviation

Table 4. Laterality index scores by motor condition

P	Fine Motor	Gross Motor	No Motor
1	0.43	0.27	1.00
2	1.00	-0.43	0.00
3	1.00	0.69	-1.00
4	1.00	0.00	0.00
5	1.00	-1.00	-1.00
6	0.67	0.00	-0.43
7	0.00	0.33	0.80
8	-0.33	-1.00	-1.00
9	0.43	0.11	0.14
10	0.73	-0.50	0.00
11	0.33	-0.50	-1.00
12	1.00	0.71	0.00
13	1.00	0.00	-0.33
14	0.50	0.20	-0.20
M	0.63	-0.08	-0.22
SD	0.42	0.54	0.64

P = participant, M = mean, SD = standard deviation