



BIROn - Birkbeck Institutional Research Online

Litak, T. and Mikulás, Szabolcs and Hidders, Jan (2016) Relational lattices: from databases to Universal Algebra. *Journal of Logical and Algebraic Methods in Programming* 85 (4), pp. 540-573. ISSN 2352-2208.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/13368/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

Relational Lattices

Tadeusz Litak^{a,*}, Szabolcs Mikulás^b, Jan Hidders^c

^a *Informatik 8, Friedrich-Alexander-Universität Erlangen-Nürnberg
Martensstraße 3, 91058 Erlangen, Germany*

^b *School of Computer Science and Information Systems,
Birkbeck, University of London, WC1E 7HX London, UK*

^c *Delft University of Technology, Elektrotechn., Wisk. and Inform.,
Mekelweg 4, 2628CD Delft, The Netherlands*

Abstract

Relational lattices are obtained by interpreting lattice connectives as *natural join* and *inner union* between database relations. Our study of their equational theory reveals that the variety generated by relational lattices has not been discussed in the existing literature. Furthermore, we show that addition of just the *header constant* to the lattice signature leads to undecidability of the quasiequational theory. Nevertheless, we also demonstrate that relational lattices are not as intangible as one may fear: for example, they do form a pseudoelementary class. We also apply the tools of Formal Concept Analysis and investigate the structure of relational lattices via their standard contexts. Furthermore, we show that the addition of typing rules and *singleton constants* allows a direct comparison with *monotonic relational expressions* of Sagiv and Yannakakis.

Keywords: relational lattices, relational algebra, database theory, algebraic logic, lattice theory

1. Introduction

We study a class of lattices with a natural database interpretation proposed by Vadim Tropashko [34, 29, 33]. It does not seem to have attracted the attention of algebraists, even those investigating the connections between algebraic logic and relational databases (see, e.g., Imieliński and Lipski [13] or Duntsch and Mikulás [6]).

The connective *natural join* (which we will interpret as lattice meet!) is one of the basic operations of Codd’s (*named*) *relational algebra* [1, 4]. Incidentally, it is also one of its total operations—i.e., defined for all arguments. In general, Codd’s “algebra” is only a *partial algebra*: some operations are defined only between relations with suitable headers, e.g., the (set) union or the difference operator. Apart from the issues of mathematical elegance and generality, this partial nature of operations has also unpleasant practical consequences. For example, queries which do not observe constraints on headers can *crash* [35].

It turns out, however, that it is possible to generalize the union operation to *inner union* defined on all elements of the algebra and lattice-dual to natural join. This approach appears more natural and has several advantages over the embedding of relational “algebras” in cylindric algebras proposed in [13]. For example, we avoid an artificial uniformization of headers and hence queries formed with the use of proposed connectives enjoy the *domain independence property* [36], [1, Ch. 5]. We discuss d.i.p. and related properties formally in Section 2.1 below.

We focus here on the (quasi)equational theory of natural join and inner union. Apart from an obvious mathematical interest, Birkhoff-style equational inference is the basis for certain query optimization

*Corresponding author

Email addresses: szabolcs@dcs.bbk.ac.uk (Szabolcs Mikulás), A.J.H.Hidders@tudelft.nl (Jan Hidders)

URL: <https://www8.informatik.uni-erlangen.de/~litak> (Tadeusz Litak)

techniques where algebraic expressions represent query evaluation plans and are rewritten by the optimizer into equivalent but more efficient expressions. As for *quasiequations*, i.e., definite Horn clauses over equalities, reasoning over many database constraints such as key constraints and foreign keys can be reduced to quasiequational reasoning. Note that an optimizer can consider more equivalent alternatives for the original expression if it can take the specified database constraints into account.

Strikingly, it turned out that relational lattices does not seem to fit anywhere into the rather well-investigated landscape of equational theories of lattices [15, 16]; we will discuss this in detail in Section 3 below. Nevertheless, there were some indications that the considered choice of connectives may lead to positive results concerning decidability/axiomatizability.

On the database side, expressions of our formalisms are closely related to (*unions*) of *conjunctive queries* [1, Ch. 4], [3] and even more so to *monotonic relational expressions* of Sagiv and Yannakakis [28]; the relationship with these classes will be discussed in more detail in Section 6 below. Such classes of queries enjoy decision procedures for problems of containment and equivalence based on so-called Homomorphism Theorem [3, 28], [1, Ch. 6]. In fact, Johnson and Klug [17] show that even in presence of *inclusion dependencies*, the containment problem for conjunctive queries remains in NP when infinite database instances are allowed—and presence of inclusion dependencies gives the containment problem distinctly quasi-equational character.

Another reason for our initial optimism came from algebraic logic itself: a somewhat (unjustly!) forgotten book of Craig [5] showed that the *finitization problem* of algebraic logic allows a positive solution when relations are allowed to contain tuples of varying arity. Note that Craig’s setting was even more liberal than our present one: while we do happily allow relations with differing headers, we assume that all tuples within one relation are defined on a fixed set of attributes.

To our surprise, however, it turned out that — at least when it comes to decidability — expansions of relational lattices share the curse of “untamed” structures from algebraic logic such as Tarski’s relation algebras or cylindric algebras. As soon as an additional *header constant* H is added to the language, one can encode the word problem for semigroups in the quasiequational theory using a technique introduced by Maddux [21]. This means that decidability of query equivalence under constraints for restricted positive database languages does not translate into decidability of corresponding quasiequational theories. However, our Theorem 4.7 and Corollary 4.8 do not rule out possible finite axiomatization results (except for quasiequational theory of *finite* structures) or decidability of equational theory.¹ And with H removed, i.e., in the pure lattice signature, the picture is completely open. Of course, such a language would be rather weak from a database point of view, but natural for an algebraist.

In the final analysis, the difference between (expansions of) relational lattices and settings allowing positive results boils down to *typed* vs. *untyped* (which should not be confused with *named* vs. *unnamed*!) and also possibly *equational* vs. *quasiequational*. Regarding the first of those distinctions, see Section 6 and in particular Section 6.2: the addition of typing discipline to an expansion of our signature allows a direct comparison with *monotonic relational expressions* of Sagiv and Yannakakis [28, Sec. 2.2], a class of queries with perfectly tractable containment and equivalence problems.

We also obtained a number of positive results. First of all, concrete relational lattices are pseudoelementary and hence their closure under subalgebras and products is a quasivariety—Theorem 4.1 and Corollary 4.3. The proof yields an encoding into a sufficiently rich (many-sorted) first-order theory with finitely many axioms. This opens up the possibility of using generic proof assistants like Isabelle or Coq in future investigations—so far, we have only used Prover9/Mace4 to study interderivability of interesting (quasi)equations.² We have also used the tools of Formal Concept Analysis (Theorem 5.3) to investigate the dual structure of full concrete relational lattices and establish, e.g., their subdirect irreducibility (Corollary 5.4). Theorem 5.3 is likely to have further applications—see the discussion of Problem 7.1.

The structure of the paper is as follows. In Section 2.1, we provide basic definitions, including the notion of *domain independence* and its natural strengthening *strict independence* (which does not seem

¹Note, however, that an extension of our signature to a language with EDPC or a discriminator term would result in an undecidable *equational* theory.

²It is worth mentioning that the database inventor of relational lattices has in the meantime developed a dedicated tool [34].

to have been explicitly defined before). In Section 2.2, we establish that relational lattices are indeed lattices and in Section 2.3, we note in passing a potential connection with category theory. Section 3 reports our findings about the (quasi)equational theory of relational lattices: the failure of most standard properties such as weakening of distributivity (Theorem 3.2), those surprising equations and properties that still hold (Theorem 3.5) and dependencies between them (Theorem 3.4). In Section 4, we focus on quasiequations and prove some of most interesting results discussed above, both positive (Theorem 4.1 and Corollaries 4.2–4.4) and negative ones (Theorem 4.7 and Corollaries 4.8–4.9). Section 5 analyzes *standard contexts, incidence and arrow relations* [8] of relational lattices. Section 6 discusses possible extensions of the signature leading towards *expressive completeness* and addition of typing information, which in turn allows a direct comparison with the setting of (monotonic) relational expressions. Section 7 concludes and discusses future work.

This paper is a significantly extended and rewritten version of an earlier conference version [20]. Some of numerous changes introduced are:

- A new version of Section 6, including in particular new Subsection 6.2, providing a direction with fragments of relational algebra, in particular with monotonic relational expressions of Sagiv and Yannakakis [28, Sec. 2.2].
- Theorem 3.5, whose lack was in fact a significant omission.
- Rewritten and extended Theorem 3.4, including derivations of new (quasi-)equations and some dependencies we failed to realize in the new version.
- All the missing proofs, for some clauses of Theorem 3.4 including long Prover9 derivations in the appendix.
- An explicit discussion of domain independence in Section 2.1 (again, its lack in the previous version was an omission) including a new, natural notion *strict independence* which for some reason did not seem to appear in the existing literature, yet for us provides an important criterion by which to judge possible extensions of the language.
- Completely rewritten Section 2.3.
- Added discussion of relationship with numerous references (scattered throughout the text), resulting in a much more extensive bibliography.

2. Basic Definitions

2.1. Domains, Relations and Independence

Let \mathcal{A} be a set of *attribute names* and \mathcal{D} be a set of *domain values*. For $H \subseteq \mathcal{A}$, a *H-sequence from \mathcal{D}* or an *H-tuple over \mathcal{D}* is a function $x : H \rightarrow \mathcal{D}$, i.e., an element of ${}^H\mathcal{D}$. H is called the *header* of x and denoted as $h(x)$. The *restriction of x to H'* is defined as $x[H'] := \{(a, v) \in x \mid a \in H'\}$, in particular $x[H'] = \emptyset$ if $H' \cap h(x) = \emptyset$. We generalize this to the *projection of a set of H-sequences X to a header H'* which is $X[H'] := \{x[H'] \mid x \in X\}$.

A *relation* is a pair $r = (H_r, B_r)$, where $H_r \subseteq \mathcal{A}$ is the *header* of r and $B_r \subseteq {}^{H_r}\mathcal{D}$ the *body* of r . The collection of all relations over \mathcal{D} whose headers are contained in \mathcal{A} will be denoted as $R(\mathcal{D}, \mathcal{A})$. Define the proper class $\mathcal{F} := \{R(\mathcal{D}, \mathcal{A}) \mid \mathcal{D}, \mathcal{A} \in \text{Set}\}$. For a fixed $\mathcal{A} \in \text{Set}$, it is also convenient to isolate the subclass of \mathcal{F} determined by it, i.e., $\mathcal{F}_{\mathcal{A}} := \{R(\mathcal{D}, \mathcal{A}) \mid \mathcal{D} \in \text{Set}\}$; we have thus $\mathcal{F} = \bigcup_{\mathcal{A} \in \text{Set}} \mathcal{F}_{\mathcal{A}}$. A (*n-ary*) *relational query* is a *n-ary* operation ϕ defined on all members of \mathcal{F} :

$$R(\mathcal{D}, \mathcal{A})^n \ni (r_1, \dots, r_n) \mapsto \phi^{\mathcal{D}, \mathcal{A}}(r_1, \dots, r_n) \in R(\mathcal{D}, \mathcal{A}).$$

We say that a query ϕ is *domain independent* [36], [1, Ch. 5] if for all $\mathcal{D}, \mathcal{D}', \mathcal{A}$, it holds that $\phi^{\mathcal{D}, \mathcal{A}}(r_1, \dots, r_n) = \phi^{\mathcal{D}', \mathcal{A}}(r_1, \dots, r_n)$ whenever $r_i \in R(\mathcal{D}_i, \mathcal{A}) \cap R(\mathcal{D}'_i, \mathcal{A})$ ($i \in \{1, \dots, n\}$).




Figure 1: Natural join and inner union. In this example, $\mathcal{A} = \{a, b, c\}$, $D = \{1, 2, 3, 4\}$.

For the purpose of the discussion in Section 6, it is also convenient to define explicitly a stronger property, which appears to be taken for granted in references like [1, Ch. 5]. Namely, say that a query ϕ is *strictly independent* if for all $\mathcal{D}, \mathcal{D}', \mathcal{A}, \mathcal{A}'$, it holds that $\phi^{\mathcal{D}, \mathcal{A}}(r_1, \dots, r_n) = \phi^{\mathcal{D}', \mathcal{A}'}(r_1, \dots, r_n)$ whenever $r_i \in R(\mathcal{D}_i, \mathcal{A}) \cap R(\mathcal{D}'_i, \mathcal{A}')$ ($i \in \{1, \dots, n\}$). That is, the outcome of ϕ is not only independent of irrelevant domain elements, but also of irrelevant attributes. In most of the paper, the operations under consideration are strictly independent (Lemma 2.2 below); only in Section 6.3 we will see examples of domain independent queries which are not strictly independent.

Examples of queries which do not have even the weaker property of domain independence abound in any references discussing explicitly the difference between first-order calculus and relational algebra (which is domain-independent by design), see Abiteboul et al. [1, Ch. 5] for references. Typical examples involve unrestricted negation or universal quantification. This is not a trivial property from the point of view of first-order logic: Vardi [36] shows that for first-order queries, the property of being domain-independent is undecidable.

2.2. Introducing Relational Lattices

For the relations r, s , we define the *natural join* $r \bowtie s$, and *inner union* $r \oplus s$:

$$\begin{aligned} r \bowtie s &:= (H_r \cup H_s, \{x \in {}^{H_r \cup H_s} \mathcal{D} \mid x[H_r] \in B_r \text{ and } x[H_s] \in B_s\}) \\ r \oplus s &:= (H_r \cap H_s, \{x \in {}^{H_r \cap H_s} \mathcal{D} \mid x \in B_r[H_s] \text{ or } x \in B_s[H_r]\}) \end{aligned}$$

In our notation, \bowtie always binds stronger than \oplus . The *header constant* $H := (\emptyset, \emptyset)$ plays a special role: for any r , $(H_r, B_r) \bowtie H = (H_r, \emptyset)$ and hence r_1 and r_2 have the same headers iff $H \bowtie r_1 = H \bowtie r_2$. Note also that the projection of r_1 to H_{r_2} can be defined as $r_1 \oplus (H \bowtie r_2)$. In fact, we can identify $H \bowtie r$ and H_r . We denote $(R(\mathcal{D}, \mathcal{A}), \bowtie, \oplus, H)$ as $\mathfrak{R}^H(\mathcal{D}, \mathcal{A})$, with \mathcal{L}_H denoting the corresponding algebraic signature. $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ is its reduct to the signature $\mathcal{L} := \{\bowtie, \oplus\}$.

Lemma 2.1. *For any \mathcal{D} and \mathcal{A} , $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ is a lattice.*

Proof. This result is due to Tropashko [34, 29, 33], but let us provide an alternative proof. Define $Dom := \mathcal{A} \cup {}^{\mathcal{A}} \mathcal{D}$ and for any $X \subseteq Dom$ set

$$Cl(X) := X \cup \{x \in {}^{\mathcal{A}} \mathcal{D} \mid \exists y \in (X \cap {}^{\mathcal{A}} \mathcal{D}). x[\mathcal{A} - X] = y[\mathcal{A} - X]\}.$$

In other words, $Cl(X)$ is the sum of $X \cap \mathcal{A}$ (the set of attributes contained in X) with the cylindrification of $X \cap {}^{\mathcal{A}} \mathcal{D}$ along the axes in $X \cap \mathcal{A}$. It is straightforward to verify Cl is a closure operator and hence Cl -closed sets form a lattice, with the order being obviously \subseteq inherited from the powerset of Dom . It remains to observe $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ is isomorphic to this lattice and the isomorphism is given by

$$(H, B) \mapsto (\mathcal{A} - H) \cup \{x \in {}^{\mathcal{A}} \mathcal{D} \mid x[H] \in B\}. \quad \square$$

We call $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ the (full) relational lattice over $(\mathcal{D}, \mathcal{A})$. We also use the alternative name *Tropashko lattices* to honor the inventor of these structures. The lattice order given by \bowtie and \oplus is

$$(H_r, B_r) \sqsubseteq (H_s, B_s) \text{ iff } H_s \subseteq H_r \text{ and } B_r[H_s] \subseteq B_s.$$

For classes of algebras, we use $\mathbb{H}, \mathbb{S}, \mathbb{P}$ to denote closures under, respectively, homomorphisms, (isomorphic copies of) subalgebras and products. Let

$$\mathcal{R}_{\text{fin}}^{\mathbb{H}} := \mathbb{S}\{\mathfrak{R}^{\mathbb{H}}(\mathcal{D}, \mathcal{A}) \mid \mathcal{D}, \mathcal{A} \text{ finite}\}, \quad \mathcal{R}_{\text{unr}}^{\mathbb{H}} := \mathbb{S}\{\mathfrak{R}^{\mathbb{H}}(\mathcal{D}, \mathcal{A}) \mid \mathcal{D}, \mathcal{A} \text{ unrestricted}\}$$

and let \mathcal{R}_{fin} and \mathcal{R}_{unr} denote the lattice reducts of respective classes.

Lemma 2.2. *All the operations in the signature $\mathcal{L}_{\mathbb{H}}$ are strictly independent.*

Proof. Straightforward. □

2.3. Relational Lattices, (Op-)Fibrations and the Grothendieck Construction

The reader disinterested in category theory can skip this section without loss of continuity. Given \mathcal{D} and \mathcal{A} , a category theorist may note that $H_{(\cdot)}$, i.e., the mapping sending every relation $r = (H_r, B_r)$ to its header H_r is a (*Grothendieck*) *opfibration* from $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ ordered by \sqsubseteq to $\mathcal{P}^{\supseteq}(\mathcal{A})$, the latter being of course the poset with reverse inclusion order. As we are talking posets here, the action of $H_{(\cdot)}$ on arrows and its functoriality are obvious. However, as most standard references in category theory (see in particular Jacobs [14, Ch. 1]) introduce opfibrations well after fibrations, it is easier to pattern-match all results and notions without having to reverse arrows all the time. So we recall that opfibration $E \rightarrow D$ is just a fibration $E^{op} \rightarrow D^{op}$ and therefore our observation can be reformulated as $H_{(\cdot)}$ being a (*Grothendieck*) *fibration* of $\mathfrak{R}^{\supseteq}(\mathcal{D}, \mathcal{A})$ over $\mathcal{P}^{\supseteq}(\mathcal{A})$, where $\mathfrak{R}^{\supseteq}(\mathcal{D}, \mathcal{A})$ is $\mathfrak{R}(\mathcal{D}, \mathcal{A})$, but ordered by \supseteq rather than \sqsubseteq . The crucial thing to note is that an arrow $r' \supseteq r$ is “cartesian” [14, Def. 1.1.3] iff $B_{r'} = B_r[H_{r'}]$, i.e., if r' is the projection of r to the header of r' . With this, one can note that $H_{(\cdot)}$ is in fact a *split fibration* [14, Def. 1.4.3].

It is thus most natural to view relational lattices themselves as obtained via the so-called Grothendieck construction [14, Def. 1.10] associated with this particular fibration. This construction is obtained via the *quasifunctor* or *pseudo-functor* [14, Def. 1.4.4] defined as follows

$$F_{\mathcal{D}}^{\mathcal{A}} : \mathcal{P}^{\supseteq}(\mathcal{A}) \ni H \longrightarrow \mathcal{P}^{\supseteq}(H\mathcal{D}) \in \text{Pos}$$

$$F_{\mathcal{D}}^{\mathcal{A}}(H \subseteq H') := (H\mathcal{D} \supseteq B \mapsto B[H'] \subseteq H'\mathcal{D})$$

We get then that $\mathfrak{R}^{\supseteq}(\mathcal{D}, \mathcal{A})$ is the Grothendieck completion $\int_{\mathcal{P}^{\supseteq}(\mathcal{A})} F_{\mathcal{D}}^{\mathcal{A}}$.

Curiously enough, a number of recent references mentioned (op-)fibrations and the Grothendieck construction in the database context [19, 18, 30]. The focus and the use seems somehow different: that connection arose in the study of queries, views and RDF triples, but it would be interesting to connect it with the Grothendieck perspective on relational lattices sketched above. Our personal belief is there is even closer relationship with a categorical approach to relational databases proposed recently by Abramsky [2], which moreover yields a surprising connection with Bell’s Theorem from theoretical physics. Our belief is motivated by the central role played by \ast in Abramsky’s work [2, Sec. 2.2] and other similarities. It is worth noting that Abramsky [2, Sec. 3] suggests that this categorical approach may yield a natural connection with (and unifying perspective on) probabilistic databases and provenance semirings.

3. Towards the Equational Theory of Relational Lattices

Let us begin the section with

Open Problem 3.1. *Are $\text{SP}(\mathcal{R}_{\text{unr}}^{\mathbb{H}}) = \text{HSPP}(\mathcal{R}_{\text{unr}}^{\mathbb{H}})$ and $\text{SP}(\mathcal{R}_{\text{unr}}) = \text{HSPP}(\mathcal{R}_{\text{unr}})$?*

If the answer is “no”, it would mean that relational lattices should be considered a quasiequational rather than equational class (cf. Corollary 4.3 below). Note also that the decidability of equational theories seems of less importance from a database point of view than decidability of quasiequational theories. Nevertheless, relating to already investigated varieties of lattices seems a good first step. It turns out that weak forms of distributivity and similar properties studied in standard references [15, 16, 32] tend to fail dramatically:

Theorem 3.2. \mathcal{R}_{fin} (and hence \mathcal{R}_{unr}) does not have any of the following properties (see the above references or the proof below for definitions):

1. upper- and lower-semidistributivity,
2. almost distributivity and neardistributivity,
3. upper- or lower-semimodularity (and hence also modularity),
4. local distributivity/local modularity,
5. the Jordan–Dedekind chain condition,
6. supersolvability.

Proof. For most clauses, it is enough to observe that $\mathfrak{R}(\{0, 1\}, \{0\})$ is isomorphic to L_4 , one of the covers of the non-modular lattice N_5 [23, 16]: a routine counterexample in such cases. In more detail:

Clause 1: Recall that *semidistributivity* is the property:

$$a \oplus b = a \oplus c \text{ implies } a \oplus b = a \oplus (b \star c).$$

Now take a to be \mathbf{H} and b and c to be the atoms with the header $\{0\}$.

Clause 2: This is a corollary of Clause 1 [15, Th 4.2 and Sec 4.3].

Clause 3: Recall that *semimodularity* is the property:

if $a \star b$ covers a and b , then $a \oplus b$ is covered by a and b .

Again, take a to be \mathbf{H} and b to be either of the atoms with the header $\{0\}$.

Clause 4: This is a corollary of Clause 3 [22].

Clause 5: Recall that *the Jordan-Dedekind chain condition* is the property that the cardinalities of two maximal chains between common end points are equal. This obviously fails in N_5 .

Clause 6: Recall that for finite lattices, *supersolvability* [31] boils down to the existence of a maximal chain generating a distributive lattice with any other chain. Again, this fails in N_5 . \square

Remark 3.3. *Theorem 3.2 has an additional consequence regarding the notion called rather misleadingly boundedness in most standard references (see e.g., Jipsen and Rose [15, p. 27]): being an image of a freely generated lattice by a bounded morphism. We use the term McKenzie-bounded, as McKenzie showed that for finite subdirectly irreducible lattices, this property amounts to splitting the lattice of varieties of lattices [15, Theorem 2.25]. Finite Tropashko lattices are subdirectly irreducible (Corollary 5.4 below) but Clause 1 of Theorem 3.2 entails they are not McKenzie-bounded [15, Lemma 2.30].*

Nevertheless, Tropashko lattices do not generate the variety of all lattices. The results of our investigations so far on valid (quasi)equations are summarized in the remainder of this section. First, let us note the following dependencies between equations and quasiequations in Table 1:

Theorem 3.4. *Assuming all lattice axioms, the following statements hold:*

1. Axioms of $\underline{R}^{\mathbf{H}}$ in Table 1 are mutually independent. Similarly, axioms of \underline{R} are mutually independent.
2. $AxRL1$ forces $Qu1$ [25].
3. $Eq1$ implies $Eq2$ and $Eq3$.
4. $Qu2$ together with $Eq1$ imply both $AxRL1$ and $AxRL2$.
5. $Eq1$ is implied by $AxRH1$. The converse implication does not hold even in presence of $AxRL1$.
6. $AxRH1$ and $AxRH2$ jointly imply $Qu2$, although each of the two equations separately is too weak to entail $Qu2$. In the converse direction, $Qu2$ implies $AxRH2$ but not $AxRH1$.
7. $AxRH1$ and $AxRH2$ jointly imply $Qu3$, although each of the two equations separately is too weak to entail $Qu3$ (in the case of $AxRH2$ even in presence of $Eq1$).
8. $AxRH1$ implies $Eq4$.

Proof. Clause 1: For mutual independence of the two axioms of $\underline{R}^{\mathbf{H}}$, counterexamples can be obtained by appropriate choices of the interpretation of \mathbf{H} in the pentagon lattice. As for \underline{R} , the example showing that the validity of $AxRL2$ does not imply the validity of $AxRL1$ is the non-distributive diamond lattice M_3 , while the reverse implication can be disproved with an eight-element model:

Table 1: (Quasi)equations Valid in Tropashko Lattices

Class \underline{R}^H in the signature \mathcal{L}_H :

all lattice axioms

$$\text{AxRH1} \quad H \times x \times (y \oplus z) \oplus y \times z = (H \times x \times y \oplus z) \times (H \times x \times z \oplus y)$$

$$\text{AxRH2} \quad x \times (y \oplus z) = x \times (z \oplus H \times y) \oplus x \times (y \oplus H \times z)$$

Class \underline{R} in the signature \mathcal{L} (without H):

all lattice axioms

$$\text{AxRL1} \quad x \times y \oplus x \times z = x \times (y \times (x \oplus z) \oplus z \times (x \oplus y))$$

$$\begin{aligned} \text{AxRL2} \quad t \times ((x \oplus y) \times (x \oplus z) \oplus (u \oplus w) \times (u \oplus v)) = \\ = t \times ((x \oplus y) \times (x \oplus z) \oplus u \oplus w \times v) \oplus t \times ((u \oplus w) \times (u \oplus v) \oplus x \oplus y \times z) \end{aligned}$$

(in \mathcal{L}_H , AxRL1 and AxRL2 are derivable from AxRH1 and AxRH2, see Theorem 3.4)

Additional (quasi)equations derivable in \underline{R}^H and \underline{R} :

$$\text{Qu1} \quad x \oplus y = x \oplus z \Rightarrow x \times (y \oplus z) = x \times y \oplus x \times z$$

$$\text{Qu2} \quad H \times (x \oplus y) = H \times (x \oplus z) \Rightarrow x \times (y \oplus z) = x \times y \oplus x \times z$$


$$\text{Qu3} \quad H \times (x \oplus y) = H \times (x \oplus z) = H \times (y \oplus z) \Rightarrow x \oplus y \times z = (x \oplus y) \times (x \oplus z)$$

$$\text{Eq1} \quad H \times x \times (y \oplus z) = H \times x \times y \oplus H \times x \times z$$

$$\text{Eq2} \quad H \times (y \oplus z) = H \times y \oplus H \times z$$

$$\text{Eq3} \quad H \times t \times (x \oplus y) \times (x \oplus z) = H \times t \times x \oplus H \times t \times y \times z$$

$$\text{Eq4} \quad H \times x \oplus x \times y = x \times (y \oplus H \times x)$$



Clause 3: Eq2 is obvious by lattice laws: substitute $y \oplus z$ for x and use absorption. For Eq3, we reason as follows:

$$\begin{aligned}
H \times t \times (x \oplus y) \times (x \oplus z) &= (H \times t \times x \oplus H \times t \times y) \times (H \times t \times x \oplus H \times t \times z) \\
&= H \times (t \times x \oplus t \times y) \times (t \times x \oplus t \times z) \\
&= H \times (t \times x \oplus t \times y) \times t \times x \oplus H \times (t \times x \oplus t \times y) \times t \times z \\
&= H \times t \times x \oplus H \times (t \times x \oplus t \times y) \times t \times z \\
&= H \times t \times x \oplus H \times t \times x \times z \oplus H \times t \times y \times z \\
&= H \times t \times x \oplus H \times t \times y \times z
\end{aligned}$$

Clause 4: Direct computation. In more detail: for AxRL1, substitute $y \times (x \oplus z)$ for y and $z \times (x \oplus y)$ for z in the antecedent of Qu2. We get then the consequent of Qu2, as $H \times (x \oplus y \times (x \oplus z)) = H \times x \oplus H \times y \times (x \oplus z) = H \times x \oplus H \times y \times x \oplus H \times y \times z = H \times x \oplus H \times y \times z = H \times x \oplus H \times x \times z \oplus H \times y \times z = H \times (x \oplus z \times (x \oplus y))$ (we are obviously using Eq1 here). Thus, the right side of AxRL1 is equal to $x \times y \times (x \oplus z) \oplus x \times z \times (x \oplus y)$. But this, by the absorption law, is equal to $x \times y \oplus x \times z$, i.e., the left side of AxRL1.

For the seemingly monstrous AxRL2, the trick is similar. Consider

$$\begin{aligned}
H \times ((x \oplus y) \times (x \oplus z) \oplus u \oplus w \times v) &= H \times (x \oplus y) \times (x \oplus z) \oplus H \times u \oplus H \times w \times v \\
&= H \times x \oplus H \times y \times z \oplus H \times (u \oplus w) \times (u \oplus v) \\
&= H \times (x \oplus y \times z) \oplus H \times (u \oplus w) \times (u \oplus v)
\end{aligned}$$

This allows us to use Qu2 to rewrite the right side of AxRL2:

$$\begin{aligned}
t \times ((x \oplus y) \times (x \oplus z) \oplus u \oplus w \times v) \oplus t \times ((u \oplus w) \times (u \oplus v) \oplus x \oplus y \times z) \\
&= t \times ((x \oplus y) \times (x \oplus z) \oplus u \oplus w \times v \oplus (u \oplus w) \times (u \oplus v) \oplus x \oplus y \times z) \\
&= t \times ((x \oplus y) \times (x \oplus z) \oplus (u \oplus w) \times (u \oplus v))
\end{aligned}$$

(the second equality obtaining by lattice laws).

Clause 5: The first part has been proved with the help of Prover9 (66 lines of proof—see Appendix). The counterexample for the converse has been found by Mace4: it is obtained by choosing H to be the top element of the pentagon lattice.

Clause 6: Prover9 was able to prove the first statement both in presence and in absence of AxRL1, although there was a significant difference in the length of both proofs (38 lines vs. 195 lines—see Appendix). The implication from Qu2 to AxRH2 is straightforward. All the necessary counterexamples have been found by Mace4 by appropriate choices of the interpretation of H in the pentagon lattice.

Clause 7: The positive statement was proved by Prover9 (mere 196 lines—see Appendix). Again, counterexamples for all the negative statements can be found using 5-element models.

Clause 8: Substitute x for z and use the absorption law. \square

AxRL1 comes from Padmanabhan et al. [25] as an example of an equation which forces *the Huntington property* (distributivity under unique complementation). Qu1 is a form of weak distributivity, denoted as CD_{\vee} [25] or WD_{\wedge} [16].

Theorem 3.5.

- $AxRH1$ and $AxRH2$ are valid in $\mathcal{R}_{\text{unr}}^H$ (and consequently in $\mathcal{R}_{\text{fin}}^H$).
- $AxRH1$, $AxRH2$ and $Eq1$ are valid in $\mathcal{R}_{\text{unr}}^H$ (and consequently in $\mathcal{R}_{\text{fin}}^H$).
- Axioms of $\mathcal{R}_{\text{unr}}^H$ are valid in $\mathcal{R}_{\text{unr}}^H$ (and consequently in $\mathcal{R}_{\text{fin}}^H$). Similarly, axioms of \underline{R} are valid in \mathcal{R}_{unr} (and consequently \mathcal{R}_{fin}).
- All formulas in Table 1 are valid in $\mathcal{R}_{\text{unr}}^H$ (and consequently in $\mathcal{R}_{\text{fin}}^H$). Those not involving H are valid in \mathcal{R}_{unr} (and consequently in \mathcal{R}_{fin}).

Proof. Theorem 3.4 implies that all clauses are equivalent, so we can choose whichever we want to prove. Perhaps the most convenient is the second one. Validity of $Eq1$ is immediate, as the sublattice of relations with empty body (we can call it the *header sublattice*) is obviously distributive. In presence of $Eq1$, we obtain automatically

$$H \times x \times (y \oplus z) \oplus y \times z \leq (H \times x \times y \oplus z) \times (H \times x \times z \oplus y),$$

so to establish $AxRH1$, it is enough to establish the other inequality. Denote $H_{x \times y} \cap H_z$ as H_1 and $H_{x \times z} \cap H_y$ as H_2 ; note that both $H_1 \cap H_2 = H_y \cap H_z$. A tuple t belongs to the body of $(H \times x \times y \oplus z) \times (H \times x \times z \oplus y)$ iff there exists $t_1 \in B_z[H_1]$ and $t_2 \in B_z[H_2]$ s.t. $t_1[H_1 \cap H_2] = t_2[H_1 \cap H_2]$ and t is the concatenation of t_1 and t_2 . This is in turn is equivalent to the existence of $t'_1 \in B_z$ and $t'_2 \in B_y$ s.t. $t'_1[H_1 \cap H_2] = t'_2[H_1 \cap H_2]$ (t_1 being $t'_1[H_1]$ and $t_2 = t'_2[H_2]$); by our earlier observation, $H_1 \cap H_2$ is precisely the set of attributes on which headers of t'_1 and t'_2 overlap, so we can see t as restriction of concatenation of t'_1 and t'_2 to $H_1 \cup H_2$. But this means that t belongs to the body of $H \times x \times (y \oplus z) \oplus y \times z$.

Finally, let us consider $AxRH2$. Lattice laws yield that

$$x \times (y \oplus z) \geq x \times (z \oplus H \times y) \oplus x \times (y \oplus H \times z),$$

so we only need to establish the opposite inequality. Pick any t in the body of $x \times (y \oplus z)$. Clearly, there exist $t_x \in B_x$ and $t_2 \in B_{y \oplus z}$ overlapping on $H_x \cap H_y \cap H_z$ and t is their concatenation. Now, t_2 is either a restriction of some $t_y \in B_y$ or of some $t_z \in B_z$. Assume the first case; we get then that t_2 belongs to the body of $y \oplus H \times z$ and consequently $t \in x \times (y \oplus H \times z)$. Similarly, in the other case we get that $t \in x \times (z \oplus H \times y)$. \square

Open Problem 3.6. *Are the equational theories of $\mathcal{R}_{\text{unr}}^H$ (\mathcal{R}_{unr}) and $\mathcal{R}_{\text{fin}}^H$ (\mathcal{R}_{fin}) equal? How about quasiequational ones?*

Open Problem 3.7. *Is the equational theory of $\mathcal{R}_{\text{unr}}^H$ (\mathcal{R}_{unr}) equal to \underline{R}^H (\underline{R} , respectively)? If not, is it finitely axiomatizable at all?*

If the answer to the last question is in the negative, one can perhaps attempt a rainbow-style argument from algebraic logic [12].

4. Relational Lattices as a Quasiequational Class

In the introduction, we discussed why an axiomatization of valid *quasiequations* is desirable from a DB point of view. There is also an algebraic reason: the class of representable Tropashko lattices (i.e., the SP-closure of concrete ones) is a *quasivariety*. This is a corollary of a more powerful result; recall that being *pseudoelementary* means being a reduct of an elementary class in a richer (possibly multi-sorted) language and that this notion plays a central role in algebraic study of axiomatizability and representability [12]:

Theorem 4.1. *$\mathcal{R}_{\text{unr}}^H$ and \mathcal{R}_{unr} are pseudoelementary classes.*

Proof. (sketch) Assume a language with sorts A, F, D and R . The connectives of \mathcal{L}_H live in R , we also have a relation symbol $inR : (F \cup A) \times R$ and a function symbol $assign : (F \times A) \mapsto D$. The interpretation is suggested by the closure system used in the proof of Lemma 2.1. That is, A denotes \mathcal{A} , F denotes ${}^A\mathcal{D}$, D denotes \mathcal{D} and R denotes the family of Cl -closed subsets of Dom . Moreover, $assign(f, a)$ denotes the value of the \mathcal{A} -sequence denoted by f on the attribute a and $inR(x, r)$ —the membership of an attribute/sequence

in the closed subset of Dom denoted by r . One needs to postulate the following axioms: “ F and R are extensional” (the first via axioms of *assign*, the second via axioms on inR); “each element of R is Cl -closed”; “ \times and \oplus are genuine infimum/supremum on R ”. For \mathcal{R}_{unr}^H , we add an axiom “ inR assigns no elements of F and all elements of A (the latter means all attributes are *irrelevant* for the element under consideration!) to H ”. \square

Corollary 4.2. \mathcal{R}_{unr}^H and \mathcal{R}_{unr} are closed under ultraproducts.

Corollary 4.3. The $\mathbb{S}\mathbb{P}$ -closures of \mathcal{R}_{unr}^H and \mathcal{R}_{unr} are quasiequational classes.

Corollary 4.4. The quasiequational, universal and elementary theories of \mathcal{R}_{unr}^H and \mathcal{R}_{unr} are recursively enumerable.

Proof. The proof of Theorem 4.1 uses finitely many axioms. \square

Note that postulating that headers are *finite* subsets of \mathcal{A} would break the proof of Theorem 4.1: such conditions are not first-order. However, concrete database instances always belong to \mathcal{R}_{fin}^H and we will show now that the decidability status of the quasiequational theory of \mathcal{R}_{unr}^H and \mathcal{R}_{fin}^H is the same. Moreover, an undecidability result also obtains for the corresponding abstract class, much like for relation algebras and cylindric algebras—in fact, we build on a proof of Maddux [21] for CA_3 —and we *do not even need all the axioms* of \underline{R}^H to show this! Let $\underline{RH1}$ be the variety of \mathcal{L}_H -algebras axiomatized by the lattice axioms and $AxRH1$. Let us list some basic observations:

Proposition 4.5.

1. $\mathcal{R}_{fin}^H \subset \mathcal{R}_{unr}^H \subset \mathbb{S}\mathbb{P}(\mathcal{R}_{unr}^H) \subseteq \underline{R}^H \subset \underline{RH1}$.
2. *Eq4* holds in $\underline{RH1}$.
3. *AxRH1* holds whenever H is interpreted as the bottom of a bounded lattice.
4. *AxRH1* holds for an arbitrary choice of H in a distributive lattice.

Proof. Clause 2 holds by clause 8 of Theorem 3.4. The remaining ones are straightforward to verify. \square

Note, e.g., that interpreting H as \perp in $AxRH2$ would only work if the lattice is distributive, so Clause 3 would not hold in general for $AxRH2$. In order to state our undecidability result, we need first

Definition 4.6. Let $\bar{e} = (u_0, u_1, u_2, e_0, e_1)$ be an arbitrary 5-tuple of variables. We abbreviate $u_0 \times u_1 \times u_2$ as u . For arbitrary L -terms s, t define

$$\begin{aligned} \mathbf{c}_0^{\bar{e}} \langle t \rangle &:= u \times (H \times u_1 \times u_2 \oplus u \times t), \\ \mathbf{c}_1^{\bar{e}} \langle t \rangle &:= u \times (H \times u_0 \times u_2 \oplus u \times t), \\ \mathbf{c}_2^{\bar{e}} \langle t \rangle &:= u \times (H \times u_0 \times u_1 \oplus u \times t), \\ s \circ^{\bar{e}} t &:= \mathbf{c}_2^{\bar{e}} \langle \mathbf{c}_1^{\bar{e}} \langle e_0 \times \mathbf{c}_2^{\bar{e}} \langle s \rangle \rangle \times \mathbf{c}_0^{\bar{e}} \langle e_1 \times \mathbf{c}_2^{\bar{e}} \langle t \rangle \rangle \rangle. \end{aligned}$$

Let $T_n(x_1, \dots, x_n)$ be the collection of all semigroup terms in n variables. Whenever $\bar{e} = (x_{n+1}, \dots, x_{n+5})$ define the translation $\tau^{\bar{e}}$ of semigroup terms as follows: $\tau^{\bar{e}}(x_i) := x_i$ for $i \leq n$ and $\tau^{\bar{e}}(s \circ t) := s \circ^{\bar{e}} t$ for any $s, t \in T_n(x_1, \dots, x_n)$.

Whenever \bar{e} is clear from the context, we will drop it to ensure readability. Now we can formulate

Theorem 4.7. For any $p_0, \dots, p_m, r_0, \dots, r_m, s, t \in T_n(x_1, \dots, x_n)$, the following conditions are equivalent:

(I) The quasiequation

$$(Qu4) \quad \forall x_1, \dots, x_n. (p_0 = r_0 \ \& \ \dots \ \& \ p_m = r_m \Rightarrow s = t)$$

holds in all semigroups (finite semigroups).

(II) For $\bar{e} = (x_{n+1}, \dots, x_{n+5})$ as in Definition 4.6, the quasiequation

$$(Qu5) \quad \begin{aligned} & \forall x_0, x_1, \dots, x_{n+5}. (\tau^{\bar{e}}(p_0) = \tau^{\bar{e}}(r_0) \& \dots \& \tau^{\bar{e}}(p_m) = \tau^{\bar{e}}(r_m) \& \\ & \& x_{n+4} = \mathbf{c}_0^{\bar{e}} \langle x_{n+4} \rangle \& x_{n+5} = \mathbf{c}_1^{\bar{e}} \langle x_{n+5} \rangle) \Rightarrow \\ & \Rightarrow \tau^{\bar{e}}(s) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle x_0 \rangle = \tau^{\bar{e}}(t) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle x_0 \rangle) \end{aligned}$$

holds in every member of $\mathcal{R}_{\text{unr}}^{\text{H}}$ (every member of $\mathcal{R}_{\text{fin}}^{\text{H}}$).

(III) Qu5 above holds in every member of RH1 (finite member of RH1).

Proof. (I) \Rightarrow (III). By contraposition:

Take any $\mathfrak{A} \in \underline{\text{RH1}}$ and arbitrarily chosen elements $u_0, u_1, u_2 \in \mathfrak{A}$. In order to use Maddux's technique, we have to prove that for any $a, b \in \mathfrak{A}$ and $k, l < 3$

- (b) $\mathbf{c}_k \langle \mathbf{c}_k \langle a \rangle \rangle = \mathbf{c}_k \langle a \rangle$,
- (c) $\mathbf{c}_k \langle a \times \mathbf{c}_k \langle b \rangle \rangle = \mathbf{c}_k \langle a \rangle \times \mathbf{c}_k \langle b \rangle$,
- (d) $\mathbf{c}_k \langle \mathbf{c}_l \langle a \rangle \rangle = \mathbf{c}_l \langle \mathbf{c}_k \langle a \rangle \rangle$

(we deliberately keep the same labels as in the quoted paper), where $\mathbf{c}_k \langle a \rangle$ is defined in the same way as in Definition 4.6 above. We will denote by $u_{\hat{k}}$ the product of u_i 's such that $i \in \{0, 1, 2\} - \{k\}$. For example, $u_{\hat{0}} = u_1 \times u_2$.

For (b):

$$\begin{aligned} L &= u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a)) \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \times (u \oplus \mathbf{H} \times u_{\hat{k}} \oplus u \times a) \oplus u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a)) && \text{by lattice laws} \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \times u \oplus \mathbf{H} \times u_{\hat{k}} \oplus u \times a) \times (\mathbf{H} \times u_{\hat{k}} \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a) \oplus u) && \text{by AxRH1} \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a) \times (\mathbf{H} \times u_{\hat{k}} \oplus u) && \text{by lattice laws} \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a) && \text{by lattice laws} \\ &= R. \end{aligned}$$

(c) is proved using a similar trick:

$$\begin{aligned} L &= u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times b)) \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \times (u \times a \oplus \mathbf{H} \times u_{\hat{k}} \oplus u \times b) \oplus u \times a \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times b)) && \text{by lattice laws} \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \times u \times a \oplus \mathbf{H} \times u_{\hat{k}} \oplus u \times b) \times (\mathbf{H} \times u_{\hat{k}} \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times b) \oplus u \times a) && \text{by AxRH1} \\ &= u \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times b) \times (\mathbf{H} \times u_{\hat{k}} \oplus u \times a) && \text{by lattice laws} \\ &= R. \end{aligned}$$

(d) is obviously true for $k = l$, hence we can restrict attention to $k \neq l$. Let j be the remaining element of $\{0, 1, 2\}$. Thus,

$$\begin{aligned} L &= u \times (\mathbf{H} \times u_l \times u_j \oplus u \times (\mathbf{H} \times u_k \times u_j \oplus u \times a)) \\ &= u \times (\mathbf{H} \times u_l \times u_j \oplus u_l \times (\mathbf{H} \times u_k \times u_j \oplus u \times a)) && \text{by Eq4} \\ &= u \times (\mathbf{H} \times u_l \times u_j \times (u_l \oplus \mathbf{H} \times u_k \times u_j \oplus u \times a) \oplus u_l \times (\mathbf{H} \times u_k \times u_j \oplus u \times a)) && \text{by lattice laws} \\ &= u \times (\mathbf{H} \times u_l \times u_j \oplus \mathbf{H} \times u_k \times u_j \oplus u \times a) \times (\mathbf{H} \times u_l \times u_j \times (\mathbf{H} \times u_k \times u_j \oplus u \times a) \oplus u_l) && \text{by AxRH1} \\ &= u \times (\mathbf{H} \times u_l \times u_j \oplus \mathbf{H} \times u_k \times u_j \oplus u \times a) \times u_l && \text{by lattice laws} \\ &= u \times (\mathbf{H} \times u_l \times u_j \oplus \mathbf{H} \times u_k \times u_j \oplus u \times a) && \text{by lattice laws} \end{aligned}$$

and in the last term, u_l and u_k may be permuted by commutativity. We then obtain the right side of the equation via an analogous sequence of transformations in the reverse direction, with the roles of u_k and u_l replaced.

The rest of the proof mimics the one by Maddux [21]. In some detail: assume there is $\bar{e} = (u_0, u_1, u_2, e_0, e_1) \in \mathfrak{A}$ such that

$$(a) \quad \mathbf{c}_0^{\bar{e}} \langle e_0 \rangle = e_0, \mathbf{c}_1^{\bar{e}} \langle e_1 \rangle = e_1$$

holds. Using (a)–(d) we prove that for every $a, b \in \mathfrak{A}$ the following hold:

- (i) $\mathbf{c}_1^{\bar{e}} \langle a \circ^{\bar{e}} b \rangle = a \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle b \rangle$,
- (ii) $a \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle b \rangle = \mathbf{c}_1^{\bar{e}} \langle \mathbf{c}_2^{\bar{e}} \langle a \rangle \times \mathbf{c}_0^{\bar{e}} \langle \mathbf{c}_2^{\bar{e}} \langle e_0 \times e_1 \times \mathbf{c}_2^{\bar{e}} \langle \mathbf{c}_1^{\bar{e}} \langle b \rangle \rangle \rangle \rangle \rangle$,
- (iii) $(a \circ^{\bar{e}} b) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle c \rangle = a \circ^{\bar{e}} (b \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle c \rangle)$,
- (iv) $((a \circ^{\bar{e}} b) \circ^{\bar{e}} c) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle d \rangle = (a \circ^{\bar{e}} (b \circ^{\bar{e}} c)) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle d \rangle$.

Now pick \mathfrak{A} witnessing the failure of Qu5 together with $\bar{e} = (u_0, u_1, u_2, e_0, e_1)$ such that elements of \bar{e} interpret variables $(x_{n+1}, \dots, x_{n+5})$ in Qu5. This means (a) is satisfied, hence (i)–(iv) hold for every element of \mathfrak{A} . We define an equivalence relation \equiv on \mathfrak{A} :

$$a \equiv b \text{ iff for all } c \in \mathfrak{A}, a \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle c \rangle = b \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle c \rangle.$$

We take $\circ^{\bar{e}}$ to be the semigroup operation on \mathfrak{A}/\equiv . Following Maddux [21], we use (i)–(iv) to prove that this operation is well-defined (i.e., independent of the choice of representatives) and satisfies semigroup axioms. It follows from the assumptions that the semigroup thus defined fails Qu4.

(III) \Rightarrow (II). Immediate.

(II) \Rightarrow (I). In analogy to Maddux [21], given a semigroup $\mathfrak{B} = (B, \circ, \mathbf{u})$ failing Qu4 and a valuation v witnessing this failure, consider $\mathfrak{R}(B, \{0, 1, 2\})$ with a valuation w defined as follows:

$$\begin{aligned} w(x_0) &:= (\{0, 1, 2\}, \{\{(0, v(r)), (1, a), (2, b)\} \mid a, b \in \mathfrak{B}\}), \\ w(x_i) &:= (\{0, 1, 2\}, \{\{(0, a), (1, a \circ v(x_i)), (2, b)\} \mid a, b \in \mathfrak{B}\}), & i \leq n, \\ w(x_{n+i}) &:= (\{i\}, \{\{(i, b)\} \mid b \in \mathfrak{B}\}), & (0 < i \leq 3), \\ w(x_{n+4}) &:= (\{0, 1, 2\}, \{\{(0, a), (1, b), (2, b)\} \mid a, b \in \mathfrak{B}\}), \\ w(x_{n+5}) &:= (\{0, 1, 2\}, \{\{(0, b), (1, a), (2, b)\} \mid a, b \in \mathfrak{B}\}). \end{aligned}$$

It is proved by induction that

$$w(\tau^{\bar{e}}(t)) = (\{0, 1, 2\}, \{\{(0, a), (1, a \circ v(t)), (2, b)\} \mid a, b \in \mathfrak{B}\})$$

(where $e = (x_{n+1}, \dots, x_{n+5})$) for every $t \in T(x_1, \dots, x_n)$ and also

$$\begin{aligned} w(\tau^{\bar{e}}(s) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle x_0 \rangle) &= (\{0, 1, 2\}, \{\{(0, a), (1, b), (2, c)\} \mid a, b, c \in \mathfrak{B}, v(r) \circ a = v(s)\}), \\ w(\tau^{\bar{e}}(r) \circ^{\bar{e}} \mathbf{c}_1^{\bar{e}} \langle x_0 \rangle) &= (\{0, 1, 2\}, \{\{(0, a), (1, b), (2, c)\} \mid a, b, c \in \mathfrak{B}, v(r) \circ a = v(r)\}). \end{aligned}$$

Any tuple whose value for attribute 0 is \mathbf{u} belongs to the first relation, but not to the second. Thus w is a valuation refuting Qu5. \square

Corollary 4.8. *The quasiequational theory of any class of algebras between $\mathcal{R}_{\text{fin}}^{\text{H}}$ and $\underline{RH1}$ is undecidable.*

Proof. Follows from Theorem 4.7 and theorems of Gurevič [9, 10] and Post [26] (for finite and arbitrary semigroups, respectively). \square

Corollary 4.9. *The quasiequational theory of $\mathcal{R}_{\text{fin}}^{\text{H}}$ is not finitely axiomatizable.*

Proof. Follows from Theorem 4.7 and the Harrop criterion [11]. \square

Open Problem 4.10. *Are the quasiequational theories of \mathcal{R}_{umr} and \mathcal{R}_{fin} (i.e., of lattice reducts) decidable?*

5. The Concept Structure of Tropashko Lattices

Given a finite lattice \mathcal{L} with $\mathfrak{J}(\mathcal{L})$ and $\mathfrak{M}(\mathcal{L})$ being the sets of its, respectively, join- and meet-irreducibles, let us follow Formal Concept Analysis [8] and investigate the structure of \mathcal{L} via its *standard context* $\text{con}(\mathcal{L}) := (\mathfrak{J}(\mathcal{L}), \mathfrak{M}(\mathcal{L}), \mathbf{l}_{\leq})$, where $\mathbf{l}_{\leq} := \leq \cap (\mathfrak{J}(\mathcal{L}) \times \mathfrak{M}(\mathcal{L}))$. Set

$$\begin{aligned} g \swarrow m &: g \text{ is } \leq\text{-minimal in } \{h \in \mathfrak{J}(\mathcal{L}) \mid \text{not } h \mathbf{l}_{\leq} m\}, \\ g \nearrow m &: m \text{ is } \leq\text{-maximal in } \{n \in \mathfrak{M}(\mathcal{L}) \mid \text{not } g \mathbf{l}_{\leq} n\}, \\ g \nearrow\swarrow m &: g \swarrow m \ \& \ g \nearrow m. \end{aligned}$$

Let also $\swarrow\swarrow$ be the smallest relation containing \swarrow and satisfying the condition

$$g \swarrow\swarrow m, h \nearrow m \text{ and } h \swarrow n \text{ imply } g \swarrow\swarrow n;$$

in a more compact notation, $\swarrow\swarrow \circ \nearrow \circ \swarrow \subseteq \swarrow\swarrow$. We have the following

Proposition 5.1. [8, Theorem 17] *A finite lattice is*

- *subdirectly irreducible iff there is $m \in \mathfrak{M}(\mathcal{L})$ such that $\swarrow\swarrow \supseteq \mathfrak{J}(\mathcal{L}) \times \{m\}$,*
- *simple iff $\swarrow\swarrow = \mathfrak{J}(\mathcal{L}) \times \mathfrak{M}(\mathcal{L})$.*

Let us describe $\mathfrak{J}(\mathfrak{R}(\mathcal{D}, \mathcal{A}))$ and $\mathfrak{M}(\mathfrak{R}(\mathcal{D}, \mathcal{A}))$ for finite \mathcal{D} and \mathcal{A} . Set

$$\begin{aligned} \mathcal{A}Dom_{\mathcal{D}, \mathcal{A}} &:= \{\text{adom}(x) \mid x \in {}^{\mathcal{A}}\mathcal{D}\} & \text{where } \text{adom}(x) &:= (\mathcal{A}, \{x\}), \\ \mathcal{A}Att_{\mathcal{D}, \mathcal{A}} &:= \{\text{aatt}(a) \mid a \in \mathcal{A}\} & \text{where } \text{aatt}(a) &:= (\mathcal{A} - \{a\}, \emptyset), \\ \text{CoDom}_{\mathcal{D}, H} &:= \{\text{codom}^H(x) \mid x \in {}^H\mathcal{D}\} & \text{where } \text{codom}^H(x) &:= (H, {}^H\mathcal{D} - \{x\}), \\ \text{CoAtt}_{\mathcal{D}, \mathcal{A}} &:= \{\text{coatt}(a) \mid a \in \mathcal{A}\} & \text{where } \text{coatt}(a) &:= (\{a\}, \{a\}\mathcal{D}), \\ \mathcal{J}_{\mathcal{D}, \mathcal{A}} &:= \mathcal{A}Dom_{\mathcal{D}, \mathcal{A}} \cup \mathcal{A}Att_{\mathcal{D}, \mathcal{A}}, \\ \mathcal{M}_{\mathcal{D}, \mathcal{A}} &:= \text{CoAtt}_{\mathcal{D}, \mathcal{A}} \cup \bigcup_{H \subseteq \mathcal{A}} \text{CoDom}_{\mathcal{D}, H}. \end{aligned}$$

It is worth noting that $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ naturally divides into what we may call *boolean H -fibres*—i.e., the powerset algebras of ${}^H\mathcal{D}$ for each $H \subseteq \mathcal{A}$. Furthermore, the projection mapping from H -fibre to H' -fibre where $H' \subseteq H$ is a join-homomorphism. Lastly, note that the bottom elements of H -fibres—i.e., elements of the form (H, \emptyset) —and top elements of the form $(H, {}^H\mathcal{D})$ form two additional boolean slices, which we may call the *lower attribute slice* and the *upper attribute slice*, respectively. Both are obviously isomorphic copies of the powerset algebra of \mathcal{A} . The intention of our definition should be clear then:

- The join-irreducibles are only the atoms of the \mathcal{A} -fibre (i.e., the fibre with the longest tuples) plus the atoms of the lower attribute slice.
- The set of meet-irreducibles is much richer: it consists of the coatoms of *all H -fibres* (note $\mathcal{M}_{\mathcal{D}, \mathcal{A}}$ includes \mathbf{H} as the sole element of $\text{CoDom}_{\mathcal{D}, \emptyset}$) plus all coatoms of the *upper attribute slice*.

Let us formalize these two itemized points as

Theorem 5.2. *For any finite \mathcal{A} and \mathcal{D} such that $|\mathcal{D}| \geq 2$, we have*

$$\begin{aligned} \mathcal{J}_{\mathcal{D}, \mathcal{A}} &= \mathfrak{J}(\mathfrak{R}(\mathcal{D}, \mathcal{A})), & (\text{join-irreducibles}) \\ \mathcal{M}_{\mathcal{D}, \mathcal{A}} &= \mathfrak{M}(\mathfrak{R}(\mathcal{D}, \mathcal{A})). & (\text{meet-irreducibles}) \end{aligned}$$

Proof. (join-irreducibles): To prove the \subseteq -direction, simply observe that the elements of $\mathcal{J}_{\mathcal{D},\mathcal{A}}$ are exactly the atoms of $\mathfrak{R}(\mathcal{D},\mathcal{A})$. For the converse, note that

- every element in a H -fibre is a join of the atoms of this fibre, as each H -fibre has a boolean structure and in the boolean case atomic = atomistic,
- the header elements (H, \emptyset) are joins of elements of $\mathcal{A}Att_{\mathcal{D},\mathcal{A}}$,
- the atoms of H -fibres are joins of header elements with elements of $\mathcal{A}Att_{\mathcal{D},\mathcal{A}}$.

Hence, no element of $\mathfrak{R}(\mathcal{D},\mathcal{A})$ outside $\mathcal{A}Att_{\mathcal{D},\mathcal{A}}$ can be join-irreducible.

(meet-irreducibles): This time, the \supseteq -direction is easier to show: $\mathcal{M}_{\mathcal{D},\mathcal{A}}$ includes the coatoms of H -fibres and of the upper attribute slice. Hence, the basic properties of finite boolean algebras imply all meet-irreducibles must be contained in $\mathcal{M}_{\mathcal{D},\mathcal{A}}$: every element of $\mathfrak{R}(\mathcal{D},\mathcal{A})$ can be obtained as an intersection of elements of $\mathcal{M}_{\mathcal{D},\mathcal{A}}$. For the \subseteq -direction, it is clear that elements of $CoAtt_{\mathcal{D},\mathcal{A}}$ are meet-irreducible, as they are coatoms of the whole $\mathfrak{R}(\mathcal{D},\mathcal{A})$. This also applies to $H \in CoDom_{\mathcal{D},\emptyset}$. Now take $\text{codom}^H(x) = (H, {}^H\mathcal{D} - \{x\})$ for a non-empty $H = \{1, \dots, h\}$ and $x = (x_1, \dots, x_h) \in {}^H\mathcal{D}$ and assume $\text{codom}^H(x) = r \ast s$ for $r, s \neq \text{codom}^H(x)$. That is, $H = H_r \cup H_s$ and

$${}^H\mathcal{D} - \{x\} = \{y \in {}^{H_r \cup H_s}\mathcal{D} \mid y[H_r] \in B_r \text{ and } y[H_s] \in B_s\}.$$

Note that wlog $H_r \subsetneq H$ and $r \subseteq \text{codom}^{H_r}(z)$ for some $z \in {}^{H_r}\mathcal{D}$; otherwise, if both r and s were top elements of their respective fibres, their meet would be $(H, {}^H\mathcal{D})$. Thus

$${}^H\mathcal{D} - \{x\} \subseteq \{y \in {}^H\mathcal{D} \mid y[H_r] \neq z\}$$

and by contraposition

$$\{y \in {}^H\mathcal{D} \mid y[H_r] = z\} \subseteq \{x\}. \quad (1)$$

This means that $z = x[H_r]$. But now take any $i \in H - H_r$, pick any $d \neq x_i$ (here is where we use the assumption that $|\mathcal{D}| \geq 2$) and set

$$x' := (x_1, \dots, x_{i-1}, d, x_{i+1}, \dots, x_h).$$

Clearly, $x'[H_r] = x[H_r] = z$, contradicting (1). \square

Theorem 5.3. *Assume \mathcal{D}, \mathcal{A} are finite sets such that $|\mathcal{D}| \geq 2$ and $\mathcal{A} \neq \emptyset$. Then $\mathbb{1}_{\leq}$, \swarrow , \nearrow and $\swarrow\swarrow$ look for $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ as follows:*

$r =$	$\text{adom}(x)$	$\text{aatt}(a)$	$\text{adom}(x)$	$\text{aatt}(a)$
$s =$	$\text{coatt}(a)$	$\text{coatt}(b)$	$\text{codom}^H(y)$	$\text{codom}^H(y)$
$r \mathbb{1}_{\leq} s$	<i>always</i>	$a \neq b$	$x[H] \neq y$	$a \notin H$
$r \swarrow s$	<i>never</i>	$a = b$	$x[H] = y$	$a \in H$
$r \nearrow s$	<i>never</i>	$a = b$	$x[H] = y$	<i>never</i>
$r \swarrow\swarrow s$	<i>never</i>	$a = b$	<i>always</i>	<i>always</i>

Sketch. For the $\mathbb{1}_{\leq}$ -row: this is just spelling out the definition of \leq on $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ as restricted to $\mathcal{J}_{\mathcal{D},\mathcal{A}} \times \mathcal{M}_{\mathcal{D},\mathcal{A}}$.

For the \swarrow -row: the set of join-irreducibles consists of only of the atoms of the whole lattice, hence \swarrow is just the complement of \leq .

This observation already yields $\nearrow \subseteq \swarrow$ and $\swarrow = \nearrow$. The last missing piece of information to define \nearrow is provided by the analysis of restriction of \leq to $\mathcal{M}_{\mathcal{D},\mathcal{A}} \times \mathcal{M}_{\mathcal{D},\mathcal{A}}$:

$$\begin{array}{llll}
& r = \text{coatt}(a), & s = \text{coatt}(b), & \text{never,} \\
\text{for} & r = \text{coatt}(a), & s = \text{codom}^H(x), & r \leq s \quad \text{iff} \quad \text{never,} \\
& r = \text{codom}^H(x), & s = \text{coatt}(a), & a \in H, \\
& r = \text{codom}^H(x), & s = \text{codom}^H(y), & \text{never.}
\end{array}$$

Finally, for \swarrow we need to observe that composing \swarrow with $\nearrow \circ \swarrow$ does not allow to reach any new elements of $\mathcal{CoAtt}_{\mathcal{D},\mathcal{A}}$. As for elements of $\mathcal{M}_{\mathcal{D},\mathcal{A}}$ of the form $\text{codom}^H(y)$, note that

$$\exists h.(h \nearrow \text{coatt}(a) \& h \swarrow \text{codom}^H(y)) \text{ if } a \in H, \quad (2)$$

$$\exists h.(h \nearrow \text{codom}^{H_x}(x) \& h \swarrow \text{codom}^{H_y}(y)) \text{ if } x[H_x \cap H_y] = y[H_x \cap H_y]. \quad (3)$$

Furthermore, we have that

- for any $x \in {}^A\mathcal{D}$ and any $H \subseteq \mathcal{A}$, $\text{adom}(x) \swarrow \text{codom}^H(x[H])$,
- for any $a \in \mathcal{A}$ and any $x \in {}^A\mathcal{D}$, $\text{aatt}(a) \swarrow \text{codom}^{\mathcal{D}}(x)$.

Using (3), we obtain then that $\mathcal{J}_{\mathcal{D},\mathcal{A}} \times \{\mathbf{H}\} \subseteq \swarrow$ and using (3) again—that $\mathcal{J}_{\mathcal{D},\mathcal{A}} \times \{\text{codom}^H(y)\} \subseteq \swarrow$ for any $y \in {}^A\mathcal{D}$ and any $H \subseteq \mathcal{A}$. \square

Corollary 5.4. *If \mathcal{D}, \mathcal{A} are finite sets such that $|\mathcal{D}| \geq 2$ and $\mathcal{A} \neq \emptyset$, then $\mathfrak{R}(\mathcal{D}, \mathcal{A})$ is subdirectly irreducible but not simple.*

Proof. Follows immediately from Proposition 5.1 and Theorem 5.3. \square

6. Extending the Signature and Adding Typing Information

Clearly, it is possible to define more operations on $\mathcal{R}_{\text{unr}}^H$ than those present in \mathcal{L}_H . Thus, our first proposal for future study, regardless of the negative result in Corollary 4.8, is a systematic investigation of extensions of the signature. Let us discuss several natural ones; see also [29, 34].

6.1. Safe Extensions with Constants and Monotonic Relational Expressions

Let us begin with most natural additional constants. As we will see, just by adding a family of constants, we can express *monotonic relational expressions* of Sagiv and Yannakakis [28, Sec. 2.2].

The only extension we need are thus *unary singleton constants* introduced below. However, let us also mention several other safe extensions of the language, which will turn out to be expressible using these constants:

The top element $\top := (\emptyset, \{\emptyset\})$. Its inclusion in the signature would be harmless, but at the same time does not appear to improve expressivity in a significant way. Note, however, that if relations with empty header are seen as boolean predicates, then \mathbf{H} plays the role of **false** and \top is necessary to encode **true**. Also, in presence of at least one unary singleton constant and the header constant, it is definable anyway, as noted above.

Attribute constants $\underline{\mathbf{a}} := (\{\mathbf{a}\}, \emptyset)$, for $\mathbf{a} \in \mathcal{A}$. We touch upon an important difference between our setting and that of both *named SPJR algebra* and *unnamed SPC algebra* in [1, Ch. 4], which are *typed*: expressions come with an explicit information about their headers (*arities* in the unnamed case). Our expressions are untyped *query schemes*. On the one hand, \mathcal{L}_H allows, e.g., *projection of r to the header of s* : $r \oplus (s \times \mathbf{H})$, which does not correspond to any *single* SPJR expression. On the other hand, only with attribute constants we can write the SPJR *projection of r to a concrete header* $\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$: $\pi_{\underline{\mathbf{a}}_1, \dots, \underline{\mathbf{a}}_n}(r) := r \oplus \underline{\mathbf{a}}_1 \times \dots \times \underline{\mathbf{a}}_n$.

Unary singleton constants $(\underline{\mathbf{a}} : \mathbf{d}) := (\{\mathbf{a}\}, \{(\mathbf{a} : \mathbf{d})\})$, for $\mathbf{a} \in \mathcal{A}$, $\mathbf{d} \in \mathcal{D}$. These are among the *base SPJR queries* [1, p. 58]. Note they add more expressivity than attribute constants: whenever the signature includes $(\underline{\mathbf{a}} : \mathbf{d})$ for some $\mathbf{d} \in \mathcal{D}$, we have $\underline{\mathbf{a}} = (\underline{\mathbf{a}} : \mathbf{d}) \times \mathbf{H}$. They also allow defining \top as $\top = (\underline{\mathbf{a}} : \mathbf{d}) \oplus \mathbf{H}$ and, more importantly, the SPJR *constant-based selection queries* $\sigma_{\underline{\mathbf{a}}=\mathbf{d}}(r) := r \times (\underline{\mathbf{a}} : \mathbf{d})$.

Table 2: Equivalence between typed expressions and monotonic relational expressions [28, Sec. 2.2].

Typing system for positive expressions

Σ is a supply of relational symbols \mathbf{r} together with typing information, i.e., $\Sigma = \{\mathbf{r}_1 : H_1, \dots, \mathbf{r}_n : H_n\}$, where $H \subseteq_{fin} \mathcal{A}$

$$\frac{\mathbf{r} : H \in \Sigma}{\Sigma \vdash \mathbf{r} : H} \qquad \frac{d \in \mathcal{D}, a \in \mathcal{A}}{\Sigma \vdash \langle \mathbf{a} : d \rangle : \{a\}} \qquad \Sigma \vdash \mathbf{H} : \emptyset$$

$$\frac{\Sigma \vdash r_1 : H_1 \quad \Sigma \vdash r_2 : H_2}{\Sigma \vdash r_1 \bowtie r_2 : H_1 \cup H_2} \qquad \frac{\Sigma \vdash r_1 : H_1 \quad \Sigma \vdash r_2 : H_2}{\Sigma \vdash r_1 \oplus r_2 : H_1 \cap H_2}$$

Translation $\langle \cdot \rangle$ of monotonic relational expressions [28, Sec. 2.2] into our terms

Recall $\underline{\mathbf{a}} = \langle \mathbf{a} : d \rangle \bowtie \mathbf{H}$, for an arbitrary choice of $d \in \mathcal{D}$

$$\langle \pi_{\mathbf{a}_1, \dots, \mathbf{a}_n}(r) \rangle = \langle r \rangle \oplus \underline{\mathbf{a}_1} \bowtie \dots \bowtie \underline{\mathbf{a}_n} \qquad \langle \sigma_{\underline{\mathbf{a}}} r \rangle = \langle r \rangle \bowtie \langle \underline{\mathbf{a}} : d \rangle$$

$$\langle r_1 \bowtie r_2 \rangle = \langle r_1 \rangle \bowtie \langle r_2 \rangle \qquad \langle r_1 \cup r_2 \rangle = \langle r_1 \rangle \oplus \langle r_2 \rangle$$

Reverse translation $(\cdot)^\Sigma$

Fix $b \in \mathcal{A}$ and distinct $d, e \in \mathcal{D}$. Observe that only for the atomic expressions and \bowtie the translation is independent from the typing judgement

$$\frac{\mathbf{r} : H \in \Sigma}{(\mathbf{r})^\Sigma = \mathbf{r}} \qquad \frac{d \in \mathcal{D}, a \in \mathcal{A}}{(\langle \mathbf{a} : d \rangle)^\Sigma = \langle \mathbf{a} : d \rangle} \qquad (\mathbf{H})^\Sigma = \pi_\emptyset(\langle \mathbf{b} : d \rangle \bowtie \langle \mathbf{b} : e \rangle)$$

$$(r_1 \bowtie r_2)^\Sigma = (r_1)^\Sigma \bowtie (r_2)^\Sigma \qquad \frac{\Sigma \vdash r_1 : \{a_1, \dots, a_m, b_1, \dots, b_n\} \quad \Sigma \vdash r_2 : \{b_1, \dots, b_n, c_1, \dots, c_k\}}{(r_1 \oplus r_2)^\Sigma = \pi_{b_1, \dots, b_n}((r_1)^\Sigma) \cup \pi_{b_1, \dots, b_n}((r_2)^\Sigma)}$$

6.2. Equivalence with Monotonic Relational Expressions

As it turns out, the mere addition of unary singleton constants brings our language very close to that of monotonic relational expressions of Sagiv and Yannakakis [28, Sec. 2.2]. To be more precise, we obtain in this way untyped (but named!) counterparts of these expressions. The typing discipline necessary to connect these two formalisms is presented in Table 2.

6.3. Other Extensions

The bottom element $\perp := (\mathcal{A}, \emptyset)$. Whenever \mathcal{A} is infinite, including \perp in the signature would exclude subalgebras consisting of relations with finite headers—i.e., exactly those arising from concrete database instances. Another undesirable feature is that the interpretation of \perp depends on \mathcal{A} , i.e., the collection of all possible attributes, which is not explicitly supplied by a query expression. In other words, it is domain-independent, but not strictly independent.

The full relation $\mathbf{U} := (\mathcal{A}, \mathcal{A}\mathcal{D})$. [34, 29] Its inclusion would destroy even the ordinary domain independence property (d.i.p.). Note that for non-empty \mathcal{A} and \mathcal{D} , \mathbf{U} is a complement of \mathbf{H} .

The equality constant $\Delta := (\mathcal{A}, \{x \in \mathcal{A}\mathcal{D} \mid \forall a, a'. x(a) = x(a')\})$. With it, we can express the *equality-based selection queries*: $\sigma_{\underline{\mathbf{a}}=\underline{\mathbf{b}}}(r) := r \bowtie (\Delta \oplus \underline{\mathbf{a}} \bowtie \underline{\mathbf{b}})$. But the interpretation of Δ violates d.i.p., hence we prefer *the inner equality operator*:

$$\bar{r} := (H_r, \{x \in H_r \mathcal{D} \mid \exists x' \in r. \exists a' \in H_r. \forall a \in H_r. x(a) = x'(a')\}),$$

which also allows to define $\sigma_{\underline{\mathbf{a}}=\underline{\mathbf{b}}}(r)$ as $r \bowtie (\bar{r} \oplus \underline{\mathbf{a}} \bowtie \underline{\mathbf{b}})$.

The header-narrowing operator $r \pitchfork s := (H_r - H_s, \{x[H_r - H_s] \mid x \in H_r\})$. This one is perhaps more surprising, but now we can define the *attribute renaming* operators [1, p. 58] as $\rho_{\mathbf{a} \rightarrow \mathbf{b}}(r) := (r \bowtie \overline{(r \oplus \mathbf{a}) \bowtie (\mathbf{b} : \mathbf{d})}) \pitchfork \mathbf{a}$, where $d \in \mathcal{D}$ is arbitrary. Instead of using \pitchfork , one could add constants for elements $\mathbf{aatt}(\mathbf{a})$ introduced in Section 5, but this would lead to the same criticism as \perp above: indeed, such constants would make \perp definable as $\perp = \mathbf{aatt}(\mathbf{a}) \bowtie \mathbf{a}$ and hence fail strict independence. However, for the purpose of recovering the full setup of Codd's relational algebra, all we need are ...

The projecting-away operator(s).

$$\exists_a r := H_r - \{a\}, \{x \in H_r - \{a\} \mathcal{D} \mid \exists x' \in H_r \mathcal{D}. x' = x[H_r - \{a\}]\}$$

Attribute renaming operators can be defined now as $\rho_{\mathbf{a} \rightarrow \mathbf{b}}(r) := \pi_{\mathbf{a}}(\overline{r \bowtie \overline{(r \oplus \mathbf{a}) \bowtie (\mathbf{b} : \mathbf{d})}})$, where $d \in \mathcal{D}$ is arbitrary.

The difference operator $r - s := (H_r, \{x \in B_r \mid x \notin B_s\})$. This is a very natural extension from the DB point of view [1, Ch. 5], which leads us beyond the SPJRU setting towards the question of *relational completeness* [4]. Here again we break with the partial character of Codd's original operator. Another option would be $(H_r \cap s, \{x \in B_r[H_s] \mid x \notin B_s[H_r]\})$, but this one can be defined with the difference operator proposed here as $(r \oplus s) - (s \oplus (r \bowtie \mathbf{H}))$.

While we do not provide details here, it should be clear how to prove equipollence between the typed version of the formalism with the extensions proposed above and Codd's relational algebra in the spirit of Section 6.2 and Table 2.

7. Summary and Future Work

We have seen that relational lattices form an interesting class with rather surprising properties. Unlike Codd's relational algebra, all operations are total and in contrast to the encoding of relational algebras in cylindric algebras, the domain independence property obtains automatically. We believe that with the extensions of the language proposed in Section 6, one can ultimately obtain most natural algebraic treatment of SPRJ(U) operators and relational query languages. Besides, given how well investigated the lattice of varieties of lattices is in general [15], it is intriguing to discover a class of lattices with a natural CS motivation which does not seem to fit anywhere in the existing picture.

We posed a number of questions and problems in the text, in particular Open Problems 3.1, 3.6, 3.7 and 4.10. Without settling them we cannot claim to have grasped how relational lattices behave as an algebraic class. None of them seems trivial, even with the rich supply of algebraic logic tools available in existing literature. Comparison with other settings, like that of Craig [5], Quine [27], other (generalized) algebras of finite sequences and many-sorted cylindric/polyadic algebras [24, Sec. 7.1–7.4 and references therein] and possible attempts at transfer of methods and results would be also of interest.

We would also like to mention the natural question of *representability*:

Open Problem 7.1 (Hirsch). *Given a finite algebra in the signature $\mathcal{L}_{\mathbf{H}}$ (\mathcal{L}), is it decidable whether it belongs to $\mathbb{SP}(\mathcal{R}_{\text{unr}}^{\mathbf{H}})$, $\mathbb{SP}(\mathcal{R}_{\text{fin}}^{\mathbf{H}})$ ($\mathbb{SP}(\mathcal{R}_{\text{unr}})$, $\mathbb{SP}(\mathcal{R}_{\text{fin}})$)?*

We believe that the analysis of the concept structure of finite relational lattices in Section 5 may lead to an algorithm recognizing whether the concept lattice of a given context belongs to $\mathbb{SP}(\mathcal{R}_{\text{fin}}^{\mathbf{H}})$ (or $\mathbb{SP}(\mathcal{R}_{\text{fin}})$). It also opens the door to a systematic investigation of a research problem suggested by Yde Venema: *duality theory of relational lattices*. Given that relational lattices have much more meet-irreducible than join-irreducible elements, it is tempting to apply the recent duality of Frittella and Santocanale [7], which is particularly well-tailored for lattices with non-isomorphic sets of meet-irreducible and join-irreducible elements. See also Section 2.3 above for other category-theoretical connections: as suggested therein, the relationship with the work of Abramsky [2] would be of particular interest.

Acknowledgements. We would like to thank *Vadim Tropashko* and *Marshall Spight* for introducing the subject to the third author (who in turn introduced it to the other two) and discussing it in the usenet group `comp.databases.theory`, *Maarten Marx*, *Balder ten Cate*, *Jan Paredaens* for additional discussions and general support in an early phase of our cooperation and the *referees* of several incarnations of this work for the comments. The first author would also like to acknowledge: a) *Peter Jipsen* for discussions in September 2013 at the Chapman University leading to recovery, rewrite and extension of the material (in particular for Sec. 5) b) Peter Jipsen and *Drew Moshier* for even more discussions and suggestions during their (re-)visit in 2014, c) suggestions by participants of: TACL'09, ALCOP 2010, RAMiCS 2014, the Birmingham TCS seminar and TU Delft Applied Logic Seminar (in particular for Sec. 2.3, 6 and 7), d) *Neil Ghani* for mentioning opfibrations in a conversation, *Alexander Kurz* for making this conversation possible and *Stefan Milius* for an interesting discussion of Abramsky's paper (which sadly was not reflected in the present work).

- [1] Abiteboul, S., Hull, R., Vianu, V., 1995. Foundations of Databases. Addison-Wesley.
- [2] Abramsky, S., 2013. Relational databases and Bell's theorem. In: Tannen, V., Wong, L., Libkin, L., Fan, W., Tan, W.-C., Fourman, M. (Eds.), In Search of Elegance in the Theory and Practice of Computation. Vol. 8000 of Lecture Notes in Computer Science. Springer Berlin Heidelberg, pp. 13–35.
URL http://dx.doi.org/10.1007/978-3-642-41660-6_2
- [3] Chandra, A. K., Merlin, P. M., 1977. Optimal implementation of conjunctive queries in relational data bases. In: Proceedings of the Ninth Annual ACM Symposium on Theory of Computing. STOC '77. ACM, New York, NY, USA, pp. 77–90.
URL <http://doi.acm.org/10.1145/800105.803397>
- [4] Codd, E. F., 1970. A relational model of data for large shared data banks. Commun. ACM 13 (6), 377–387.
- [5] Craig, W., 1974. Logic in Algebraic Form. Three Languages and Theories. Vol. 72 of Studies in Logic and the Foundations of Mathematics. North Holland.
- [6] Düntsch, I., Mikuláš, S., 2001. Cylindric structures and dependencies in relational databases. Theor. Comput. Sci. 269 (1–2), 451–468.
- [7] Frittella, S., Santocanale, L., 2014. Fixed-point theory in the varieties \mathcal{D}_n . In: Höfner, P., Jipsen, P., Kahl, W., Müller, M. E. (Eds.), Relational and Algebraic Methods in Computer Science - 14th International Conference, RAMiCS 2014, Marienstatt, Germany, April 28-May 1, 2014. Proceedings. Vol. 8428 of Lecture Notes in Computer Science. Springer, pp. 446–462.
URL http://dx.doi.org/10.1007/978-3-319-06251-8_27
- [8] Ganter, B., Wille, R., 1996. Applied lattice theory: Formal concept analysis. In: Grätzer, G. (Ed.), General Lattice Theory, second edition, 2nd Edition. Birkhäuser, pp. 591–606.
- [9] Gurevich, Y., 1966. The word problem for certain classes of semigroups. Algebra and Logic 5, 25–35.
- [10] Gurevich, Y., Lewis, H. R., 1984. The word problem for cancellation semigroups with zero. The Journal of Symbolic Logic 49 (1), 184–191.
URL <http://www.jstor.org/stable/2274101>
- [11] Harrop, R., 1958. On the existence of finite models and decision procedures for propositional calculi. Mathematical Proceedings of the Cambridge Philosophical Society 54, 1–13.
URL <http://journals.cambridge.org/article.S0305004100033120>
- [12] Hirsch, R., Hodkinson, I., 2002. Relation Algebras by Games. Vol. 147 of Studies in Logic and the Foundations of Mathematics. Elsevier.
- [13] Imieliński, T., Lipski, W., 1984. The relational model of data and cylindric algebras. Journal of Computer and System Sciences 28 (1), 80–102.
URL <http://www.sciencedirect.com/science/article/pii/002200084900771>
- [14] Jacobs, B., 1999. Categorical Logic and Type Theory. No. 141 in Studies in Logic and the Foundations of Mathematics. North Holland, Amsterdam.
- [15] Jipsen, P., Rose, H., 1992. Varieties of Lattices. Vol. 1533 of Lecture Notes in Mathematics. Springer Verlag.
- [16] Jipsen, P., Rose, H., 1998. Varieties of lattices. In: Grätzer, G. (Ed.), General Lattice Theory. Birkhäuser Verlag, pp. 555–574, appendix F to the second edition.
- [17] Johnson, D. S., Klug, A., 1982. Testing containment of conjunctive queries under functional and inclusion dependencies. In: Proceedings of the 1st ACM SIGACT-SIGMOD Symposium on Principles of Database Systems. PODS '82. ACM, New York, NY, USA, pp. 164–169.
URL <http://doi.acm.org/10.1145/588111.588138>
- [18] Johnson, M., Rosebrugh, R., Dec. 2007. Fibrations and universal view updatability. Theoretical Computer Science 388 (1–3), 109–129.
URL <http://dx.doi.org/10.1016/j.tcs.2007.06.004>
- [19] Johnson, M., Rosebrugh, R., Wood, R. J., Feb. 2012. Lenses, fibrations and universal translations. Mathematical Structures in Computer Science 22 (1), 25–42.
URL <http://dx.doi.org/10.1017/S0960129511000442>
- [20] Litak, T., Mikuláš, S., Hidders, J., 2014. Relational lattices. In: Höfner, P., Jipsen, P., Kahl, W., Müller, M. E. (Eds.), Relational and Algebraic Methods in Computer Science 2014 (RAMiCS). Vol. 8428 of Lecture Notes in Computer Science.

- Springer International Publishing, pp. 327–343.
URL http://dx.doi.org/10.1007/978-3-319-06251-8_20
- [21] Maddux, R., 1980. The equational theory of CA_3 is undecidable. *The Journal of Symbolic Logic* 45 (2), 311–316.
URL <http://www.jstor.org/stable/2273191>
- [22] Maeda, S., 1974. Locally modular lattices and locally distributive lattices. *Proceedings of the American Mathematical Society* 44 (2), pp. 237–243.
URL <http://www.jstor.org/stable/2040413>
- [23] McKenzie, R., 1972. Equational bases and non-modular lattice varieties. *Trans. Amer. Math. Soc.* 174, 1–43.
- [24] Némethi, I., 1991. Algebraization of quantifier logics, an introductory overview. *Studia Logica: An International Journal for Symbolic Logic* 50 (3/4), pp. 485–569.
URL <http://www.jstor.org/stable/20015599>
- [25] Padmanabhan, R., McCune, W., Veroff, R., 2007. Lattice laws forcing distributivity under unique complementation. *Houston Journal of Mathematics* 33 (2), 391–401.
- [26] Post, E. L., 1947. Recursive unsolvability of a problem of Thue. *The Journal of Symbolic Logic* 12 (1), 1–11.
URL <http://www.jstor.org/stable/2267170>
- [27] Quine, W. V., 1936. Toward a calculus of concepts. *The Journal of Symbolic Logic* 1 (1), pp. 2–25.
URL <http://www.jstor.org/stable/2269323>
- [28] Sagiv, Y., Yannakakis, M., Oct. 1980. Equivalences among relational expressions with the union and difference operators. *J. ACM* 27 (4), 633–655.
URL <http://doi.acm.org/10.1145/322217.322221>
- [29] Spight, M., Tropashko, V., 2006. First steps in relational lattice. <http://arxiv.org/abs/cs/0603044>.
URL <http://www.citebase.org/abstract?id=oai:arXiv.org:cs/0603044>
- [30] Spivak, D. I., 12 2014. Database queries and constraints via lifting problems. *Mathematical Structures in Computer Science* 24.
URL <http://journals.cambridge.org/article.S0960129513000479>
- [31] Stanley, R. P., 1972. Supersolvable lattices. *Algebra Universalis* 2 (1), 197–217.
URL <http://dx.doi.org/10.1007/BF02945028>
- [32] Stern, M., 1999. *Semimodular Lattices*. Vol. 73 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press.
- [33] Tropashko, V., 2005. Relational algebra as non-distributive lattice. <http://arxiv.org/abs/cs/0501053>.
URL <http://arxiv.org/abs/cs/0501053>
- [34] Tropashko, V., 2009–13. The website of QBQL: Prototype of relational lattice system, <https://code.google.com/p/qbql/>.
- [35] Van den Bussche, J., Van Gucht, D., Vansummeren, S., 2007. A crash course on database queries. In: *PODS '07: Proceedings of the twenty-sixth ACM SIGMOD-SIGACT-SIGART symposium on Principles of Database Systems*. ACM, New York, NY, USA, pp. 143–154.
- [36] Vardi, M. Y., 1981. The decision problem for database dependencies. *Inf. Process. Lett.* 12 (5), 251–254.
URL [http://dx.doi.org/10.1016/0020-0190\(81\)90025-9](http://dx.doi.org/10.1016/0020-0190(81)90025-9)

Appendix A. Theorem 3.4, Clause 5:

formulas (assumptions) .

```

x ^ y = y ^ x .
(x ^ y) ^ z = x ^ (y ^ z) .
x v y = y v x .
(x v y) v z = x v (y v z) .
x v (x ^ y) = x .
x ^ (x v y) = x .
UpMe(x,y,z) = x ^ (y v z) .
LoMe(x,y,z) = (x ^ y) v (x ^ z) .
UpJo(x,y,z) = (x v y) ^ (x v z) .
LoJo(x,y,z) = x v (y ^ z) .
end_of_list .

```

formulas (goals) .

```

(all x1 all y1 all w UpMe(a ^ x1,y1,w) v (y1 ^ w) = (((a ^ x1) ^ y1) v w)
  ^ (((a ^ x1) ^ w) v y1)) ->

UpMe(a ^ z1,z2,z3) = LoMe(a ^ z1,z2,z3) .

```

end_of_list.

.....

===== PROOF =====

% Proof 1 at 45.03 (+ 0.26) seconds.

% Length of proof is 66.

% Level of proof is 14.

% Maximum clause weight is 35.

% Given clauses 464.

```
1 (all x1 all y1 all w UpMe(a ^ x1,y1,w) v (y1 ^ w) = (((a ^ x1) ^ y1) v w
   ) ^ (((a ^ x1) ^ w) v y1)) -> UpMe(a ^ z1,z2,z3) = LoMe(a ^ z1,z2,z3) #
   label(non_clause) # label(goal). [goal].
2 x ^ y = y ^ x. [assumption].
3 (x ^ y) ^ z = x ^ (y ^ z). [assumption].
4 x v y = y v x. [assumption].
5 (x v y) v z = x v (y v z). [assumption].
6 x v (x ^ y) = x. [assumption].
7 x ^ (x v y) = x. [assumption].
8 UpMe(x,y,z) = x ^ (y v z). [assumption].
9 LoMe(x,y,z) = (x ^ y) v (x ^ z). [assumption].
12 UpMe(a ^ x,y,z) v (y ^ z) = (((a ^ x) ^ y) v z) ^ (((a ^ x) ^ z) v y).
   [deny(1)].
13 (x ^ y) v (a ^ (z ^ (x v y))) = ((a ^ (z ^ x)) v y) ^ ((a ^ (z ^ y)) v
   x). [copy(12),rewrite([8(3),3(4),4(6),3(9),3(13)])].
14 LoMe(a ^ c1,c2,c3) != UpMe(a ^ c1,c2,c3). [deny(1)].
15 (c2 ^ (a ^ c1)) v (c3 ^ (a ^ c1)) != a ^ (c1 ^ (c2 v c3)). [copy(14),
   rewrite([9(6),2(5),2(10),8(17),3(18)])].
16 x ^ (y ^ z) = z ^ (x ^ y). [para(3(a,1),2(a,1))].
17 x ^ (y ^ z) = y ^ (x ^ z). [para(2(a,1),3(a,1,1)),rewrite([3(2)])].
18 (a ^ (c1 ^ c2)) v (a ^ (c1 ^ c3)) != a ^ (c1 ^ (c2 v c3)). [
   back_rewrite(15),rewrite([16(5),2(4),17(5),16(10),2(9),17(10)])].
20 x v (y v z) = y v (x v z). [para(4(a,1),5(a,1,1)),rewrite([5(2)])].
21 x v (y ^ x) = x. [para(2(a,1),6(a,1,2))].
22 (x ^ y) v (x ^ (y ^ z)) = x ^ y. [para(3(a,1),6(a,1,2))].
23 x v ((x ^ y) v z) = x v z. [para(6(a,1),5(a,1,1)),flip(a)].
26 x ^ (y ^ ((x ^ y) v z)) = x ^ y. [para(7(a,1),3(a,1)),flip(a)].
27 x ^ (y v x) = x. [para(4(a,1),7(a,1,2))].
28 (x v y) ^ (x v (y v z)) = x v y. [para(5(a,1),7(a,1,2))].
29 x v x = x. [para(7(a,1),6(a,1,2))].
30 x ^ x = x. [para(6(a,1),7(a,1,2))].
38 (x ^ (y ^ z)) v (a ^ (u ^ ((x ^ y) v z))) = ((a ^ (u ^ (x ^ y))) v z) ^
   ((a ^ (u ^ z)) v (x ^ y)). [para(3(a,1),13(a,1,1))].
63 x ^ (y ^ (x v z)) = y ^ x. [para(7(a,1),17(a,1,2)),flip(a)].
76 x v (y v x) = y v x. [para(29(a,1),5(a,2,2)),rewrite([4(2)])].
79 x ^ (x ^ y) = x ^ y. [para(30(a,1),3(a,1,1)),flip(a)].
81 x ^ (y ^ x) = y ^ x. [para(30(a,1),3(a,2,2)),rewrite([2(2)])].
82 (x ^ y) v (a ^ (x v y)) = ((a ^ x) v y) ^ ((a ^ y) v x). [para(30(a,1),
   13(a,1,2,2)),rewrite([2(8),7(8),2(11),27(11)])].
```

88 $x \vee (y \wedge (z \wedge x)) = x$. [para(3(a,1),21(a,1,2))].
89 $x \vee ((y \wedge x) \vee z) = x \vee z$. [para(21(a,1),5(a,1,1)), flip(a)].
95 $x \wedge ((y \vee x) \wedge z) = x \wedge z$. [para(27(a,1),3(a,1,1)), flip(a)].
97 $x \wedge (y \vee (z \vee x)) = x$. [para(5(a,1),27(a,1,2))].
111 $x \vee (y \vee (z \wedge x)) = y \vee x$. [para(21(a,1),20(a,1,2)), flip(a)].
132 $x \vee (y \vee (z \vee x)) = y \vee (z \vee x)$. [para(5(a,1),76(a,1,2)), rewrite([5(5)])].
141 $(x \wedge y) \vee ((x \wedge (y \wedge z)) \vee u) = (x \wedge y) \vee u$. [para(22(a,1),5(a,1,1)), flip(a)].
150 $x \wedge (y \wedge (z \wedge x)) = y \wedge (z \wedge x)$. [para(3(a,1),81(a,1,2)), rewrite([3(5)])].
155 $(x \wedge y) \vee (y \wedge x) = x \wedge y$. [para(81(a,1),22(a,1,2))].
158 $x \vee ((y \wedge (z \wedge x)) \vee u) = x \vee u$. [para(88(a,1),5(a,1,1)), flip(a)].
176 $a \wedge (x \wedge ((y \vee z) \wedge (u \vee ((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)))) = a \wedge (x \wedge (y \vee z))$. [para(13(a,1),97(a,1,2,2)), rewrite([3(15),3(14)])].
195 $(x \vee y) \wedge ((x \wedge z) \vee y) = (x \wedge z) \vee y$. [para(23(a,1),27(a,1,2)), rewrite([2(4)])].
197 $(x \wedge y) \vee (z \vee x) = z \vee x$. [para(76(a,1),23(a,2)), rewrite([132(4)])].
220 $(x \wedge y) \vee (x \wedge (y \vee z)) = x \wedge (y \vee z)$. [para(63(a,1),21(a,1,2)), rewrite([4(4)])].
228 $(x \vee y) \wedge ((z \wedge x) \vee y) = (z \wedge x) \vee y$. [para(89(a,1),27(a,1,2)), rewrite([2(4)])].
244 $(x \vee y) \wedge (z \wedge y) = z \wedge y$. [para(81(a,1),95(a,2)), rewrite([150(4)])].
321 $(a \wedge x) \vee (((a \wedge x) \vee y) \wedge z) = ((a \wedge x) \vee z) \wedge ((a \wedge x) \vee y)$. [para(26(a,1),13(a,2,1,1)), rewrite([5(9),26(11),4(7),20(17),141(17)])].
336 $(x \wedge y) \vee (z \vee (y \wedge x)) = z \vee (y \wedge x)$. [para(155(a,1),5(a,2,2)), rewrite([4(4)])].
341 $(x \wedge (y \wedge z)) \vee (y \wedge x) = x \wedge y$. [para(155(a,1),23(a,2)), rewrite([3(3),336(6)])].
398 $(x \vee y) \wedge (y \vee x) = x \vee y$. [para(76(a,1),28(a,1,2))].
565 $(x \vee (y \vee z)) \wedge (z \vee y) = z \vee y$. [para(398(a,1),244(a,1,2)), rewrite([398(7)])].
1281 $(x \vee y) \wedge ((z \wedge (u \wedge x)) \vee y) = (z \wedge (u \wedge x)) \vee y$. [para(158(a,1),27(a,1,2)), rewrite([2(5)])].
3918 $(x \vee y) \wedge ((z \wedge y) \vee x) = (z \wedge y) \vee x$. [para(111(a,1),565(a,1,1))].
6421 $x \wedge ((x \wedge y) \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$. [para(6(a,1),195(a,1,1))].
6979 $(x \wedge (y \wedge z)) \vee (x \wedge (u \vee y)) = x \wedge (u \vee y)$. [para(197(a,1),220(a,1,2,2)), rewrite([197(8)])].
7074 $x \wedge ((y \wedge z) \vee (y \wedge x)) = x \wedge y$. [para(341(a,1),228(a,1,2)), rewrite([2(5),3(5),6421(4),341(8)])].
7542 $x \wedge ((y \wedge z) \vee (x \wedge y)) = x \wedge y$. [para(2(a,1),7074(a,1,2,2))].
7744 $x \wedge ((y \wedge z) \vee (x \wedge (u \vee z))) = x \wedge (u \vee z)$. [para(244(a,1),7542(a,1,2,1))].
8021 $(a \wedge x) \vee (a \wedge y) = a \wedge (x \vee y)$. [para(82(a,2),38(a,1,2,2)), rewrite([7744(10),6979(7),2(10),3(10),27(9),79(7),2(11),26(12),3918(12)]), flip(a)].
8518 $a \wedge ((c1 \wedge c2) \vee (c1 \wedge c3)) \neq a \wedge (c1 \wedge (c2 \vee c3))$. [back_rewrite(18), rewrite([8021(11)])].
9294 $a \wedge ((a \wedge x) \vee y) = a \wedge (x \vee y)$. [para(79(a,1),8021(a,1,1)), rewrite([8021(5)]), flip(a)].

10727 $a \wedge (x \wedge ((a \wedge y) \vee z)) = x \wedge (a \wedge (y \vee z))$. [para(9294(a,1),17(a,1,2)),flip(a)].
15426 $x \wedge (a \wedge ((x \wedge y) \vee z)) = a \wedge (x \wedge (y \vee z))$. [para(6(a,1),176(a,1,2,2,2)),rewrite([1281(7),10727(7)])].
36559 $a \wedge ((x \wedge y) \vee (x \wedge z)) = a \wedge (x \wedge (y \vee z))$. [para(341(a,1),321(a,1,2,1)),rewrite([3(6),8021(7),341(15),2(12),3(12),10727(12),15426(10)])].
36560 \$F. [resolve(36559,a,8518,a)].

===== **end of proof**=====

===== **STATISTICS**=====

Given=464. Generated=384816. Kept=36556. proofs=1.
Usable=432. Sos=19999. Demods=18086. Limbo=2, Disabled=16134. Hints=0.
Kept_by_rule=0, Deleted_by_rule=385.
Forward_subsumed=290602. Back_subsumed=991.
Sos_limit_deleted=57273. Sos_displaced=7640. Sos_removed=0.
New_demodulators=28671 (6 lex), Back_demodulated=7489. Back_unit_deleted=0.
Demod_attempts=7921128. Demod_rewrites=1153227.
Res_instance_prunes=0. Para_instance_prunes=0. Basic_paramod_prunes=0.
Nonunit_fsub_feature_tests=0. Nonunit_bsub_feature_tests=0.
Megabytes=75.19.
User_CPU=45.03, System_CPU=0.26, Wall_clock=46.

===== **end of statistics**=====

===== **end of search**=====

THEOREM PROVED

Appendix B. Theorem 3.4, Clause 6:

formulas(assumptions).
 $x \wedge y = y \wedge x$.
 $(x \wedge y) \wedge z = x \wedge (y \wedge z)$.
 $x \vee y = y \vee x$.
 $(x \vee y) \vee z = x \vee (y \vee z)$.
 $x \vee (x \wedge y) = x$.
 $x \wedge (x \vee y) = x$.
UpMe(x,y,z) = $x \wedge (y \vee z)$.
LoMe(x,y,z) = $(x \wedge y) \vee (x \wedge z)$.
UpJo(x,y,z) = $(x \vee y) \wedge (x \vee z)$.
LoJo(x,y,z) = $x \vee (y \wedge z)$.

```
UpMe(a ^ x1,y1,z1) v (y1 ^ z1) = (((a ^ x1) ^ y1) v z1) ^ (((a ^ x1) ^ z1)
v y1).
```

```
UpMe(x,y,z) = UpMe(x,y,a ^ z) v UpMe(x,z,a ^ y).
```

```
end_of_list.
```

```
formulas(goals).
```

```
(all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> UpMe(x2,y2,z2) =
LoMe(x2,y2,z2))).
```

```
end_of_list.
```

```
....
```

```
===== PROOF =====
```

```
\% Proof 1 at 222.55 (+ 1.51) seconds.
```

```
\% Length of proof is 195.
```

```
\% Level of proof is 24.
```

```
\% Maximum clause weight is 47.
```

```
\% Given clauses 1611.
```

```
1 (all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> UpMe(x2,y2,z2) =
LoMe(x2,y2,z2))) # label(non_clause) # label(goal). [goal].
```

```
2 x ^ y = y ^ x. [assumption].
```

```
3 (x ^ y) ^ z = x ^ (y ^ z). [assumption].
```

```
4 x v y = y v x. [assumption].
```

```
5 (x v y) v z = x v (y v z). [assumption].
```

```
6 x v (x ^ y) = x. [assumption].
```

```
7 x ^ (x v y) = x. [assumption].
```

```
8 UpMe(x,y,z) = x ^ (y v z). [assumption].
```

```
9 LoMe(x,y,z) = (x ^ y) v (x ^ z). [assumption].
```

```
12 UpMe(a ^ x,y,z) v (y ^ z) = (((a ^ x) ^ y) v z) ^ (((a ^ x) ^ z) v y).
[assumption].
```

```
13 (x ^ y) v (a ^ (z ^ (x v y))) = ((a ^ (z ^ x)) v y) ^ ((a ^ (z ^ y)) v
x). [copy(12),rewrite([8(3),3(4),4(6),3(9),3(13)])].
```

```
14 UpMe(x,y,z) = UpMe(x,y,a ^ z) v UpMe(x,z,a ^ y). [assumption].
```

```
15 (x ^ (y v (a ^ z))) v (x ^ (z v (a ^ y))) = x ^ (y v z). [copy(14),
rewrite([8(1),8(5),8(9)]),flip(a)].
```

```
16 UpMe(a,c1,c3) = UpMe(a,c1,c2). [deny(1)].
```

```
17 a ^ (c1 v c3) = a ^ (c1 v c2). [copy(16),rewrite([8(4),8(9)])].
```

```
18 LoMe(c1,c2,c3) != UpMe(c1,c2,c3). [deny(1)].
```

```
19 (c1 ^ c2) v (c1 ^ c3) != c1 ^ (c2 v c3). [copy(18),rewrite([9(4),8(11)
])].
```

```
21 x ^ (y ^ z) = y ^ (x ^ z). [para(2(a,1),3(a,1,1)),rewrite([3(2)])].
```

```
23 x v (y v z) = y v (x v z). [para(4(a,1),5(a,1,1)),rewrite([5(2)])].
```

```
24 x v (y ^ x) = x. [para(2(a,1),6(a,1,2))].
```

```
25 (x ^ y) v (x ^ (y ^ z)) = x ^ y. [para(3(a,1),6(a,1,2))].
```

```
26 x v ((x ^ y) v z) = x v z. [para(6(a,1),5(a,1,1)),flip(a)].
```

```
27 x v (y v ((x v y) ^ z)) = x v y. [para(6(a,1),5(a,1)),flip(a)].
```

```
28 x ^ ((x v y) ^ z) = x ^ z. [para(7(a,1),3(a,1,1)),flip(a)].
```

```
29 x ^ (y ^ ((x ^ y) v z)) = x ^ y. [para(7(a,1),3(a,1)),flip(a)].
```

```
30 x ^ (y v x) = x. [para(4(a,1),7(a,1,2))].
```


31 $(x \vee y) \wedge (x \vee (y \vee z)) = x \vee y.$ $[para(5(a,1),7(a,1,2))].$
32 $x \vee x = x.$ $[para(7(a,1),6(a,1,2))].$
33 $x \wedge x = x.$ $[para(6(a,1),7(a,1,2))].$
36 $(x \wedge y) \vee (a \wedge ((x \vee y) \wedge z)) = ((a \wedge (z \wedge x)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x).$ $[para(2(a,1),13(a,1,2,2))].$
37 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = ((a \wedge (x \wedge z)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x).$ $[para(2(a,1),13(a,2,1,1,2))].$
41 $(x \wedge (y \wedge z)) \vee (a \wedge (u \wedge ((x \wedge y) \vee z))) = ((a \wedge (u \wedge (x \wedge y))) \vee z) \wedge ((a \wedge (u \wedge z)) \vee (x \wedge y)).$ $[para(3(a,1),13(a,1,1))].$
44 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = (y \vee (a \wedge (z \wedge x))) \wedge ((a \wedge (z \wedge y)) \vee x).$ $[para(4(a,1),13(a,2,1))].$
55 $((a \wedge x) \vee y) \wedge (x \vee ((a \wedge (x \wedge y)) \vee z)) = (a \wedge x) \vee ((x \vee z) \wedge y).$ $[para(7(a,1),13(a,2,1,1,2)),rewrite([5(5),7(6),4(5),23(13)]),flip(a)].$
58 $((a \wedge (x \wedge y)) \vee z) \wedge u \vee (((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)) = (y \wedge z) \vee ((a \wedge (x \wedge (y \vee z))) \vee (((a \wedge (x \wedge y)) \vee z) \wedge u)).$ $[para(13(a,2),9(a,2,2)),rewrite([9(9),23(27),4(26)])].$
68 $(x \wedge (y \vee (a \wedge z))) \vee ((z \vee (a \wedge y)) \wedge x) = x \wedge (y \vee z).$ $[para(2(a,1),15(a,1,2))].$
83 $a \wedge ((c1 \vee c3) \wedge x) = a \wedge ((c1 \vee c2) \wedge x).$ $[para(17(a,1),3(a,1,1)),rewrite([3(6)]),flip(a)].$
90 $x \vee (y \wedge (x \wedge z)) = x.$ $[para(21(a,1),6(a,1,2))].$
91 $x \wedge (y \wedge (x \vee z)) = y \wedge x.$ $[para(7(a,1),21(a,1,2)),flip(a)].$
92 $(x \wedge (y \wedge z)) \vee (a \wedge (u \wedge (y \vee (x \wedge z)))) = ((a \wedge (u \wedge y)) \vee (x \wedge z)) \wedge ((a \wedge (u \wedge (x \wedge z))) \vee y).$ $[para(21(a,1),13(a,1,1))].$
108 $x \vee (y \vee x) = y \vee x.$ $[para(32(a,1),5(a,2,2)),rewrite([4(2)])].$
112 $x \wedge (x \wedge y) = x \wedge y.$ $[para(33(a,1),3(a,1,1)),flip(a)].$
114 $x \wedge (y \wedge x) = y \wedge x.$ $[para(33(a,1),3(a,2,2)),rewrite([2(2)])].$
115 $(x \wedge y) \vee (a \wedge (x \vee y)) = ((a \wedge x) \vee y) \wedge ((a \wedge y) \vee x).$ $[para(33(a,1),13(a,1,2,2)),rewrite([2(8),7(8),2(11),30(11)])].$
132 $x \vee (y \wedge (z \wedge x)) = x.$ $[para(3(a,1),24(a,1,2))].$
133 $x \vee ((y \wedge x) \vee z) = x \vee z.$ $[para(24(a,1),5(a,1,1)),flip(a)].$
140 $(x \wedge y) \vee (x \wedge (z \wedge y)) = x \wedge y.$ $[para(21(a,1),24(a,1,2))].$
141 $x \wedge ((y \vee x) \wedge z) = x \wedge z.$ $[para(30(a,1),3(a,1,1)),flip(a)].$
142 $x \wedge (y \wedge (z \vee (x \wedge y))) = x \wedge y.$ $[para(30(a,1),3(a,1)),flip(a)].$
143 $x \wedge (y \vee (z \vee x)) = x.$ $[para(5(a,1),30(a,1,2))].$
146 $((a \wedge x) \vee y) \wedge ((a \wedge (x \wedge y)) \vee (z \vee x)) = (a \wedge x) \vee (y \wedge (z \vee x)).$ $[para(30(a,1),13(a,2,2,1,2)),rewrite([143(6),4(5),2(14)]),flip(a)].$
149 $x \wedge (y \wedge (z \vee x)) = y \wedge x.$ $[para(30(a,1),21(a,1,2)),flip(a)].$
152 $x \vee (y \vee (x \wedge z)) = y \vee x.$ $[para(6(a,1),23(a,1,2)),flip(a)].$
153 $x \wedge (y \vee (x \vee z)) = x.$ $[para(23(a,1),7(a,1,2))].$
159 $x \vee (y \vee (z \wedge x)) = y \vee x.$ $[para(24(a,1),23(a,1,2)),flip(a)].$
160 $(x \vee y) \wedge (x \vee (z \vee y)) = x \vee y.$ $[para(23(a,1),30(a,1,2))].$
163 $x \vee ((y \wedge (x \wedge z)) \vee u) = x \vee u.$ $[para(90(a,1),5(a,1,1)),flip(a)].$
170 $x \wedge (y \wedge (x \wedge z)) = y \wedge (x \wedge z).$ $[para(90(a,1),30(a,1,2)),rewrite([2(3)])].$
171 $x \vee (y \vee (z \wedge (x \wedge u))) = y \vee x.$ $[para(90(a,1),23(a,1,2)),flip(a)].$
174 $(a \wedge x) \vee ((x \vee y) \wedge z) = ((a \wedge x) \vee z) \wedge (x \vee y).$ $[back_rewrite(55),rewrite([163(8)]),flip(a)].$
185 $x \vee (y \vee (z \vee x)) = y \vee (z \vee x).$ $[para(5(a,1),108(a,1,2)),rewrite([5(5)])].$
196 $(x \wedge y) \vee ((x \wedge (y \wedge z)) \vee u) = (x \wedge y) \vee u.$ $[para(25(a,1),5(a,1,1)),flip(a)].$

207 $x \wedge (y \wedge (z \wedge x)) = y \wedge (z \wedge x)$. $[para(3(a,1),114(a,1,2)),rewrite([3(5)])]$.
213 $(x \wedge y) \vee (y \wedge x) = x \wedge y$. $[para(114(a,1),25(a,1,2))]$.
215 $x \vee (y \wedge (z \wedge (u \wedge x))) = x$. $[para(3(a,1),132(a,1,2,2))]$.
216 $x \vee ((y \wedge (z \wedge x)) \vee u) = x \vee u$. $[para(132(a,1),5(a,1,1)),flip(a)]$.
217 $x \vee (y \vee (z \wedge (u \wedge (x \vee y)))) = x \vee y$. $[para(132(a,1),5(a,1)),flip(a)]$.
231 $x \wedge (y \wedge (z \vee (u \vee (x \wedge y)))) = x \wedge y$. $[para(143(a,1),3(a,1)),flip(a)]$.
232 $x \wedge (y \vee (z \vee (u \vee x))) = x$. $[para(5(a,1),143(a,1,2,2))]$.
236 $a \wedge (x \wedge ((y \vee z) \wedge (u \vee (((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)))) = a \wedge (x \wedge (y \vee z))$. $[para(13(a,1),143(a,1,2,2)),rewrite([3(15),3(14)])]$.
259 $x \vee (((a \wedge (y \wedge x)) \vee z) \wedge ((a \wedge (y \wedge z)) \vee x)) = x \vee (a \wedge (y \wedge (x \vee z)))$. $[para(13(a,1),26(a,1,2))]$.
261 $(x \vee y) \wedge ((x \wedge z) \vee y) = (x \wedge z) \vee y$. $[para(26(a,1),30(a,1,2)),rewrite([2(4)])]$.
262 $x \vee (y \vee ((x \wedge z) \vee u)) = y \vee (x \vee u)$. $[para(26(a,1),23(a,1,2)),flip(a)]$.
263 $(x \wedge y) \vee (z \vee x) = z \vee x$. $[para(108(a,1),26(a,2)),rewrite([185(4)])]$.
267 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee ((x \vee u) \wedge y)))) = (x \vee (a \wedge (z \wedge ((x \vee u) \wedge y)))) \wedge ((a \wedge (z \wedge x)) \vee ((x \vee u) \wedge y))$. $[para(28(a,1),13(a,1,1)),rewrite([4(20),2(21)])]$.
276 $(x \wedge y) \vee ((x \vee z) \wedge y) = (x \vee z) \wedge y$. $[para(28(a,1),24(a,1,2)),rewrite([4(4)])]$.
277 $(x \vee y) \wedge (z \wedge x) = z \wedge x$. $[para(114(a,1),28(a,2)),rewrite([207(4)])]$.
280 $x \wedge (y \wedge (z \wedge (x \vee u))) = y \wedge (z \wedge x)$. $[para(3(a,1),91(a,1,2)),rewrite([3(6)])]$.
292 $(x \wedge y) \vee (x \wedge (y \vee z)) = x \wedge (y \vee z)$. $[para(91(a,1),24(a,1,2)),rewrite([4(4)])]$.
297 $x \vee (((a \wedge (y \wedge z)) \vee x) \wedge ((a \wedge (y \wedge x)) \vee z)) = x \vee (a \wedge (y \wedge (z \vee x)))$. $[para(13(a,1),133(a,1,2))]$.
301 $(x \wedge y) \vee ((x \wedge (z \wedge y)) \vee u) = (x \wedge y) \vee u$. $[para(21(a,1),133(a,1,2,1))]$.
302 $(x \vee y) \wedge ((z \wedge x) \vee y) = (z \wedge x) \vee y$. $[para(133(a,1),30(a,1,2)),rewrite([2(4)])]$.
304 $(x \wedge y) \vee (z \vee y) = z \vee y$. $[para(108(a,1),133(a,2)),rewrite([185(4)])]$.
321 $(x \vee y) \wedge (z \wedge y) = z \wedge y$. $[para(114(a,1),141(a,2)),rewrite([207(4)])]$.
338 $x \vee (y \vee (((x \wedge z) \vee y) \wedge u)) = x \vee y$. $[para(27(a,1),26(a,1,2)),rewrite([26(3)]),flip(a)]$.
352 $a \wedge (c3 \wedge (c1 \vee c2)) = a \wedge c3$. $[para(17(a,1),149(a,1,2)),rewrite([21(7)])]$.
353 $(x \wedge y) \vee (x \wedge (z \vee y)) = x \wedge (z \vee y)$. $[para(149(a,1),24(a,1,2)),rewrite([4(4)])]$.
380 $(x \vee y) \wedge (x \vee (z \wedge y)) = x \vee (z \wedge y)$. $[para(159(a,1),30(a,1,2)),rewrite([2(4)])]$.
403 $x \wedge (y \wedge ((y \wedge x) \vee z)) = x \wedge y$. $[para(2(a,1),29(a,1,2,2,1))]$.

407 $(a \wedge x) \vee (((a \wedge x) \vee y) \wedge z) = ((a \wedge x) \vee z) \wedge ((a \wedge x) \vee y)$. [*para* (29(a,1),13(a,2,1,1)), *rewrite* ([5(9),29(11),4(7),23(17),196(17)])].
424 $(x \wedge y) \vee (z \vee (y \wedge x)) = z \vee (y \wedge x)$. [*para* (213(a,1),5(a,2,2)), *rewrite* ([4(4)])].
428 $x \wedge (y \wedge (z \vee (y \wedge x))) = x \wedge y$. [*para* (213(a,1),143(a,1,2,2)), *rewrite* ([3(4)])].
429 $(x \wedge (y \wedge z)) \vee (y \wedge x) = x \wedge y$. [*para* (213(a,1),26(a,2)), *rewrite* ([3(3),424(6)])].
431 $x \vee ((y \wedge (z \wedge (u \wedge x))) \vee w) = x \vee w$. [*para* (215(a,1),5(a,1,1)), *flip* (a)].
458 $a \wedge (x \wedge ((y \vee z) \wedge (u \vee (w \vee ((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)))) = a \wedge (x \wedge (y \vee z))$. [*para* (13(a,1),232(a,1,2,2,2)), *rewrite* ([3(16),3(15)])].
461 $x \wedge ((y \vee (a \wedge z)) \wedge (u \vee (w \vee (x \wedge (z \vee y)))) = x \wedge (y \vee (a \wedge z))$. [*para* (15(a,1),232(a,1,2,2,2)), *rewrite* ([3(9)])].
483 $(x \vee y) \wedge (y \vee (x \vee z)) = y \vee x$. [*para* (4(a,1),31(a,1,1))].
493 $(x \vee y) \wedge (y \vee x) = x \vee y$. [*para* (108(a,1),31(a,1,2))].
495 $(x \wedge (y \wedge z)) \vee (u \vee (x \wedge y)) = u \vee (x \wedge y)$. [*para* (3(a,1),263(a,1,1))]].
503 $(x \wedge (y \wedge z)) \vee (u \vee y) = u \vee y$. [*para* (21(a,1),263(a,1,1))].
512 $(a \wedge x) \vee (y \wedge (z \vee x)) = ((a \wedge x) \vee y) \wedge (z \vee x)$. [*back_rewrite* (146), *rewrite* ([503(8)]), *flip* (a)].
514 $(x \vee y) \wedge (z \wedge (u \wedge x)) = z \wedge (u \wedge x)$. [*para* (3(a,1),277(a,1,2)), *rewrite* ([3(6)])].
516 $(x \wedge y) \vee (a \wedge (z \wedge (u \vee y))) = (y \vee u) \wedge ((a \wedge (z \wedge (y \vee u))) \vee (x \wedge y))$. [*para* (277(a,1),13(a,1,1)), *rewrite* ([5(5),159(5),23(18),431(18),2(14)])].
534 $(x \wedge (y \wedge z)) \vee (u \vee z) = u \vee z$. [*para* (3(a,1),304(a,1,1))].
566 $((a \wedge (x \wedge y)) \vee z) \wedge (((a \wedge (x \wedge z)) \vee y) \wedge (u \wedge (a \wedge (x \wedge (y \vee z)))) = u \wedge (a \wedge (x \wedge (y \vee z)))$. [*para* (13(a,1),321(a,1,1)), *rewrite* ([3(15)])].
582 $(x \vee y) \wedge (z \wedge (x \vee (u \wedge y))) = z \wedge (x \vee (u \wedge y))$. [*para* (159(a,1),321(a,1,1))].
674 $c1 \vee (c2 \vee (a \wedge c3)) = c1 \vee c2$. [*para* (352(a,1),132(a,1,2)), *rewrite* ([4(7),23(7),4(6)])].
695 $x \vee (y \vee (z \wedge (y \vee x))) = x \vee y$. [*para* (493(a,1),132(a,1,2,2)), *rewrite* ([5(4)])].
698 $(x \vee (y \vee z)) \wedge (z \vee y) = z \vee y$. [*para* (493(a,1),321(a,1,2)), *rewrite* ([493(7)])].
712 $a \wedge (c3 \wedge (x \vee (c1 \vee c2))) = a \wedge c3$. [*para* (674(a,1),232(a,1,2,2)), *rewrite* ([3(8)])].
847 $x \wedge (y \wedge (z \wedge (u \vee (x \wedge z)))) = x \wedge (y \wedge z)$. [*para* (140(a,1),143(a,1,2,2)), *rewrite* ([3(5),3(4)])].
851 $(x \wedge (y \wedge z)) \vee (x \wedge (y \wedge (z \vee u))) = x \wedge (y \wedge (z \vee u))$. [*para* (91(a,1),140(a,1,2,2)), *rewrite* ([4(6)])].
870 $(a \wedge x) \vee (y \wedge (z \vee (a \wedge x))) = (z \vee (a \wedge x)) \wedge ((a \wedge x) \vee y)$. [*para* (142(a,1),13(a,2,2,1)), *rewrite* ([231(11),4(7),495(14)])].
904 $(x \vee y) \wedge (x \vee (z \wedge (u \wedge y))) = x \vee (z \wedge (u \wedge y))$. [*para* (132(a,1),160(a,1,2,2)), *rewrite* ([2(5)])].
1373 $(x \wedge y) \vee z = z \vee (((a \wedge (x \wedge z)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x))$. [*para* (37(a,1),171(a,1,2)), *flip* (a)].

1688 $(x \vee y) \wedge ((z \wedge (u \wedge x)) \vee y) = (z \wedge (u \wedge x)) \vee y$. [para(216(a,1),30(a,1,2)),rewrite([2(5)])].
2786 $(a \wedge x) \vee (y \wedge (z \wedge (x \vee u))) = (x \vee u) \wedge ((a \wedge x) \vee (y \wedge z))$. [para(7(a,1),41(a,2,2,1,2)),rewrite([153(8),4(6),23(12),163(12)])].
4300 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = (y \vee (a \wedge (z \wedge x))) \wedge (x \vee (a \wedge (z \wedge y)))$. [para(4(a,1),44(a,2,2))].
4305 $(x \wedge y) \vee (a \wedge x) = x \wedge (y \vee (a \wedge x))$. [para(7(a,1),44(a,1,2,2)),rewrite([33(6),4(11),90(11),2(8)])].
4488 $x \wedge ((a \wedge y) \vee (a \wedge x)) = a \wedge x$. [para(403(a,1),44(a,1,2)),rewrite([3(3),4305(6),112(10),114(10),429(15),2(12),3(12),428(12)])].
4553 $x \vee ((x \vee (a \wedge (y \wedge z))) \wedge (z \vee (a \wedge (y \wedge x)))) = x \vee (a \wedge (y \wedge z))$. [para(44(a,2),695(a,1,2,2)),rewrite([4300(9),870(13),23(12),216(12)])].
.
5121 $(x \vee y) \wedge ((z \wedge y) \vee x) = (z \wedge y) \vee x$. [para(159(a,1),698(a,1,1))].
5163 $a \wedge (c3 \wedge (c1 \vee (x \vee c2))) = a \wedge c3$. [para(23(a,1),712(a,1,2,2))].
5929 $x \wedge ((a \wedge x) \vee (a \wedge y)) = a \wedge x$. [para(4(a,1),4488(a,1,2))].
6005 $a \wedge (x \wedge (c3 \wedge (c1 \vee (y \vee c2)))) = x \wedge (a \wedge c3)$. [para(5163(a,1),21(a,1,2)),flip(a)].
6223 $x \vee (a \wedge (y \wedge (x \vee z))) = x \vee (a \wedge (y \wedge z))$. [para(483(a,1),58(a,1,1)),rewrite([5(14),407(13),534(8),259(10),483(20),23(15),4(14),851(14),23(12),4300(11),4553(15)])].
6337 $(a \wedge x) \vee (a \wedge ((x \vee (a \wedge y)) \wedge z)) = ((a \wedge x) \vee (a \wedge y)) \wedge (x \vee (a \wedge (z \wedge ((a \wedge x) \vee (a \wedge y))))$. [para(5929(a,1),36(a,1,1)),rewrite([133(9),23(18),301(18),4(23)])].
6572 $c1 \vee (c3 \vee (a \wedge ((c1 \vee c2) \wedge x))) = c1 \vee c3$. [para(83(a,1),90(a,1,2)),rewrite([5(10)])].
6603 $(c1 \vee c3) \wedge (x \wedge a) = (c1 \vee c2) \wedge (x \wedge a)$. [para(83(a,1),207(a,1)),rewrite([207(8)]),flip(a)].
7269 $(x \vee y) \wedge (x \vee (z \wedge (u \wedge (x \vee y)))) = x \vee (z \wedge (u \wedge (x \vee y)))$. [para(217(a,1),160(a,1,2)),rewrite([2(6)])].
7441 $(x \wedge ((a \wedge y) \vee (a \wedge z))) \vee ((z \vee (a \wedge y)) \wedge x) = x \wedge ((a \wedge y) \vee z)$. [para(112(a,1),68(a,1,2,1,2))].
8538 $x \wedge ((x \wedge y) \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$. [para(6(a,1),261(a,1,1))].
8763 $x \vee ((x \vee y) \wedge (x \vee z)) = (x \vee y) \wedge (x \vee z)$. [para(7(a,1),276(a,1,1))].
9151 $(x \wedge (y \wedge (z \vee u))) \vee ((w \wedge (x \wedge (y \wedge z))) \vee v5) = (x \wedge (y \wedge (z \vee u))) \vee v5$. [para(280(a,1),216(a,1,2,1,2))].
9174 $(x \wedge (y \wedge (z \wedge u))) \vee (w \vee (y \wedge (z \wedge (u \vee v5)))) = w \vee (y \wedge (z \wedge (u \vee v5)))$. [para(280(a,1),534(a,1,1,2))].
9221 $(x \wedge (y \wedge z)) \vee (x \wedge (u \vee y)) = x \wedge (u \vee y)$. [para(263(a,1),292(a,1,2,2)),rewrite([263(8)])].
9343 $x \wedge ((y \wedge z) \vee (y \wedge x)) = x \wedge y$. [para(429(a,1),302(a,1,2)),rewrite([2(5),3(5),8538(4),429(8)])].
9373 $x \vee (a \wedge (y \wedge (z \vee x))) = x \vee (a \wedge (y \wedge z))$. [para(302(a,1),58(a,1,1)),rewrite([5(15),297(14),9174(10),302(20),9151(16),4(11),23(12),4300(11),4553(15)])].
9434 $x \wedge ((y \wedge z) \vee (z \wedge x)) = x \wedge z$. [para(2(a,1),9343(a,1,2,1))].
9435 $x \wedge ((y \wedge z) \vee (x \wedge y)) = x \wedge y$. [para(2(a,1),9343(a,1,2,2))].
9874 $x \wedge ((y \wedge z) \vee (x \wedge (u \vee z))) = x \wedge (u \vee z)$. [para(321(a,1),9435(a,1,2,1))].

11144 $x \vee ((x \wedge y) \vee (x \wedge z)) \wedge u = x$. $[para(338(a,1),26(a,1)),rewrite([6(2)]),flip(a)]$.
11205 $x \vee (y \vee ((x \wedge z) \vee (y \wedge u)) \wedge w) = x \vee y$. $[para(338(a,1),262(a,1,2)),rewrite([152(3)]),flip(a)]$.
11222 $x \vee (y \wedge ((x \wedge z) \vee (x \wedge u))) = x$. $[para(2(a,1),11144(a,1,2))]$.
11735 $x \wedge (y \wedge ((x \wedge z) \vee (x \wedge u))) = y \wedge ((x \wedge z) \vee (x \wedge u))$. $[para(11222(a,1),30(a,1,2)),rewrite([2(5)])]$.
11820 $(a \wedge x) \vee (a \wedge ((x \vee (a \wedge y)) \wedge z)) = ((a \wedge x) \vee (a \wedge y)) \wedge (x \vee (z \wedge ((a \wedge x) \vee (a \wedge y))))$. $[back_rewrite(6337),rewrite([11735(2)])]$.
12718 $x \vee ((y \vee x) \wedge (y \vee z)) = x \vee y$. $[para(483(a,1),353(a,1,2)),rewrite([23(5),4(4),8763(4),483(8)])]$.
12799 $x \vee ((y \vee z) \wedge (y \vee x)) = x \vee y$. $[para(2(a,1),12718(a,1,2))]$.
12819 $x \vee (((y \wedge z) \vee x) \wedge (u \vee z)) = x \vee (y \wedge z)$. $[para(304(a,1),12718(a,1,2,2))]$.
12946 $x \vee ((y \vee z) \wedge ((u \wedge z) \vee x)) = x \vee (u \wedge z)$. $[para(304(a,1),12799(a,1,2,1))]$.
13031 $(a \wedge x) \vee (y \wedge ((x \vee z) \wedge u)) = ((a \wedge x) \vee (y \wedge u)) \wedge (x \vee z)$. $[para(7(a,1),92(a,2,1,1,2)),rewrite([5(7),7(8),4(6),23(16),163(16)])]$.
13409 $((a \wedge x) \vee (a \wedge y)) \wedge (x \vee (a \wedge z)) = ((a \wedge x) \vee (a \wedge z)) \wedge (x \vee (y \wedge ((a \wedge x) \vee (a \wedge z))))$. $[back_rewrite(11820),rewrite([13031(9)])]$.
16801 $x \vee (a \wedge (x \vee y)) = x \vee (a \wedge y)$. $[para(115(a,1),26(a,1,2)),rewrite([12946(8)]),flip(a)]$.
16811 $x \vee (a \wedge (y \vee x)) = x \vee (a \wedge y)$. $[para(115(a,1),133(a,1,2)),rewrite([12819(8)]),flip(a)]$.
16931 $(a \wedge x) \vee (a \wedge y) = a \wedge (x \vee y)$. $[para(115(a,2),41(a,1,2,2)),rewrite([9874(10),9221(7),2(10),3(10),30(9),112(7),2(11),29(12),5121(12)]),flip(a)]$.
17283 $a \wedge (x \vee (y \wedge (a \wedge (x \vee z)))) = a \wedge ((x \vee y) \wedge (x \vee (a \wedge z)))$. $[back_rewrite(13409),rewrite([16931(5),3(7),16931(12),16931(15),3(16),7269(15)]),flip(a)]$.
17360 $(x \wedge (a \wedge (y \vee z))) \vee ((z \vee (a \wedge y)) \wedge x) = x \wedge ((a \wedge y) \vee z)$. $[back_rewrite(7441),rewrite([16931(5)])]$.
17661 $c1 \vee (a \wedge c3) = c1 \vee (a \wedge c2)$. $[para(17(a,1),16801(a,1,2)),rewrite([16801(7)]),flip(a)]$.
17662 $a \wedge (x \vee (a \wedge y)) = a \wedge (x \vee y)$. $[para(16801(a,1),30(a,1,2)),rewrite([3(7),380(6)])]$.
17667 $x \vee (a \wedge ((y \wedge x) \vee z)) = x \vee (a \wedge z)$. $[para(16801(a,1),133(a,1,2)),rewrite([133(5)]),flip(a)]$.
17670 $a \wedge (x \wedge (y \vee (a \wedge z))) = x \wedge (a \wedge (y \vee z))$. $[para(16801(a,1),149(a,1,2,2)),rewrite([3(8),582(7)])]$.
17727 $(x \wedge a) \vee (a \wedge y) = a \wedge ((x \wedge a) \vee y)$. $[para(16801(a,1),9434(a,1,2)),rewrite([380(9),2(10)])]$.
17951 $a \wedge (x \vee (y \wedge (a \wedge (x \vee z)))) = a \wedge ((x \vee y) \wedge (x \vee z))$. $[back_rewrite(17283),rewrite([17670(14),21(12)])]$.
18506 $a \wedge (c1 \vee (c2 \vee c3)) = a \wedge (c1 \vee c2)$. $[para(17661(a,1),16811(a,1,2,2)),rewrite([17662(10),16931(9),23(6),4(5),4(14),16931(14),17(12)])]$.
18675 $a \wedge ((x \wedge a) \vee y) = a \wedge (x \vee y)$. $[para(2(a,1),16931(a,1,1)),rewrite([17727(5)])]$.
18680 $(a \wedge x) \vee (y \wedge (a \wedge z)) = a \wedge (x \vee (y \wedge z))$. $[para(21(a,1),16931(a,1,2))]$.

18682 $a^{\wedge}((a^{\wedge}x) \vee y) = a^{\wedge}(x \vee y)$. [para(112(a,1),16931(a,1,1)),
rewrite([16931(5)]),flip(a)].
18698 $a^{\wedge}(x \vee (y^{\wedge}(a^{\wedge}z))) = a^{\wedge}(x \vee (y^{\wedge}z))$. [para(170(a,1),16931(a,
1,2)),rewrite([18680(6)]),flip(a)].
18803 $(x^{\wedge}a) \vee (a^{\wedge}y) = a^{\wedge}(x \vee y)$. [back_rewrite(17727),rewrite([
18675(10)])].
19197 $a^{\wedge}(x \vee (y^{\wedge}(x \vee z))) = a^{\wedge}((x \vee y)^{\wedge}(x \vee z))$. [back_rewrite
(17951),rewrite([18698(7)])].
19661 $a^{\wedge}((c1 \vee (c2 \vee c3))^{\wedge}x) = a^{\wedge}((c1 \vee c2)^{\wedge}x)$. [para(18506(a,1)
,3(a,1,1)),rewrite([3(6)]),flip(a)].
19740 $a^{\wedge}(x^{\wedge}((a^{\wedge}y) \vee z)) = a^{\wedge}(x^{\wedge}(y \vee z))$. [para(68(a,1),18675(a,
2,2)),rewrite([2(7),17670(7),17360(10)])].
19911 $a^{\wedge}(((a^{\wedge}x) \vee y)^{\wedge}z) = a^{\wedge}((x \vee y)^{\wedge}z)$. [para(18682(a,1),3(a,
1,1)),rewrite([3(4)]),flip(a)].
21622 $(a^{\wedge}x) \vee (x^{\wedge}y) = x^{\wedge}((a^{\wedge}x) \vee y)$. [para(6(a,1),174(a,1,2,1)),
rewrite([6(9),2(8)])].
23068 $(x^{\wedge}(y^{\wedge}z)) \vee (x^{\wedge}(z \vee u)) = (z \vee u)^{\wedge}x$. [para(514(a,1),429(a,
1,1))].
25533 $a^{\wedge}(x^{\wedge}((x^{\wedge}y) \vee z)) = a^{\wedge}(x^{\wedge}(y \vee z))$. [para(6(a,1),236(a,
1,2,2,2)),rewrite([1688(7),19740(7)])].
30690 $c1 \vee (a^{\wedge}(c3 \vee ((c1 \vee c2)^{\wedge}x))) = c1 \vee (a^{\wedge}c2)$. [para(6572(a,1)
,16801(a,1,2,2)),rewrite([17(6),16801(7),17662(16)]),flip(a)].
30792 $a^{\wedge}(x \vee ((c1 \vee c2)^{\wedge}y)) = a^{\wedge}((x \vee (c1 \vee c2))^{\wedge}(x \vee y))$. [para
(6603(a,1),267(a,2,1,2,2)),rewrite([2(9),16801(10),17670(11),21(9)
,83(9),18803(10),19197(8),2(15),21(16),112(17),6223(16),83(21),2(24)
,23068(25),2(18),2(19),3(19),904(18),17662(17)]),flip(a)].
30797 $c1 \vee (a^{\wedge}((c1 \vee c2)^{\wedge}(c3 \vee x))) = c1 \vee (a^{\wedge}c2)$. [back_rewrite
(30690),rewrite([30792(9),23(7),4(6),19661(11)])].
50699 $x \vee (a^{\wedge}((x \vee y)^{\wedge}z)) = x \vee (a^{\wedge}(y^{\wedge}z))$. [para(512(a,1),17667(a,
1,2,2)),rewrite([19911(7),9373(6),9373(10)])].
50705 $c1 \vee (a^{\wedge}(c2^{\wedge}(c3 \vee x))) = c1 \vee (a^{\wedge}c2)$. [back_rewrite(30797),
rewrite([50699(10)])].
50709 $c1 \vee (a^{\wedge}(c2^{\wedge}c3)) = c1 \vee (a^{\wedge}c2)$. [para(6(a,1),50705(a,1,2,2,2)
)].
50727 $c1 \vee (c2^{\wedge}(c3 \vee (a^{\wedge}c2))) = c1 \vee (c2^{\wedge}c3)$. [para(50709(a,1),159(
a,1,2)),rewrite([23(9),4(8),21622(8),4(7)])].
50801 $a^{\wedge}(x \vee (y^{\wedge}z)) = a^{\wedge}((x \vee z)^{\wedge}(x \vee y))$. [para(516(a,1),18682(a,
1,2)),rewrite([4(9),2786(9),16931(8),21(7),21(8),112(7),112(7)
,9373(11),17662(11)]),flip(a)].
50871 $a^{\wedge}(c2^{\wedge}(x \vee (c1 \vee (c2^{\wedge}c3)))) = a^{\wedge}c2$. [para(50727(a,1),461(a,
1,2,2,2)),rewrite([16931(8),3(13),21(14),28(13),16931(18),21(17),7(16)
])].
50874 $a^{\wedge}(c2^{\wedge}(c1 \vee ((c2^{\wedge}c3) \vee x))) = a^{\wedge}c2$. [para(4(a,1),50871(a,
1,2,2)),rewrite([5(8)])].
50901 $a^{\wedge}(c2^{\wedge}(c3 \vee (x \vee c1))) = a^{\wedge}c2$. [para(185(a,1),50874(a,1,2,2))
,rewrite([25533(10)])].
53798 $a^{\wedge}(c3^{\wedge}((c1^{\wedge}c3) \vee (x \vee c2))) = a^{\wedge}c3$. [para(6005(a,1),566(a,
1,2,2)),rewrite([2(4),4(16),112(21),2(20),3(20),17670(20),50801(17)
,2(18),83(19),160(18),21(15),352(15),2(12),3(12),19740(12),5163(19)
,112(15)])].
53820 $a^{\wedge}(c3^{\wedge}(c2 \vee (c1^{\wedge}c3))) = a^{\wedge}c3$. [para(32(a,1),53798(a,
1,2,2,2)),rewrite([4(7)])].

54317 $c3 \vee (a \wedge (c2 \wedge (x \vee c1))) = c3 \vee (a \wedge c2)$. [para(50901(a,1),6223(a,1,2)),flip(a)].
54322 $c2 \vee (a \wedge (c1 \wedge c3)) = c2 \vee (a \wedge c3)$. [para(53820(a,1),6223(a,1,2)),rewrite([114(12)]),flip(a)].
54473 $c3 \vee (a \wedge (c1 \wedge c2)) = c3 \vee (a \wedge c2)$. [para(32(a,1),54317(a,1,2,2,2)),rewrite([2(5)])].
60183 $a \wedge (x \wedge (y \vee (z \wedge u))) = x \wedge (a \wedge ((y \vee u) \wedge (y \vee z)))$. [para(11205(a,1),458(a,1,2,2,2)),rewrite([280(8),2(4),50801(4)]),flip(a)].
73833 $(c1 \wedge c3) \vee (c1 \wedge (c2 \vee (a \wedge c3))) \neq c1 \wedge (c2 \vee c3)$. [para(1373(a,1),19(a,1)),rewrite([112(9),4(10),54322(10),2(14),21(14),4(17),90(17),2(10)])].
73856 \$F. [para(1373(a,1),73833(a,1)),rewrite([112(17),60183(16),4(15),2(16),28(17),21(14),4(16),9373(16),54473(14),2(22),21(22),847(23),4(19),90(19),2(14),15(15)]),xx(a)].

===== **end of proof**=====

===== **STATISTICS**=====

Given=1611. Generated=4107893. Kept=73850. proofs=1.
Usable=1438. Sos=19999. Demods=19736. Limbo=0, Disabled=52427. Hints=0.
Kept_by_rule=0, Deleted_by_rule=5024.
Forward_subsumed=2472886. Back_subsumed=870.
Sos_limit_deleted=1556132. Sos_displaced=38067. Sos_removed=0.
New_demodulators=64146 (6 lex), Back_demodulated=13474. Back_unit_deleted=0.
Demod_attempts=108087980. Demod_rewrites=13862951.
Res_instance_prunes=0. Para_instance_prunes=0. Basic_paramod_prunes=0.
Nonunit_fsub_feature_tests=0. Nonunit_bsub_feature_tests=0.
Megabytes=107.30.
User_CPU=222.55, System_CPU=1.51, Wall_clock=391.

===== **end of statistics**=====

===== **end of search**=====

THEOREM PROVED

Just for comparison, see how much simpler the proof gets when AxRL1 is assumed:

formulas (assumptions).

$x \wedge y = y \wedge x$.
 $(x \wedge y) \wedge z = x \wedge (y \wedge z)$.
 $x \vee y = y \vee x$.
 $(x \vee y) \vee z = x \vee (y \vee z)$.
 $x \vee (x \wedge y) = x$.
 $x \wedge (x \vee y) = x$.
UpMe(x,y,z) = $x \wedge (y \vee z)$.
LoMe(x,y,z) = $(x \wedge y) \vee (x \wedge z)$.
UpJo(x,y,z) = $(x \vee y) \wedge (x \vee z)$.
LoJo(x,y,z) = $x \vee (y \wedge z)$.
UpMe(a \wedge x1,y1,z1) \vee (y1 \wedge z1) = $((a \wedge x1) \wedge y1) \vee z1) \wedge ((a \wedge x1) \wedge z1) \vee y1)$.

```

LoMe(x,y,z) = UpMe(x,UpMe(y,x,z),UpMe(z,x,y)).
UpMe(x,y,z) = UpMe(x,y,a ^ z) v UpMe(x,z,a ^ y).
end_of_list .

```

```

formulas(goals) .

```

```

(all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> UpMe(x2,y2,z2) =
  LoMe(x2,y2,z2))).
end_of_list .

```

```

.....

```

```

===== PROOF =====

```

```

% Proof 1 at 88.76 (+ 0.59) seconds.

```

```

% Length of proof is 38.

```

```

% Level of proof is 8.

```

```

% Maximum clause weight is 29.

```

```

% Given clauses 936.

```

```

1 (all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> UpMe(x2,y2,z2) =
  LoMe(x2,y2,z2))) # label(non_clause) # label(goal). [goal].
2 x ^ y = y ^ x. [assumption].
3 (x ^ y) ^ z = x ^ (y ^ z). [assumption].
4 x v y = y v x. [assumption].
5 (x v y) v z = x v (y v z). [assumption].
6 x v (x ^ y) = x. [assumption].
7 x ^ (x v y) = x. [assumption].
8 UpMe(x,y,z) = x ^ (y v z). [assumption].
9 LoMe(x,y,z) = (x ^ y) v (x ^ z). [assumption].
12 UpMe(a ^ x,y,z) v (y ^ z) = (((a ^ x) ^ y) v z) ^ (((a ^ x) ^ z) v y).
  [assumption].
13 (x ^ y) v (a ^ (z ^ (x v y))) = ((a ^ (z ^ x)) v y) ^ ((a ^ (z ^ y)) v
  x). [copy(12),rewrite([8(3),3(4),4(6),3(9),3(13)])].
14 LoMe(x,y,z) = UpMe(x,UpMe(y,x,z),UpMe(z,x,y)). [assumption].
15 (x ^ y) v (x ^ z) = x ^ ((y ^ (x v z)) v (z ^ (x v y))). [copy(14),
  rewrite([9(1),8(4),8(6),8(8)])].
16 UpMe(x,y,z) = UpMe(x,y,a ^ z) v UpMe(x,z,a ^ y). [assumption].
17 (x ^ (y v (a ^ z))) v (x ^ (z v (a ^ y))) = x ^ (y v z). [copy(16),
  rewrite([8(1),8(5),8(9)]),flip(a)].
18 UpMe(a,c1,c3) = UpMe(a,c1,c2). [deny(1)].
19 a ^ (c1 v c3) = a ^ (c1 v c2). [copy(18),rewrite([8(4),8(9)])].
20 LoMe(c1,c2,c3) != UpMe(c1,c2,c3). [deny(1)].
21 (c1 ^ c2) v (c1 ^ c3) != c1 ^ (c2 v c3). [copy(20),rewrite([9(4),8(11)
  ])].
23 x ^ (y ^ z) = y ^ (x ^ z). [para(2(a,1),3(a,1,1)),rewrite([3(2)])].
28 x v ((x ^ y) v z) = x v z. [para(6(a,1),5(a,1,1)),flip(a)].
30 x ^ ((x v y) ^ z) = x ^ z. [para(7(a,1),3(a,1,1)),flip(a)].
32 x ^ (y v x) = x. [para(4(a,1),7(a,1,2))].
34 x v x = x. [para(7(a,1),6(a,1,2))].
67 x ^ ((y ^ (x v z)) v (z ^ (x v y))) = (y ^ x) v (x ^ z). [para(2(a,1)
  ,15(a,1,1)),flip(a)].

```


132 $a \wedge (x \wedge (c1 \vee c3)) = x \wedge (a \wedge (c1 \vee c2))$. $[para(19(a,1),23(a,1,2)),$
 $flip(a)]$.
143 $x \vee (y \vee x) = y \vee x$. $[para(34(a,1),5(a,2,2)),rewrite([4(2)])]$.
183 $x \wedge (y \vee (z \vee x)) = x$. $[para(5(a,1),32(a,1,2))]$.
186 $((a \wedge x) \vee y) \wedge ((a \wedge (x \wedge y)) \vee (z \vee x)) = (a \wedge x) \vee (y \wedge (z \vee x))$.
 $[para(32(a,1),13(a,2,2,1,2)),rewrite([183(6),4(5),2(14)]),flip(a)]$.
189 $x \wedge (y \wedge (z \vee x)) = y \wedge x$. $[para(32(a,1),23(a,1,2)),flip(a)]$.
231 $x \vee (y \vee (z \vee x)) = y \vee (z \vee x)$. $[para(5(a,1),143(a,1,2)),rewrite([$
 $5(5)])]$.
312 $(x \wedge y) \vee (z \vee x) = z \vee x$. $[para(143(a,1),28(a,2)),rewrite([231(4)])]$
.
401 $a \wedge (c3 \wedge (c1 \vee c2)) = a \wedge c3$. $[para(19(a,1),189(a,1,2)),rewrite([$
 $23(7)])]$.
553 $(x \wedge (y \wedge z)) \vee (u \vee y) = u \vee y$. $[para(23(a,1),312(a,1,1))]$.
562 $(a \wedge x) \vee (y \wedge (z \vee x)) = ((a \wedge x) \vee y) \wedge (z \vee x)$. $[back_rewrite(186)$
 $,rewrite([553(8)]),flip(a)]$.
721 $(x \wedge (y \vee (a \wedge c3))) \vee (x \wedge ((c3 \wedge (c1 \vee c2)) \vee (a \wedge y))) = x \wedge (y \vee ($
 $c3 \wedge (c1 \vee c2)))$. $[para(401(a,1),17(a,1,1,2,2))]$.
51633 $(c1 \wedge c2) \vee (c1 \wedge c3) = c1 \wedge (c2 \vee c3)$. $[para(67(a,1),721(a,2)),$
 $rewrite([4(10),562(10),4(6),2(10),30(11),132(20),23(20),32(19),4(17)$
 $,562(17),4(13),2(17),30(18),17(15),2(8)]),flip(a)]$.
51634 \$F. $[resolve(51633,a,21,a)]$.

===== **end of proof**=====

===== **STATISTICS**=====

Given=936. Generated=1315923. Kept=51627. proofs=1.
Usable=861. Sos=19999. Demods=18987. Limbo=1, Disabled=30780. Hints=0.
Kept_by_rule=0, Deleted_by_rule=3409.
Forward_subsumed=884193. Back_subsumed=793.
Sos_limit_deleted=376694. Sos_displaced=18619. Sos_removed=0.
New_demodulators=43355 (6 lex), Back_demodulated=11351. Back_unit_deleted
=0.
Demod_attempts=32232244. Demod_rewrites=4313279.
Res_instance_prunes=0. Para_instance_prunes=0. Basic_paramod_prunes=0.
Nonunit_fsub_feature_tests=0. Nonunit_bsub_feature_tests=0.
Megabytes=87.32.
User_CPU=88.76, System_CPU=0.59, Wall_clock=143.

===== **end of statistics**=====

===== **end of search**=====

THEOREM PROVED

Appendix C. Theorem 3.4, Clause 7:

formulas(assumptions).

```

x ^ y = y ^ x.
(x ^ y) ^ z = x ^ (y ^ z).
x v y = y v x.
(x v y) v z = x v (y v z).
x v (x ^ y) = x.
x ^ (x v y) = x.
UpMe(x,y,z) = x ^ (y v z).
LoMe(x,y,z) = (x ^ y) v (x ^ z).
UpJo(x,y,z) = (x v y) ^ (x v z).
LoJo(x,y,z) = x v (y ^ z).
UpMe(a ^ x1,y1,z1) v (y1 ^ z1) = (((a ^ x1) ^ y1) v z1) ^ (((a ^ x1) ^ z1)
v y1).
UpMe(x,y,z) = UpMe(x,y,a ^ z) v UpMe(x,z,a ^ y).
end_of_list.

```

formulas(goals).

```

(all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> (UpMe(a,x2,y2) =
UpMe(a,y2,z2) ->
UpJo(x2,y2,z2) = LoJo(x2,y2,z2))))).
end_of_list.

```

.....

===== PROOF =====

```

% Proof 1 at 963.33 (+ 15.06) seconds.
% Length of proof is 196.
% Level of proof is 20.
% Maximum clause weight is 47.000.
% Given clauses 4826.

```

```

1 (all x2 all y2 all z2 (UpMe(a,x2,y2) = UpMe(a,x2,z2) -> (UpMe(a,x2,y2) =
UpMe(a,y2,z2) -> UpJo(x2,y2,z2) = LoJo(x2,y2,z2)))) # label(non_clause
) # label(goal). [goal].
2 x ^ y = y ^ x. [assumption].
3 (x ^ y) ^ z = x ^ (y ^ z). [assumption].
4 x v y = y v x. [assumption].
5 (x v y) v z = x v (y v z). [assumption].
6 x v (x ^ y) = x. [assumption].
7 x ^ (x v y) = x. [assumption].
8 UpMe(x,y,z) = x ^ (y v z). [assumption].
9 LoMe(x,y,z) = (x ^ y) v (x ^ z). [assumption].
10 UpJo(x,y,z) = (x v y) ^ (x v z). [assumption].
11 LoJo(x,y,z) = x v (y ^ z). [assumption].

```

12 $\text{UpMe}(a \wedge x, y, z) \vee (y \wedge z) = (((a \wedge x) \wedge y) \vee z) \wedge (((a \wedge x) \wedge z) \vee y)$.
[assumption].

13 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = ((a \wedge (z \wedge x)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x)$. *[copy(12), rewrite([8(3), 3(4), 4(6), 3(9), 3(13)])]*.

14 $\text{UpMe}(x, y, z) = \text{UpMe}(x, y, a \wedge z) \vee \text{UpMe}(x, z, a \wedge y)$. *[assumption]*.

15 $(x \wedge (y \vee (a \wedge z))) \vee (x \wedge (z \vee (a \wedge y))) = x \wedge (y \vee z)$. *[copy(14), rewrite([8(1), 8(5), 8(9)]), flip(a)]*.

16 $\text{UpMe}(a, c1, c3) = \text{UpMe}(a, c1, c2)$. *[deny(1)]*.

17 $a \wedge (c1 \vee c3) = a \wedge (c1 \vee c2)$. *[copy(16), rewrite([8(4), 8(9)])]*.

18 $\text{UpMe}(a, c2, c3) = \text{UpMe}(a, c1, c2)$. *[deny(1)]*.

19 $a \wedge (c2 \vee c3) = a \wedge (c1 \vee c2)$. *[copy(18), rewrite([8(4), 8(9)])]*.

20 $\text{LoJo}(c1, c2, c3) \neq \text{UpJo}(c1, c2, c3)$. *[deny(1)]*.

21 $c1 \vee (c2 \wedge c3) \neq (c1 \vee c2) \wedge (c1 \vee c3)$. *[copy(20), rewrite([11(4), 10(9)])]*.

23 $x \wedge (y \wedge z) = y \wedge (x \wedge z)$. *[para(2(a,1), 3(a,1,1)), rewrite([3(2)])]*.

25 $x \vee (y \vee z) = y \vee (x \vee z)$. *[para(4(a,1), 5(a,1,1)), rewrite([5(2)])]*.

26 $x \vee (y \wedge x) = x$. *[para(2(a,1), 6(a,1,2))]*.

27 $(x \wedge y) \vee (x \wedge (y \wedge z)) = x \wedge y$. *[para(3(a,1), 6(a,1,2))]*.

28 $x \vee ((x \wedge y) \vee z) = x \vee z$. *[para(6(a,1), 5(a,1,1)), flip(a)]*.

29 $x \vee (y \vee ((x \vee y) \wedge z)) = x \vee y$. *[para(6(a,1), 5(a,1,1)), flip(a)]*.

30 $x \wedge ((x \vee y) \wedge z) = x \wedge z$. *[para(7(a,1), 3(a,1,1)), flip(a)]*.

31 $x \wedge (y \wedge ((x \wedge y) \vee z)) = x \wedge y$. *[para(7(a,1), 3(a,1,1)), flip(a)]*.

32 $x \wedge (y \vee x) = x$. *[para(4(a,1), 7(a,1,2))]*.

33 $(x \vee y) \wedge (x \vee (y \vee z)) = x \vee y$. *[para(5(a,1), 7(a,1,2))]*.

34 $x \vee x = x$. *[para(7(a,1), 6(a,1,2))]*.

35 $x \wedge x = x$. *[para(6(a,1), 7(a,1,2))]*.

38 $(x \wedge y) \vee (a \wedge ((x \vee y) \wedge z)) = ((a \wedge (z \wedge x)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x)$. *[para(2(a,1), 13(a,1,2,2))]*.

39 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = ((a \wedge (x \wedge z)) \vee y) \wedge ((a \wedge (z \wedge y)) \vee x)$. *[para(2(a,1), 13(a,2,1,1,2))]*.

43 $(x \wedge (y \wedge z)) \vee (a \wedge (u \wedge ((x \wedge y) \vee z))) = ((a \wedge (u \wedge (x \wedge y))) \vee z) \wedge ((a \wedge (u \wedge z)) \vee (x \wedge y))$. *[para(3(a,1), 13(a,1,1))]*.

46 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = (y \vee (a \wedge (z \wedge x))) \wedge ((a \wedge (z \wedge y)) \vee x)$. *[para(4(a,1), 13(a,2,1))]*.

57 $((a \wedge x) \vee y) \wedge (x \vee ((a \wedge (x \wedge y)) \vee z)) = (a \wedge x) \vee ((x \vee z) \wedge y)$. *[para(7(a,1), 13(a,2,1,1,2)), rewrite([5(5), 7(6), 4(5), 25(13)]), flip(a)]*.

60 $((a \wedge (x \wedge y)) \vee z) \wedge u \vee (((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)) = (y \wedge z) \vee ((a \wedge (x \wedge (y \vee z))) \vee (((a \wedge (x \wedge y)) \vee z) \wedge u))$. *[para(13(a,2), 9(a,2,2)), rewrite([9(9), 25(27), 4(26)])]*.

67 $(x \wedge (y \vee (z \wedge a))) \vee (x \wedge (z \vee (a \wedge y))) = x \wedge (y \vee z)$. *[para(2(a,1), 15(a,1,1,2,2))]*.

86 $(a \wedge (c1 \vee c2)) \vee (a \wedge (x \wedge (a \vee (c1 \vee c3)))) = a \wedge (c1 \vee (c3 \vee (a \wedge (x \wedge a))))$. *[para(17(a,1), 13(a,1,1)), rewrite([25(22), 4(21), 4(30), 6(30), 2(24)])]*.

91 $(x \wedge (c1 \vee (c3 \vee (a \wedge y)))) \vee (x \wedge (y \vee (a \wedge (c1 \vee c2)))) = x \wedge (c1 \vee (c3 \vee y))$. *[para(17(a,1), 15(a,1,2,2,2)), rewrite([5(6), 5(19)])]*.

93 $(a \wedge (c1 \vee c2)) \vee (a \wedge (x \wedge (a \vee (c2 \vee c3)))) = a \wedge (c2 \vee (c3 \vee (a \wedge (x \wedge a))))$. *[para(19(a,1), 13(a,1,1)), rewrite([25(22), 4(21), 4(30), 6(30), 2(24)])]*.

98 $(x \wedge (c2 \vee (c3 \vee (a \wedge y)))) \vee (x \wedge (y \vee (a \wedge (c1 \vee c2)))) = x \wedge (c2 \vee (c3 \vee y))$. *[para(19(a,1), 15(a,1,2,2,2)), rewrite([5(6), 5(19)])]*.

99 $x \vee (y \wedge (x \wedge z)) = x$. *[para(23(a,1), 6(a,1,2))]*.

100 $x \wedge (y \wedge (x \vee z)) = y \wedge x$. $[para(7(a,1),23(a,1,2)), flip(a)]$.
110 $a \wedge (x \wedge (c1 \vee c3)) = x \wedge (a \wedge (c1 \vee c2))$. $[para(17(a,1),23(a,1,2)), flip(a)]$.
111 $a \wedge (x \wedge (c2 \vee c3)) = x \wedge (a \wedge (c1 \vee c2))$. $[para(19(a,1),23(a,1,2)), flip(a)]$.
113 $(a \wedge (c1 \vee c2)) \vee (x \wedge a) = a \wedge (c2 \vee (c3 \vee (a \wedge (x \wedge a))))$. $[back_rewrite(93), rewrite([100(13)])]$.
114 $a \wedge (c2 \vee (c3 \vee (a \wedge (x \wedge a)))) = a \wedge (c1 \vee (c3 \vee (a \wedge (x \wedge a))))$. $[back_rewrite(86), rewrite([100(13),113(8)])]$.
117 $(a \wedge (c1 \vee c2)) \vee (x \wedge a) = a \wedge (c1 \vee (c3 \vee (a \wedge (x \wedge a))))$. $[back_rewrite(113), rewrite([114(18)])]$.
118 $x \vee (x \vee y) = x \vee y$. $[para(34(a,1),5(a,1,1)), flip(a)]$.
120 $x \vee (y \vee x) = y \vee x$. $[para(34(a,1),5(a,2,2)), rewrite([4(2)])]$.
124 $x \wedge (x \wedge y) = x \wedge y$. $[para(35(a,1),3(a,1,1)), flip(a)]$.
126 $x \wedge (y \wedge x) = y \wedge x$. $[para(35(a,1),3(a,2,2)), rewrite([2(2)])]$.
127 $(x \wedge y) \vee (a \wedge (x \vee y)) = ((a \wedge x) \vee y) \wedge ((a \wedge y) \vee x)$. $[para(35(a,1),13(a,1,2,2)), rewrite([2(8),7(8),2(11),32(11)])]$.
128 $(x \wedge y) \vee (a \wedge y) = y \wedge ((a \wedge y) \vee x)$. $[para(35(a,1),13(a,2,2,1,2)), rewrite([32(4),4(8),99(8)])]$.
144 $(a \wedge (c1 \vee c2)) \vee (x \wedge a) = a \wedge (c1 \vee (c3 \vee (x \wedge a)))$. $[back_rewrite(117), rewrite([126(15)])]$.
148 $x \vee (y \wedge (z \wedge x)) = x$. $[para(3(a,1),26(a,1,2))]$.
149 $x \vee ((y \wedge x) \vee z) = x \vee z$. $[para(26(a,1),5(a,1,1)), flip(a)]$.
150 $x \vee (y \vee (z \wedge (x \vee y))) = x \vee y$. $[para(26(a,1),5(a,1)), flip(a)]$.
157 $(x \wedge y) \vee (x \wedge (z \wedge y)) = x \wedge y$. $[para(23(a,1),26(a,1,2))]$.
158 $x \wedge ((y \vee x) \wedge z) = x \wedge z$. $[para(32(a,1),3(a,1,1)), flip(a)]$.
159 $x \wedge (y \wedge (z \vee (x \wedge y))) = x \wedge y$. $[para(32(a,1),3(a,1)), flip(a)]$.
160 $x \wedge (y \vee (z \vee x)) = x$. $[para(5(a,1),32(a,1,2))]$.
166 $x \wedge (y \wedge (z \vee x)) = y \wedge x$. $[para(32(a,1),23(a,1,2)), flip(a)]$.
169 $x \vee (y \vee (x \wedge z)) = y \vee x$. $[para(6(a,1),25(a,1,2)), flip(a)]$.
170 $x \wedge (y \vee (x \vee z)) = x$. $[para(25(a,1),7(a,1,2))]$.
176 $x \vee (y \vee (z \wedge x)) = y \vee x$. $[para(26(a,1),25(a,1,2)), flip(a)]$.
177 $(x \vee y) \wedge (x \vee (z \vee y)) = x \vee y$. $[para(25(a,1),32(a,1,2))]$.
180 $x \vee ((y \wedge (x \wedge z)) \vee u) = x \vee u$. $[para(99(a,1),5(a,1,1)), flip(a)]$.
191 $(a \wedge x) \vee ((x \vee y) \wedge z) = ((a \wedge x) \vee z) \wedge (x \vee y)$. $[back_rewrite(57), rewrite([180(8)]), flip(a)]$.
202 $x \vee (y \vee (z \vee x)) = y \vee (z \vee x)$. $[para(5(a,1),120(a,1,2)), rewrite([5(5)])]$.
213 $(x \wedge y) \vee ((x \wedge (y \wedge z)) \vee u) = (x \wedge y) \vee u$. $[para(27(a,1),5(a,1,1)), flip(a)]$.
224 $x \wedge (y \wedge (z \wedge x)) = y \wedge (z \wedge x)$. $[para(3(a,1),126(a,1,2)), rewrite([3(5)])]$.
230 $(x \wedge y) \vee (y \wedge x) = x \wedge y$. $[para(126(a,1),27(a,1,2))]$.
232 $x \vee (y \wedge (z \wedge (u \wedge x))) = x$. $[para(3(a,1),148(a,1,2,2))]$.
233 $x \vee ((y \wedge (z \wedge x)) \vee u) = x \vee u$. $[para(148(a,1),5(a,1,1)), flip(a)]$.
239 $c1 \vee (c3 \vee (x \wedge (a \wedge (c1 \vee c2)))) = c1 \vee c3$. $[para(17(a,1),148(a,1,2,2)), rewrite([5(10)])]$.
249 $x \wedge (y \wedge (z \vee (u \vee (x \wedge y)))) = x \wedge y$. $[para(160(a,1),3(a,1)), flip(a)]$.
250 $x \wedge (y \vee (z \vee (u \vee x))) = x$. $[para(5(a,1),160(a,1,2,2))]$.
267 $(a \wedge x) \vee (y \wedge (x \vee z)) = (x \vee z) \wedge ((a \wedge x) \vee y)$. $[para(170(a,1),13(a,1,2,2)), rewrite([4(5),25(10),180(10),7(9)])]$.

277 $x \vee (((a \wedge (y \wedge x)) \vee z) \wedge ((a \wedge (y \wedge z)) \vee x)) = x \vee (a \wedge (y \wedge (x \vee z)))$. $[para(13(a,1),28(a,1,2))]$.
279 $(x \vee y) \wedge ((x \wedge z) \vee y) = (x \wedge z) \vee y$. $[para(28(a,1),32(a,1,2)), rewrite([2(4)])]$.
280 $x \vee (y \vee ((x \wedge z) \vee u)) = y \vee (x \vee u)$. $[para(28(a,1),25(a,1,2)), flip(a)]$.
281 $(x \wedge y) \vee (z \vee x) = z \vee x$. $[para(120(a,1),28(a,2)), rewrite([202(4)])]$.
293 $x \wedge (y \wedge ((x \vee z) \wedge u)) = y \wedge (x \wedge u)$. $[para(30(a,1),23(a,1,2)), flip(a)]$.
294 $(x \wedge y) \vee ((x \vee z) \wedge y) = (x \vee z) \wedge y$. $[para(30(a,1),26(a,1,2)), rewrite([4(4)])]$.
295 $(x \vee y) \wedge (z \wedge x) = z \wedge x$. $[para(126(a,1),30(a,2)), rewrite([224(4)])]$.
296 $((x \vee y) \wedge z) \vee (u \wedge (x \wedge z)) = (x \vee y) \wedge z$. $[para(30(a,1),148(a,1,2,2))]$.
298 $x \wedge (y \wedge (z \wedge (x \vee u))) = y \wedge (z \wedge x)$. $[para(3(a,1),100(a,1,2)), rewrite([3(6)])]$.
310 $(x \wedge y) \vee (x \wedge (y \vee z)) = x \wedge (y \vee z)$. $[para(100(a,1),26(a,1,2)), rewrite([4(4)])]$.
315 $x \vee (((a \wedge (y \wedge z)) \vee x) \wedge ((a \wedge (y \wedge x)) \vee z)) = x \vee (a \wedge (y \wedge (z \vee x)))$. $[para(13(a,1),149(a,1,2))]$.
321 $(x \vee y) \wedge ((z \wedge x) \vee y) = (z \wedge x) \vee y$. $[para(149(a,1),32(a,1,2)), rewrite([2(4)])]$.
322 $x \vee (y \vee ((z \wedge x) \vee u)) = y \vee (x \vee u)$. $[para(149(a,1),25(a,1,2)), flip(a)]$.
323 $(x \wedge y) \vee (z \vee y) = z \vee y$. $[para(120(a,1),149(a,2)), rewrite([202(4)])]$.
338 $(x \wedge y) \vee ((z \vee x) \wedge y) = (z \vee x) \wedge y$. $[para(158(a,1),26(a,1,2)), rewrite([4(4)])]$.
340 $(x \vee y) \wedge (z \wedge y) = z \wedge y$. $[para(126(a,1),158(a,2)), rewrite([224(4)])]$.
357 $x \vee (y \vee (((x \wedge z) \vee y) \wedge u)) = x \vee y$. $[para(29(a,1),28(a,1,2)), rewrite([28(3)]), flip(a)]$.
372 $(x \wedge y) \vee (x \wedge (z \vee y)) = x \wedge (z \vee y)$. $[para(166(a,1),26(a,1,2)), rewrite([4(4)])]$.
390 $x \vee (y \vee (z \vee (u \wedge (x \vee y)))) = z \vee (x \vee y)$. $[para(176(a,1),5(a,1)), flip(a)]$.
391 $x \vee (y \vee (z \vee (u \wedge x))) = y \vee (z \vee x)$. $[para(5(a,1),176(a,1,2)), rewrite([5(6)])]$.
400 $(x \vee y) \wedge (x \vee (z \wedge y)) = x \vee (z \wedge y)$. $[para(176(a,1),32(a,1,2)), rewrite([2(4)])]$.
427 $(a \wedge x) \vee (((a \wedge x) \vee y) \wedge z) = ((a \wedge x) \vee z) \wedge ((a \wedge x) \vee y)$. $[para(31(a,1),13(a,2,1,1)), rewrite([5(9),31(11),4(7),25(17),213(17)])]$.
444 $(x \wedge y) \vee (z \vee (y \wedge x)) = z \vee (y \wedge x)$. $[para(230(a,1),5(a,2,2)), rewrite([4(4)])]$.
449 $(x \wedge (y \wedge z)) \vee (y \wedge x) = x \wedge y$. $[para(230(a,1),28(a,2)), rewrite([3(3),444(6)])]$.
451 $x \vee ((y \wedge (z \wedge (u \wedge x))) \vee w) = x \vee w$. $[para(232(a,1),5(a,1,1)), flip(a)]$.
479 $a \wedge (x \wedge ((y \vee z) \wedge (u \vee (w \vee (((a \wedge (x \wedge y)) \vee z) \wedge ((a \wedge (x \wedge z)) \vee y)))))) = a \wedge (x \wedge (y \vee z))$. $[para(13(a,1),250(a,1,2,2,2)), rewrite([$

$3(16), 3(15)]]$.
504 $(x \vee y) \wedge (y \vee (x \vee z)) = y \vee x$. [para(4(a,1),33(a,1,1))].
514 $(x \vee y) \wedge (y \vee x) = x \vee y$. [para(120(a,1),33(a,1,2))].
516 $(x \wedge (y \wedge z)) \vee (u \vee (x \wedge y)) = u \vee (x \wedge y)$. [para(3(a,1),281(a,1,1))].
537 $(x \wedge y) \vee (a \wedge (z \wedge (u \vee y))) = (y \vee u) \wedge ((a \wedge (z \wedge (y \vee u))) \vee (x \wedge y))$. [para(295(a,1),13(a,1,1)),rewrite([5(5),176(5),25(18),451(18),2(14)])].
556 $(x \wedge (y \wedge z)) \vee (u \vee z) = u \vee z$. [para(3(a,1),323(a,1,1))].
557 $(x \wedge y) \vee (z \vee (u \vee y)) = z \vee (u \vee y)$. [para(5(a,1),323(a,1,2)),rewrite([5(6)])].
606 $(x \vee y) \wedge (z \wedge (x \vee (u \wedge y))) = z \wedge (x \vee (u \wedge y))$. [para(176(a,1),340(a,1,1))].
719 $x \vee (y \vee (z \wedge (y \vee x))) = x \vee y$. [para(514(a,1),148(a,1,2,2)),rewrite([5(4)])].
722 $(x \vee (y \vee z)) \wedge (z \vee y) = z \vee y$. [para(514(a,1),340(a,1,2)),rewrite([514(7)])].
745 $(x \vee y) \wedge (y \vee (z \wedge (x \vee y))) = y \vee (z \wedge (x \vee y))$. [para(150(a,1),32(a,1,2)),rewrite([2(5)])].
890 $(x \wedge (y \wedge z)) \vee (x \wedge (y \wedge (z \vee u))) = x \wedge (y \wedge (z \vee u))$. [para(100(a,1),157(a,1,2,2)),rewrite([4(6)])].
898 $(x \wedge (y \vee z)) \vee (x \wedge (y \vee (z \vee u))) = x \wedge (y \vee (z \vee u))$. [para(33(a,1),157(a,1,2,2)),rewrite([4(6)])].
909 $(a \wedge x) \vee (y \wedge (z \vee (a \wedge x))) = (z \vee (a \wedge x)) \wedge ((a \wedge x) \vee y)$. [para(159(a,1),13(a,2,2,1)),rewrite([249(11),4(7),516(14)])].
921 $(x \wedge y) \vee (z \vee (y \wedge (u \vee (x \wedge y)))) = z \vee (y \wedge (u \vee (x \wedge y)))$. [para(159(a,1),323(a,1,1))].
941 $(a \wedge ((x \vee y) \wedge z)) \vee ((a \wedge (z \wedge x)) \vee y) = (a \wedge (z \wedge x)) \vee y$. [para(38(a,2),6(a,1,2)),rewrite([25(11),4(10),557(11)])].
964 $(a \wedge (c1 \vee c2)) \vee (a \wedge x) = a \wedge (c1 \vee (c3 \vee (x \wedge a)))$. [para(17(a,1),38(a,1,1)),rewrite([30(13),126(12),25(14),4(13),110(20),4(22),99(22),2(16)])].
1108 $(x \vee y) \wedge (x \vee ((x \vee y) \wedge z)) = x \vee ((x \vee y) \wedge z)$. [para(29(a,1),177(a,1,2)),rewrite([2(5)])].
1243 $a \wedge (c2 \vee (c3 \vee (a \wedge x))) = a \wedge (c1 \vee (c3 \vee (x \wedge a)))$. [para(19(a,1),39(a,1,1)),rewrite([100(13),144(8),124(12),25(14),4(13),111(20),4(22),99(22),2(16)]),flip(a)].
2810 $(a \wedge x) \vee (y \wedge (z \wedge (x \vee u))) = (x \vee u) \wedge ((a \wedge x) \vee (y \wedge z))$. [para(7(a,1),43(a,2,2,1,2)),rewrite([170(8),4(6),25(12),180(12)])].
4331 $(x \wedge y) \vee (a \wedge (z \wedge (x \vee y))) = (y \vee (a \wedge (z \wedge x))) \wedge (x \vee (a \wedge (z \wedge y)))$. [para(4(a,1),46(a,2,2))].
4337 $(a \wedge x) \vee (y \wedge a) = a \wedge (x \vee (y \wedge a))$. [para(46(a,1),9(a,2)),rewrite([9(5),100(7),126(9),4(13),6(13),2(10)])].
4352 $a \wedge (c1 \vee (c3 \vee (x \wedge a))) = a \wedge (c1 \vee (c2 \vee (x \wedge a)))$. [para(17(a,1),46(a,1,1)),rewrite([100(13),4337(8),5(7),126(15),5(14),110(20),4(22),99(22),2(16)]),flip(a)].
4586 $x \vee ((x \vee (a \wedge (y \wedge z))) \wedge (z \vee (a \wedge (y \wedge x)))) = x \vee (a \wedge (y \wedge z))$. [para(46(a,2),719(a,1,2,2)),rewrite([4331(9),909(13),25(12),233(12)])].
5088 $a \wedge (c2 \vee (c3 \vee (a \wedge x))) = a \wedge (c1 \vee (c2 \vee (x \wedge a)))$. [back_rewrite(1243),rewrite([4352(16)])].

5089 $(a \wedge (c1 \vee c2)) \vee (a \wedge x) = a \wedge (c1 \vee (c2 \vee (x \wedge a)))$. [*back_rewrite*
(964), *rewrite*([4352(16)])].
5201 $(x \vee y) \wedge ((z \wedge y) \vee x) = (z \wedge y) \vee x$. [*para*(176(a,1), 722(a,1,1))].
6279 $x \vee (a \wedge (y \wedge (x \vee z))) = x \vee (a \wedge (y \wedge z))$. [*para*(504(a,1), 60(a
,1,1)), *rewrite*([5(14), 427(13), 556(8), 277(10), 504(20), 25(15), 4(14)
, 890(14), 25(12), 4331(11), 4586(15)])].
6495 $((x \vee (y \wedge a)) \wedge z) \vee (z \wedge (y \vee (a \wedge x))) = z \wedge (x \vee y)$. [*para*(2(a
,1), 67(a,1,1))].
6508 $(x \wedge (y \vee (z \wedge a))) \vee (u \vee (x \wedge (z \vee (a \wedge y)))) = u \vee (x \wedge (y \vee z))$.
[*para*(67(a,1), 25(a,1,2)), *flip*(a)].
7094 $(a \wedge x) \vee (y \wedge x) = x \wedge ((a \wedge x) \vee y)$. [*para*(128(a,1), 4(a,1)), *flip*(a
)].
7695 $c1 \vee (c3 \vee (a \wedge c2)) = c1 \vee c3$. [*para*(166(a,1), 239(a,1,2,2))].
7732 $(c1 \vee c3) \wedge (c1 \vee (a \wedge c2)) = c1 \vee (a \wedge c2)$. [*para*(7695(a,1), 177(a
,1,2)), *rewrite*([2(9)])].
9660 $x \wedge ((x \wedge y) \vee (x \wedge z)) = (x \wedge y) \vee (x \wedge z)$. [*para*(6(a,1), 279(a,1,1)
)].
10028 $x \vee ((x \vee y) \wedge (x \vee z)) = (x \vee y) \wedge (x \vee z)$. [*para*(7(a,1), 294(a
,1,1))].
10127 $(x \wedge (y \wedge (z \vee u))) \vee ((w \wedge (x \wedge (y \wedge z))) \vee v5) = (x \wedge (y \wedge (z \vee u)
)) \vee v5$. [*para*(298(a,1), 233(a,1,2,1,2))].
10150 $(x \wedge (y \wedge (z \wedge u))) \vee (w \vee (y \wedge (z \wedge (u \vee v5)))) = w \vee (y \wedge (z \wedge (u
\vee v5)))$. [*para*(298(a,1), 556(a,1,1,2))].
10193 $a \wedge (c1 \vee (c2 \vee (a \wedge (c2 \vee (c3 \vee x)))))) = a \wedge (c2 \vee (c3 \vee x))$. [
para(19(a,1), 310(a,1,1)), *rewrite*([5(10), 5089(12), 2(9), 5(17)])].
10202 $(x \wedge (y \wedge z)) \vee (x \wedge (u \vee y)) = x \wedge (u \vee y)$. [*para*(281(a,1), 310(a
,1,2,2)), *rewrite*([281(8)])].
10210 $(x \wedge (y \wedge z)) \vee (z \wedge ((x \wedge (y \wedge z)) \vee u)) = z \wedge ((x \wedge (y \wedge z)) \vee u)$.
[*para*(224(a,1), 310(a,1,1))].
10338 $x \wedge ((y \wedge z) \vee (y \wedge x)) = x \wedge y$. [*para*(449(a,1), 321(a,1,2)), *rewrite*
([2(5), 3(5), 9660(4), 449(8)])].
10367 $x \vee (a \wedge (y \wedge (z \vee x))) = x \vee (a \wedge (y \wedge z))$. [*para*(321(a,1), 60(a
,1,1)), *rewrite*([5(15), 315(14), 10150(10), 321(20), 10127(16), 4(11), 25(12)
, 4331(11), 4586(15)])].
10914 $x \wedge ((y \wedge z) \vee (z \wedge x)) = x \wedge z$. [*para*(2(a,1), 10338(a,1,2,1))].
10915 $x \wedge ((y \wedge z) \vee (x \wedge y)) = x \wedge y$. [*para*(2(a,1), 10338(a,1,2,2))].
10940 $x \wedge ((y \wedge (z \wedge u)) \vee (u \wedge x)) = x \wedge u$. [*para*(224(a,1), 10338(a
,1,2,1))].
11021 $x \wedge ((y \wedge z) \vee (x \wedge z)) = x \wedge z$. [*para*(2(a,1), 10914(a,1,2,2))].
11183 $x \wedge ((y \wedge z) \vee (x \wedge (u \vee z))) = x \wedge (u \vee z)$. [*para*(340(a,1), 10915(a
,1,2,1))].
11568 $x \wedge (y \vee (x \wedge (z \vee (u \vee y)))) = x \wedge (z \vee (u \vee y))$. [*para*(160(a,1)
, 11021(a,1,2,1))].
12001 $x \vee (y \vee (z \vee (u \vee (w \wedge x)))) = y \vee (z \vee (u \vee x))$. [*para*(202(a,1)
, 322(a,1,2,2)), *rewrite*([391(9)])].
12204 $x \vee ((y \vee x) \wedge (x \vee z)) = (y \vee x) \wedge (x \vee z)$. [*para*(7(a,1), 338(a
,1,1))].
12396 $x \vee (y \vee (((x \wedge z) \vee (y \wedge u)) \wedge w)) = x \vee y$. [*para*(357(a,1), 280(a
,1,2)), *rewrite*([169(3)]), *flip*(a)].
13792 $x \vee ((y \vee x) \wedge (y \vee z)) = x \vee y$. [*para*(504(a,1), 372(a,1,2)), *rewrite*
([25(5), 4(4), 10028(4), 504(8)])].
13882 $x \vee ((y \vee z) \wedge (y \vee x)) = x \vee y$. [*para*(2(a,1), 13792(a,1,2))].

13902 $x \vee ((y \wedge z) \vee x) \wedge (u \vee z) = x \vee (y \wedge z)$. [para(323(a,1),13792(a,1,2,2))].
14400 $x \vee ((y \vee z) \wedge (z \vee x)) = x \vee z$. [para(4(a,1),13882(a,1,2,1))].
14415 $x \vee ((y \vee z) \wedge ((u \wedge z) \vee x)) = x \vee (u \wedge z)$. [para(323(a,1),13882(a,1,2,1))].
14736 $x \vee ((y \vee z) \wedge (z \vee (u \wedge x))) = x \vee z$. [para(14400(a,1),149(a,1,2)), rewrite([149(3)]), flip(a)].
17603 $x \vee (a \wedge (x \vee y)) = x \vee (a \wedge y)$. [para(127(a,1),28(a,1,2)), rewrite([14415(8)]), flip(a)].
17613 $x \vee (a \wedge (y \vee x)) = x \vee (a \wedge y)$. [para(127(a,1),149(a,1,2)), rewrite([13902(8)]), flip(a)].
17730 $(a \wedge x) \vee (a \wedge y) = a \wedge (x \vee y)$. [para(127(a,2),43(a,1,2,2)), rewrite([11183(10),10202(7),2(10),3(10),32(9),124(7),2(11),31(12),5201(12)]), flip(a)].
18007 $a \wedge (c1 \vee (c2 \vee (a \wedge (c3 \vee x)))) = a \wedge (c2 \vee (c3 \vee x))$. [back_rewrite(10193), rewrite([17603(10)])].
18151 $a \wedge (c1 \vee (c2 \vee (x \wedge a))) = a \wedge (c1 \vee (c2 \vee x))$. [back_rewrite(5089), rewrite([17730(8),5(5)]), flip(a)].
18251 $a \wedge (c2 \vee (c3 \vee (a \wedge x))) = a \wedge (c1 \vee (c2 \vee x))$. [back_rewrite(5088), rewrite([18151(16)])].
18578 $c1 \vee (a \wedge c3) = c1 \vee (a \wedge c2)$. [para(17(a,1),17603(a,1,2)), rewrite([17603(7)]), flip(a)].
18580 $a \wedge (x \vee (a \wedge y)) = a \wedge (x \vee y)$. [para(17603(a,1),32(a,1,2)), rewrite([3(7),400(6)])].
18588 $a \wedge (x \wedge (y \vee (a \wedge z))) = x \wedge (a \wedge (y \vee z))$. [para(17603(a,1),166(a,1,2,2)), rewrite([3(8),606(7)])].
18658 $a \wedge (c1 \vee (c2 \vee (c3 \vee x))) = a \wedge (c1 \vee (c3 \vee x))$. [para(91(a,1),17603(a,2)), rewrite([25(24),4(23),17730(23),25(20),25(19),5(18),25(19),25(18),118(20),18580(21),12001(18),4(17),17730(17),25(14),25(13),4(12),25(12),25(11),25(10),4(9),26(9),25(10),25(9),118(8),118(9)])].
18661 $a \wedge (c2 \vee (c3 \vee x)) = a \wedge (c1 \vee (c2 \vee x))$. [para(98(a,1),17603(a,2)), rewrite([18251(8),18251(15),25(20),4(19),898(19),11568(15),34(13)]), flip(a)].
19033 $a \wedge (c1 \vee (c2 \vee (a \wedge x))) = a \wedge (c1 \vee (c2 \vee x))$. [back_rewrite(18251), rewrite([18661(8)])].
19034 $a \wedge (c1 \vee (c3 \vee x)) = a \wedge (c1 \vee (c2 \vee x))$. [back_rewrite(18007), rewrite([19033(10),18658(8),18661(12)])].
19518 $c3 \vee (a \wedge (c1 \vee c2)) = c3 \vee (a \wedge c1)$. [para(17(a,1),17613(a,1,2))].
19519 $c3 \vee (a \wedge c2) = c3 \vee (a \wedge c1)$. [para(19(a,1),17613(a,1,2)), rewrite([19518(7)]), flip(a)].
19765 $a \wedge ((a \wedge x) \vee y) = a \wedge (x \vee y)$. [para(124(a,1),17730(a,1,1)), rewrite([17730(5)]), flip(a)].
22049 $(a \wedge x) \vee (x \wedge y) = x \wedge ((a \wedge x) \vee y)$. [para(6(a,1),191(a,1,2,1)), rewrite([6(9),2(8)])].
30100 $(x \wedge y) \vee ((x \vee z) \wedge (u \vee y)) = (x \vee z) \wedge (u \vee y)$. [para(166(a,1),296(a,1,2)), rewrite([4(5)])].
40090 $c2 \vee ((x \vee c3) \wedge (c3 \vee (a \wedge c1))) = c2 \vee c3$. [para(19519(a,1),14736(a,1,2,2))].
44335 $c1 \vee (a \wedge (c3 \vee x)) = c1 \vee (a \wedge (c2 \vee x))$. [para(19034(a,1),17603(a,1,2)), rewrite([17603(8)]), flip(a)].
46202 $(a \wedge (x \wedge y)) \vee (y \wedge z) = y \wedge ((a \wedge (x \wedge y)) \vee z)$. [para(10940(a,1),427(a,2)), rewrite([10210(12),2(9),166(9),2(10)])].

48249 $c1 \vee (a \wedge (c2 \wedge (x \vee c3))) = c1 \vee (a \wedge c2)$. [para(40090(a,1),44335(a,2,2,2)),rewrite([12204(12),18588(11),4(7),17(8),23(9),6279(10),2(6),19(14),17603(15)])].

49221 $c1 \vee (a \wedge (c2 \wedge c3)) = c1 \vee (a \wedge c2)$. [para(34(a,1),48249(a,1,2,2,2))].

49233 $c1 \vee (c2 \wedge (c3 \vee (a \wedge c1))) = c1 \vee (c2 \wedge c3)$. [para(49221(a,1),176(a,1,2)),rewrite([25(9),4(8),22049(8),4(7),19519(7)])].

52023 $a \wedge (x \vee (y \wedge z)) = a \wedge ((x \vee z) \wedge (x \vee y))$. [para(537(a,1),19765(a,1,2)),rewrite([4(9),2810(9),17730(8),23(7),23(8),124(7),124(7),10367(11),18580(11)]),flip(a)].

63667 $a \wedge (x \wedge (y \vee (z \wedge u))) = x \wedge (a \wedge ((y \vee u) \wedge (y \vee z)))$. [para(12396(a,1),479(a,1,2,2,2)),rewrite([298(8),2(4),52023(4)]),flip(a)].

63754 $x \vee (y \vee (z \wedge (x \vee (u \wedge z)))) = y \vee (x \vee (u \wedge z))$. [para(921(a,1),390(a,1,2))].

64469 $x \vee (a \wedge ((y \vee x) \wedge z)) = x \vee (a \wedge (y \wedge z))$. [para(745(a,1),941(a,1,1,2)),rewrite([52023(5),120(3),2(10),63667(11),120(8),293(11),4(10),6279(10),25(10),4(9),17730(9),30100(6),6279(6),2(10),63667(11),120(8),293(11),4(10),6279(10)])].

69039 $x \vee ((x \vee y) \wedge (z \vee (a \wedge x))) = x \vee ((x \vee y) \wedge z)$. [para(1108(a,1),7094(a,1,2)),rewrite([52023(5),118(4),25(9),46202(8),4(7),64469(7),7(4),52023(14),118(13),4(16),148(16),2(11),1108(11)])].

104056 $c1 \vee (c2 \wedge (c1 \vee c3)) = (c1 \vee c2) \wedge (c1 \vee c3)$. [para(7732(a,1),6495(a,1,2)),rewrite([2(4),2(9),4(15),5(15),267(14),25(13),4(12),19519(12),176(13),4(9),2(10),69039(11),2(6),4(13),2(14)])].

104095 $c1 \vee (c2 \wedge c3) = (c1 \vee c2) \wedge (c1 \vee c3)$. [para(49233(a,1),6508(a,1,2)),rewrite([2(5),18578(6),25(13),4(12),63754(13),25(9),4(8),22049(8),4(7),19519(7),49233(9),104056(12)])].

104096 \$F. [resolve(104095,a,21,a)].

===== **end of proof**=====

===== **STATISTICS**=====

Given=4826. Generated=39046099. Kept=104089. proofs=1.
Usable=4530. Sos=19712. Demods=22873. Limbo=11, Disabled=79850. Hints=0.
Kept_by_rule=0, Deleted_by_rule=26712.
Forward_subsumed=19936868. Back_subsumed=810.
Sos_limit_deleted=18978430. Sos_displaced=63145. Sos_removed=0.
New_demodulators=92229 (6 lex), Back_demodulated=15878. Back_unit_deleted=0.
Demod_attempts=1136610097. Demod_rewrites=133339635.
Res_instance_prunes=0. Para_instance_prunes=0. Basic_paramod_prunes=0.
Nonunit_fsub_feature_tests=0. Nonunit_bsub_feature_tests=0.
Megabytes=136.64.
User_CPU=963.34, System_CPU=15.07, Wall_clock=979.

===== **end of statistics**=====

===== **end of search**=====

THEOREM PROVED