



## BIROn - Birkbeck Institutional Research Online

Mikulás, Szabolcs and Sain, I. and Simon, A. (2015) Complexity of equational theory of relational algebras with standard projection elements. *Synthese* 192 (7), pp. 2159-2182. ISSN 0039-7857.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/13400/>

*Usage Guidelines:*

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>  
contact [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk).

or alternatively

# Complexity of Equational Theory of Relational Algebras with Standard Projection Elements\*

Szabolcs Mikulás

szabolcs@dcs.bbk.ac.uk

School of Comp. Sci. and Information Systems,  
Birkbeck College, University of London

Ildikó Sain

sain.ildiko@renyi.mta.hu

Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences

András Simon

asimon@math.bme.hu

Department of Algebra, Budapest University of Technology and Economics

February 2015

## Abstract

The class  $\text{TPA}$  of *true pairing algebras* is defined to be the class of relation algebras expanded with *concrete* set theoretical projection functions. The main results of the present paper is that neither the equational theory of  $\text{TPA}$  nor the first order theory of  $\text{TPA}$  are decidable. Moreover, we show that the set of all equations valid in  $\text{TPA}$  is exactly on the  $\Pi_1^1$  level. We consider the class  $\text{TPA}^-$  of the relation algebra reducts of  $\text{TPA}$ 's, as well. We prove that the equational theory of  $\text{TPA}^-$  is much simpler, namely, it is recursively enumerable. We also give motivation for our results and some connections to related work.

---

\*Research supported by Hungarian National Foundation for Scientific Research grant Numbers T030314, T034861, and T035192.

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Finitization Problem and True Pairing Algebras</b>	<b>3</b>
2.1	Algebraizing Logic . . . . .	3
2.2	Hilbert style axiomatizability versus finitely based variety . . . . .	4
2.3	Finitization problem . . . . .	5
2.4	Finitization of first order logic <i>with</i> equality . . . . .	6
<b>3</b>	<b>Complexity of <math>\text{Eq}(\text{TPA})</math> and some consequences</b>	<b>7</b>
<b>4</b>	<b>The complexity of <math>\text{Eq}(\text{TPA})</math> is at least <math>\Pi_1^1</math></b>	<b>9</b>
<b>5</b>	<b>The complexity of <math>\text{Eq}(\text{TPA})</math> is at most <math>\Pi_1^1</math></b>	<b>15</b>
<b>6</b>	<b><math>\text{Eq}(\text{TPA}^-)</math> is recursively enumerable</b>	<b>22</b>
<b>7</b>	<b>Related work</b>	<b>23</b>

## 1 Introduction

The topic of this paper is relevant to the following parts of science: *logic, algebraic logic, recursion theory, and theoretical computer science*. The relevance for the latter is explained e.g. in [12], [41], [43].

The results of this paper concern true pairing algebras (TPA's for short). The class TPA was introduced, e.g., in Maddux [21, Definitions 17–20], and in different form, independently, in Veloso–Haeblerer [42], [43]. TPA's in a different form are also discussed in [40, item 4.1 (iv) on p.96, pp. 251–254]. The purpose for which TPA's are used in these publications is, roughly speaking, to provide a nicer formulation of first order logic with equality. By “nice” here we mean a certain point of view. This point of view was described in various works on logic and algebraic logic, and is often referred to as the finitization problem. Section 2 serves for outlining the latter — to the extent we believed it is useful for motivating our results presented in the rest of the sections.

A positive solution for the finitization problem for first order logic (FOL) *without equality* was given in [31], and published later in [32], [33], [35] (among which, the last is the most detailed account on the subject). R. J. Thompson conjectured that there is no positive solution for the finitization problem for FOL with equality. His conjecture is motivated by the fact that he proved that the approach in [35] cannot be extended to FOL with equality, cf. [31, §5], [35, §6]. As it will be explained in subsection 2.4, TPA's were suggested as an alternative approach to solve the finitization problem for FOL with equality.

We will prove that the equational theory  $\text{Eq}(\text{TPA})$  of TPA's is (not only not recursively enumerable but) such that its degree of unsolvability is exactly  $\Pi_1^1$ . On the other hand, let  $\text{TPA}^-$  be obtained from  $\text{TPA}$  by forgetting the constant symbols naming the projection functions. I.e.,  $\text{TPA}^-$  is that reduct of  $\text{TPA}$  which belongs to the similarity class of relation algebras (RA's). Then  $\text{Eq}(\text{TPA}^-)$  is recursively enumerable but is still not finitely axiomatizable.

In [42, sec.3] the standard models  $\mathbf{U}$  of the Extended Calculus of Binary Relations (ECBR for short) are defined. These are relation algebras with some extra operators just as TPA's are such. We will not need to recall the precise definition of these algebras. Let  $\text{SECBR}$  denote the class of these standard models of ECBR'. Our Corollary 3.3 (after Theorem 3.2) shows that the programme initiated in [42] has certain limitations. It also seems to point in the direction that this approach might not be very convenient for attacking the finitization problem.

Our paper consists of sections 2–7. In section 2 we give a survey of our motivation for our results, exactly stated and proved in the later sections. These results were obtained in 1990. Until now they existed in the form of the research report [22] and were abstracted in [23]. In section 7 we mention some more results concerning the topic of our paper.

The authors are grateful to William Craig, Steve Givant, Roger Maddux, István Németi, Gábor Sági, and Tarek Sayed-Ahmed supporting us, in various ways, in completing this paper.

## 2 Finitization Problem and True Pairing Algebras

### 2.1 Algebraizing Logic

In algebraic logic, to several kinds of logical systems (*logics* for short) — like classical propositional logic, modal logics, first order logic with or without equality — we associate classes of algebras via an algorithm which is described, e.g., in [15, §4.3, pp. 258–260] or in [4, Part II], [3, section 3], [5], [34, sec.3]. We call such a class of algebras associated to a logic the algebraic counterpart of it. For example, the algebraic counterpart of classical propositional logic is the class  $\mathbf{BA}$  of Boolean algebras. The algebraic counterpart of FOL with equality (using  $\omega$  many variables) is the class  $\mathbf{RCA}_\omega$  of representable cylindric algebras (of dimension  $\omega$ ), cf. e.g. [15, §4.3] or [4, p. 237].

The point in finding the algebraic counterparts of logics is that logical properties of a logic can often be correlated with algebraic properties of the algebraic counterpart of the logic. Thus not only logical systems can have their algebraic counterparts, but logical properties can have their algebraic counterparts as well.

Going even further, an abstract notion of a logic has been worked out, cf. e.g.

in [7], [4], [5], [3], [34]. Such an abstract logic is a common generalization of the systems mentioned above (and more). We can define (abstract) logical properties in this setting, for example, *interpolation properties*, *definability properties*, *existence of derivation systems*, *decidability* can be formulated in this general setting. Then the algebraic counterparts of these properties can also be formulated. Then one can find algebraic properties which correspond to these logical properties, in the following sense.

An abstract logic  $\mathcal{L}$  has logical property  $\Phi$   
 $\iff$   
 its algebraic counterpart  $\text{Alg}(\mathcal{L})$  has algebraic property  $\text{Alg}(\Phi)$ .

Classical theorems state the equivalence of *Beth definability* of a logic  $\mathcal{L}$  and *surjectiveness of epis* in the category formed by  $\text{Alg}(\mathcal{L})$  and the homomorphisms between them (cf. e.g. [4, Thm.58]). Another example for such results is that the Craig interpolation property of  $\mathcal{L}$  is equivalent with the amalgamation property of  $\text{Alg}(\mathcal{L})$  (a precise formulation is, e.g., [4, Thm.62]).

## 2.2 Hilbert style axiomatizability versus finitely based variety

Theorems in the spirit of the ones mentioned in the previous subsection connect properties of derivation systems of a logic with axiomatizability properties of the corresponding classes of algebras. Such a theorem, interesting from the point of view of the present paper, states that

A logic  $\mathcal{L}$  has a truly finite Hilbert style inference system  
 $\iff$   
 its algebraic counterpart  $\text{Alg}(\mathcal{L})$  is a finitely based quasi-variety,

where a “*finitely based (quasi-)variety*”, by definition, is a class of algebras which can be axiomatized by a finite set of (quasi-)equations.

The classical example for a logic, where the statement on the left hand side of the above double implication is satisfied (and thus also the one on the right hand side of it), is classical propositional logic. Here we have a Hilbert style inference system with a truly finite schema of logical axioms on one hand; and, on the other hand, the corresponding class, **BA** is a finitely based variety (by Stone’s theorem).

If we consider FOL, the picture is different. The axioms and rules of the usual inference system of FOL refer to individual variables, and thus they do not form a finite schema. Looking at the other side of the implication, we find that, though the natural algebraic counterpart  $\text{RCA}_\omega$  of FOL is a variety (i.e., axiomatizable by a set of equations), it is very far from being axiomatizable by a finite set of equations. Actually, any set  $\Sigma$  of equations axiomatizing  $\text{RCA}_\omega$  must contain infinitely many

variables, moreover, to any  $n \in \omega$ , there is an  $e \in \Sigma$  containing more than  $n$  fundamental operations as well as more than  $n$  variables, a result of Andr eka (improving a result of Monk), cf. [27], [1].

## 2.3 Finitization problem

1) The finitization problem asks whether the negative result quoted in the last paragraph of subsection 2.2 is an inherent property of FOL or not. Perhaps the negative result is due to an unfortunate but not necessary choice of the basic logical connectives (the Boolean connectives plus quantifiers and equalities). Perhaps one could choose the basic logical connectives in such a way that the above negative property of the derivation system of FOL, or equivalently, the so complex non-finite axiomatizability of the algebraic counterpart of FOL would disappear.

This problem (and many variants) was formulated several times in the literature of logic and algebraic logic. We mention here only some of them: [24], [14, Problem 1], [27]. In [24], J. D. Monk formulates the finitization problem as follows:

*“Devise an algebraic version of predicate logic in which the class of representable algebras forms a finitely based equational class”.*

2) A strongly related problem is raised and discussed in [40, pp.56–62]. The research reported in [40] is aiming at finitizing set theory — finitizing this way the whole of mathematics, in that the traditional foundation of mathematics is set theory. This problem, viewed from the perspective of FOL, has to do with finitizing (algebraic counterparts of) *finite variable fragments* of FOL, cf. e.g. [28], [37], [2, p.16], while the former problem deals with usual FOL having a sequence of variables of order type  $\omega$ . In other words, in [40], instead of the expressive power of full FOL, they only require that of the three-variable fragment of FOL to be still available in the new formulation.

Alfred Tarski formulated set theory in the equational theory of *relation algebras*. The classes of algebras (QRA, TPA, and their variants) investigated in the present paper were introduced first in [40].

There are some further natural requirements, namely that all logical connectives as well as their algebraic counterparts should be permutation invariant, cf. [27], [35], [40, §3.5, p. 57 item 3.5(i)], J nsson [17], [8, p. 55 lines 16–20]. We note that this requirement is needed, for FOL, to fulfill a most natural model theoretic requirement, namely, that isomorphic models satisfy the same formulas. A discussion related to the necessity of permutation invariance is [39].

*The present paper addresses directly the second problem, stressing that we require permutation invariance briefly described in the previous paragraph.*

We note that [37] gives a good exposition of the finitization problem. Some recent results concerning the finitization problem are mentioned in our subsection 7.

## 2.4 Finitization of first order logic *with* equality

As we mentioned in the Introduction, the finitization problem for FOL without equality does have a solution. Now we discuss an approach attacking the finitization problem for FOL with equality which leads to the use of TPA's. When doing this, below we outline further facts supporting Thompson's conjecture (cf. in the Introduction) that the finitization problem might *not* admit a positive solution for FOL with equality. This is in sharp contrast with the case of logic without equality. In this connection, we feel, algebraic logic might give some useful information back to logic.

This approach seems to be similar to that of Craig–Vaught [9], and was suggested in the lecture [20] and independently (in different form) in Veloso–Haeberer [42], [43]. It is based on Tarski's *quasi-projective relation algebras* (QRA's) (see [40], to be recalled in section 3 below). The idea is that QRA is a reduct of a finitely based variety and all QRA's are representable. As it was pointed out by Leon Henkin, George McNulty, and other participants during the discussion of the lecture [20], three problems have to be solved in trying to implement this idea. These are the following.

1. QRA is *not* a variety, and the finitization problem, cf. e.g. Henkin–Monk [14], explicitly writes “. . . class of representable algebras forms a finitely based *equational class*”. So the problem asks for a variety and not for a reduct of one.
2. The projection functions are not logical in the sense of [40], i.e., they are not permutation invariant in Jónsson's sense.
3. The set QRA's have all infinite bases, which seems to mean that the corresponding logic has no finite models (at least if we use the standard translation between algebraic logic and logic as described in our subsection 2.1).

To alleviate item 1, we could add two constant symbols say  $p$  and  $q$  to the language of QRA, and add some axioms on these constants (like  $p^{-1} \circ q = 1$ ). The intended meaning is that  $p$  and  $q$  are the so called *quasi-projections*. The new finitely based variety  $\text{QRA}^+$  is such that its RA-reduct is exactly QRA. So  $\text{QRA}^+$  is a candidate for being the variety the Henkin–Monk problem asks for. The problem requires that every element of  $\text{QRA}^+$  should be *representable*. Representability of an algebra  $\mathfrak{A}$  means that  $\mathfrak{A}$  is isomorphic to some  $\mathfrak{A}^+$  *all* operations of which are *concrete* set theoretic ones (like intersection, complementation, relation composition etc.). Therefore the constant operations  $p$  and  $q$  of  $\mathfrak{A}^+$  should be concrete set theoretic constants (like the empty set, the identity relation etc.). Briefly, we say that  $p$  and  $q$  of  $\text{QRA}^+$  should be representable, too. As it was anticipated by Henkin ([13]), we will see that this is not easy to arrange. Of course, we may add finitely many new equations to those defining  $\text{QRA}^+$  to ensure representability of  $p$  and  $q$  (of some subvariety  $\mathbf{V}$  of  $\text{QRA}^+$  with  $\mathbf{V}$  sufficiently large for the original purposes). Theorem 3.4 below seems to say that this is still not easy to do. In connection with

representability of members of  $\mathbf{QRA}^+$  we note that, e.g. in Definition 3.1 below, the subclass  $\mathbf{TPA}$  of  $\mathbf{QRA}^+$  consists of representable algebras because  $p$  and  $q$  are represented as concrete set theoretic constants in the definition of  $\mathbf{TPA}$ . For the whole of  $\mathbf{QRA}^+$ , the present authors have not yet seen a proposed notion of representation that would intend to be similar in concreteness to the representation proposed by  $\mathbf{TPA}$ . Summing up,  $\mathbf{QRA}^+$  is a finitely based variety, but we do not see a representation for it. Therefore, we turn to its subclass  $\mathbf{SP}(\mathbf{TPA})$  consisting of representable algebras to see if we can obtain a finitely based variety from that. Note that our problem in item 1 above does *not* involve the question of whether the operations are logical. So if we investigate the Henkin–Monk problem without the new requirement added in [40], our problem 1 above (i.e., Theorem 3.4 below) still has to be dealt with somehow.

Let us return to a more concrete investigation of problems 1–3 above. [20] proposed to approach this problem by replacing the usual set  $\mathbf{QRA}$ 's with  $\mathbf{TPA}$ 's. Independently, the same approach is taken in [42], [43]. Roughly, a  $\mathbf{TPA}$  is a set  $\mathbf{QRA}$  with a base  $U$  and a distinguished subset  $U_0$  of  $U$  such that the elements of  $U_0$  are not pairs while  $U$  is the closure of  $U_0$  under forming ordered pairs. That is

$$(1) \quad U = \bigcup_{n \in \omega} U_n, \quad \text{where } U_{n+1} = (U_n \times U_n) \cup U_n.$$

Further, the projection functions of this  $\mathbf{QRA}$  are the standard set theoretic ones. Then we say that  $U_0$  corresponds to the universe of that model (of logic) from which our  $\mathbf{TPA}$  is obtained and the rest of  $U$ , i.e.,  $U \setminus U_0$  corresponds to (part of the) logic itself. Hence the elements of  $U \setminus U_0$  are “logical” elements. Therefore, when this approach reaches the point where the notion of “set algebra” or representable algebra is introduced, then the elements of  $U \setminus U_0$  should be required to be some “standard set theoretical” constructs (obtained from  $U_0$ ). Indeed, this is the spirit in which  $\mathbf{TPA}$ 's are defined in [20], [21], [42], [43]. Actually, in [42], [43] the word is not “ $\mathbf{TPA}$ ” but “ $\mathbf{SECBR}$ ” (cf. the relevant paragraph in the Introduction), but the results and methods of the present paper apply to those algebras the same way as to  $\mathbf{TPA}$ 's, see [43, §3]. From the point of view of the present paper it is interesting to note that, around the end of [43, §6 (comparison of  $\mathbf{TPA}$ 's and cylindric algebras)], they seem to conjecture that the equational theory of  $\mathbf{TPA}$  might be finitely axiomatizable. We will prove below that no expansion of the theory can have the conjectured property.

### 3 Complexity of $\mathbf{Eq}(\mathbf{TPA})$ and some consequences

Let us recall from e.g. [15, Def.5.3.1 p.211] that a *relation algebra* ( $\mathbf{RA}$  for short) is a system  $\mathfrak{A} = \langle A, +, \cdot, -, 0, 1, \circ, {}^{-1}, 1' \rangle$  such that  $\langle A, +, \cdot, -, 0, 1 \rangle$  is a  $\mathbf{BA}$ ,  $\circ$  is a binary operation on  $A$  (called composition),  ${}^{-1}$  is a unary operation on  $A$  (called converse),  $1' \in A$  (called identity), and  $\mathfrak{A}$  satisfies certain conditions. Recall, e.g. from Tarski–Givant [40], that the class of quasi-projective relation algebras is

$$\mathbf{QRA} \stackrel{\text{def}}{=} \{ \mathfrak{A} \in \mathbf{RA} : \mathfrak{A} \models \exists x \exists y ((x^{-1} \circ x + y^{-1} \circ y \leq 1') \wedge (x^{-1} \circ y = 1)) \}.$$



When speaking of a QRA  $\mathfrak{A}$ , its elements  $x, y$  in the above formula are called  $\mathfrak{A}$ 's *projection functions*, and are denoted by  $p$  and  $q$ . By a set QRA we understand one that is a relation set algebra in the sense of [15].  $\mathfrak{R}\mathfrak{e}(U)$  denotes the full relation set algebra with base  $U$  in the sense of [15, Def. 5.3.2]. Thus the greatest element of  $\mathfrak{R}\mathfrak{e}(U)$  is  $U \times U$  and all its elements are binary relations on  $U$ .

**3.1 DEFINITION** ([20], [21, §4 Def. 20])

1.  $\mathfrak{A}$  is called a *full true pairing algebra* (full TPA for short) iff  $\mathfrak{A} = \langle \mathfrak{R}, p, q \rangle$ , where  $\mathfrak{R} = \mathfrak{R}\mathfrak{e}(U)$  for some set  $U$  and

$$p = \{ \langle \langle x, y \rangle, x \rangle : x, y \in U \}, \quad q = \{ \langle \langle x, y \rangle, y \rangle : x, y \in U \}.$$

$p$  and  $q$  are called projection functions of  $\mathfrak{A}$ .

2.  $\text{TPA} = \mathbf{SP}(\text{full TPA})$ , i.e.,  $\mathfrak{A} \in \text{TPA}$  iff  $\mathfrak{A}$  is isomorphic to a subalgebra of a direct product of full TPA's. TPA is called the class of all true pairing algebras.

△

In formulating Theorem 3.2 below, we use the terminology of recursion theory for clarifying the “uncomputability” of non-computable functions. The complexity classes are denoted by  $\Pi_n^k$  and  $\Sigma_n^k$  ( $n, k \in \omega$ ), where if  $k > 0$ , then the set in question is so far from being computable that it is called non-arithmetical. (This implies that such sets are not definable in set theory without the Axiom of Infinity.)

**3.2 THEOREM** *The set  $\text{Eq}(\text{TPA})$  of all equations valid in TPA is  $\Pi_1^1$  complete (in the analytical hierarchy). That is, the “degree of unsolvability” or measure of definability of  $\text{Eq}(\text{TPA})$  is exactly  $\Pi_1^1$ .*

*Proof.* This is an immediate consequence of Theorems 4.1 and 5.1 below. □

**3.3 COROLLARY** *The equational theory of SECBR is  $\Pi_1^1$ -complete.*

*Proof.* The TPA-operations are term-definable in SECBR, hence our theorem immediately yields this corollary. □

**3.4 THEOREM** 1. *The variety generated by TPA is not finitely based (i.e.,  $\mathbf{HSP}(\text{TPA})$  is not finitely axiomatizable). Moreover,  $\mathbf{HSP}(\text{TPA})$  is not axiomatizable by any decidable (or even recursively enumerable) set of axioms.*

2. *The first order theory  $\text{Th}(\text{TPA})$  of TPA is not finitely axiomatizable.  $\text{Th}(\text{TPA})$  is not axiomatizable by any decidable set either.*

*Proof.* This is an immediate consequence of Theorem 3.2, but, for completeness, we give a more direct proof too.

(1) Assume that  $\mathbf{HSP}(\mathbf{TPA}) = \text{Mod}(E)$  where  $E$  is a recursively enumerable (r.e.) set of equations. Then  $\{e \in \text{Equations} : \mathbf{TPA} \models e\} = \{e \in \text{Equations} : E \vdash e\}$  is r.e., contradicting an immediate consequence of Theorem 4.1 below: since  $\text{Eq}(\mathbf{TPA})$ , i.e., all equations valid in  $\mathbf{TPA}$ , is at least on the  $\Pi_1^1$  level by Theorem 4.1, we get that  $\text{Eq}(\mathbf{TPA})$  is not r.e. Since every finite set is r.e., this completes the proof of (1).

(2) Assume that  $\text{Th}(\mathbf{TPA}) = \{\phi : Ax \vdash \phi\}$  with  $Ax$  r.e. Then  $\{e \in \text{Equations} : e \in \text{Th}(\mathbf{TPA})\}$  is r.e., too, contradicting the above fact again.  $\square$

## 4 The complexity of $\text{Eq}(\mathbf{TPA})$ is at least $\Pi_1^1$

Let  $\Omega = \langle \omega, \text{pred}, +, \cdot \rangle$ , where  $\text{pred}, +, \cdot$  are, respectively, the usual predecessor, addition, and multiplication on the set  $\omega$  of natural numbers. Thus

$$\text{pred} = \{\langle n, n - 1 \rangle : 0 < n \in \omega\}.$$

**4.1 THEOREM** *Let  $\psi$  be a  $\Pi_1^1$  sentence of arithmetic. Then there is an equation  $\epsilon_\psi$  in the language of  $\mathbf{TPA}$ , effectively calculable from  $\psi$ , such that*

$$\Omega \models \psi \iff \mathbf{TPA} \models \epsilon_\psi.$$

*Proof.* By a standard pairing structure we understand a structure  $\langle U, p, q \rangle$  where  $U$  is as in (1) at the end of the previous section, and  $p, q$  are the two projection functions, see in item 1 of Def.3.1 above.

First we write up a quasi-equation  $(e_1 \wedge e_2 \wedge e_3)(N, A, M)$  in the language of  $\mathbf{TPA}$  containing three variables  $N, A, M$ . The intended meaning of  $(e_1 \wedge e_2 \wedge e_3)(N, A, M)$  is that  $N$  is the set of natural numbers and  $A : N \times N \longrightarrow N$ ,  $M : N \times N \longrightarrow N$  are the functions addition and multiplication, respectively. (All this is meant up to isomorphism.) The first order version  $\Phi_N \wedge \Phi_A \wedge \Phi_M$  of  $e_1 \wedge e_2 \wedge e_3$  contains

$$\begin{aligned} \Phi_N \stackrel{\text{def}}{=} & (\forall x \in N)(\exists y \in N)(p(y) = x = q(y)) \wedge |N \setminus \text{Dom}(p)| = 1 \\ & \wedge (\forall x \in N \cap \text{Dom}(p))(p(x) = q(x) \in N). \end{aligned}$$

If  $\langle U, p, q \rangle$  is a standard pairing structure and  $\langle U, p, q, N \rangle \models \Phi_N$ , then  $\langle N, p \upharpoonright N \rangle \cong \langle \omega, \text{pred} \rangle$ . This is true because by the Axiom of Foundation (in the set theory we are working in)  $(\forall x \in N)(\exists n \in \omega)(p^n(x) \notin \text{Dom}(p)$  but  $p^n(x)$  is defined). Then letting  $f(x) \stackrel{\text{def}}{=} n$ , we defined a bijection  $f : N \rightsquigarrow \omega$  which turns out to be the desired isomorphism.

In describing the equational forms (of  $\Phi_N$  etc.), we will need the following fact. It is well known in RA-theory that any Boolean combination of equations is equivalent

with a single equation in the subdirectly irreducible  $\mathbf{RA}$ 's. This follows, e.g., from the fact that  $\mathbf{RA}$ 's form a discriminator variety. For a careful and illuminating exposition of all this see [16]. (We note that in algebraic logic this property of  $\mathbf{RA}$ 's was known already in the 19th century.)

The equational form of  $\Phi_N$  is (with  $N$  as a variable)

$$e_1 \stackrel{\text{def}}{=} (1.1) \wedge (1.2) \wedge (1.3) \wedge (1.4) \wedge (1.5), \text{ where}$$

$$(1.1) \stackrel{\text{def}}{=} (N \leq 1'), \text{ meaning, of course, that } N \subseteq \{\langle x, x \rangle : x \in U\}$$

$$(1.2) \stackrel{\text{def}}{=} (1 \circ N \circ (p \cap q) \geq N), \text{ i.e., } \{\langle x, y \rangle : \exists z(\langle z, \langle y, y \rangle \rangle \in N)\} \supseteq N, \\ \text{which, together with (1.1), implies that } N \text{ is closed under successor}$$

$$(1.3) \stackrel{\text{def}}{=} (1 \circ N \circ 1 = 1), \text{ i.e., } N \neq \emptyset$$

$$(1.4) \stackrel{\text{def}}{=} [1 \circ (N - (p \circ 1))] \cap [(N - (p \circ 1)) \circ 1] \leq 1', \text{ i.e., } |(U_0 \times U_0) \cap N| \leq 1, \\ \text{uniqueness of } 0$$

$$(1.5) \stackrel{\text{def}}{=} (N \circ p = N \circ q \leq 1 \circ N), \text{ i.e.,} \\ (\forall x \in N \cap \text{Dom}(p) \cap \text{Dom}(q))(p(x) = q(x) \in N), \\ \text{i.e., } N \text{ is closed under predecessor.}$$

(\*1) If  $\mathfrak{B} = \langle \mathfrak{Ae}(U), p, q \rangle \in \text{TPA}$  and  $N \subseteq U$ , then  $\mathfrak{B} \models e_1[\text{Id} \upharpoonright N]$  implies that  $\langle U, p, q, N \rangle \models \Phi_N$  hence  $\langle N, p \rangle \cong \langle \omega, \text{pred} \rangle$ .

By (\*1) we manage to “code” the set of natural numbers by an equation  $e_1$  in  $\text{TPA}$ 's. This is similar to the beginning of the proof of Thm.2 in Némethi [26], see the explanation of the idea of the proof of Thm.3.1 on p.252 therein, or the following picture of  $N$  (described by  $e_1$ ).

$$\begin{array}{c} \vdots \\ \langle \langle \langle u_0, u_0 \rangle, \langle u_0, u_0 \rangle \rangle, \langle \langle u_0, u_0 \rangle, \langle u_0, u_0 \rangle \rangle \rangle \\ \downarrow p \\ \langle \langle u_0, u_0 \rangle, \langle u_0, u_0 \rangle \rangle \\ \downarrow p \\ \langle u_0, u_0 \rangle \end{array}$$

Here  $u_0 \in U_0$ . Note that  $\text{Dom}(N)(= \text{Rng}(N))$  satisfies  $\Phi_N$  in  $\langle U, p, q, N \rangle$ .

The rest of our proof will proceed analogously to that proof of Némethi [26], too. (More precisely, the first order part of this proof is analogous to the one in [26]. Lifting this first order part to  $\Pi_1^1$  seems to involve new ideas.) We are writing up

formulas in the language of the structure  $\mathfrak{A} = \langle U, p, q, N, A, M \rangle$ , where  $A, M \subseteq U \times U$  are partial unary operations (and the rest are as above). To define  $\Phi_A$ , we introduce the abbreviation

$$+(v_0, v_1, v_2) \stackrel{\text{def}}{\iff} \exists x[(p(x) = v_0 \wedge q(x) = v_1 \wedge A(x) = v_2)].$$

Now

$$\Phi_A \stackrel{\text{def}}{=} (A.1) \wedge (A.2) \wedge (A.3) \wedge (A.4) \wedge (A.5) \wedge (A.6),$$

where

$$(A.1) \stackrel{\text{def}}{=} \forall v(p(v), q(v) \in N \rightarrow A(v) \in N)$$

$$(A.2) \stackrel{\text{def}}{=} \forall v(\forall x, y, z \in N)(+(x, y, z) \wedge +(x, y, v) \rightarrow z = v)$$

$$(A.3) \stackrel{\text{def}}{=} (\forall x, y, z \in N)(+(x, y, z) \leftrightarrow +(y, x, z))$$

$$(A.4) \stackrel{\text{def}}{=} (\forall x, y, z \in N)(+(x, p(y), p(z)) \rightarrow +(x, y, z))$$

$$(A.5) \stackrel{\text{def}}{=} (\forall x, y \in N)(y \notin \text{Dom}(p) \leftrightarrow +(x, y, x))$$

$$(A.6) \stackrel{\text{def}}{=} \forall v(v \in \text{Dom}(A) \leftrightarrow v \in \text{Dom}(M) \leftrightarrow (p(v) \in N \wedge q(v) \in N)).$$

(\*2) As in the case of  $\Phi_N$ , if  $\langle U, p, q \rangle$  is a standard pairing structure, then

$$\mathfrak{A} \stackrel{\text{def}}{=} \langle U, p, q, N, A, M \rangle \models \Phi_N \wedge \Phi_A \text{ implies } \langle N, p \upharpoonright N, +^U \rangle \cong \langle \omega, \text{pred}, + \rangle,$$

where

$$+^U \stackrel{\text{def}}{=} \{ \langle a, b, c \rangle : \langle a, b \rangle \in \text{Dom}(A) \text{ and } A(\langle a, b \rangle) = c \}.$$

The equational form of  $\Phi_A$  is ( $N, A$  and  $M$  are variables again)

$$e_2(N, A, M) \stackrel{\text{def}}{=} (2.1) \wedge (2.2) \wedge (2.3) \wedge (2.4) \wedge (2.5) \wedge (2.6),$$

where (2.1), ..., (2.6) correspond to (A.1), ..., (A.6), respectively.

$$(2.1) \stackrel{\text{def}}{=} (q \circ (p \cap q) \circ N \circ 1) \cap (p \circ (p \cap q) \circ N \circ 1) \leq A \circ (p \cap q) \circ N \circ (p \cap q)^{-1}$$

$$(2.2) \stackrel{\text{def}}{=} C^{-1} \circ C \leq 1', \text{ where } C \stackrel{\text{def}}{=} A \cap (q \circ (p \cap q) \circ N \circ 1) \cap (p \circ (p \cap q) \circ N \circ 1)$$

$$(2.3) \stackrel{\text{def}}{=} C^{-1} \circ ((p \circ q^{-1}) \cap (q \circ p^{-1})) \circ C \leq 1'$$

$$(2.4) \stackrel{\text{def}}{=} ((p \circ (p \cap q) \circ N \circ (p \cap q)^{-1} \circ p^{-1}) \cap (q \circ (p \cap q) \circ N \circ q^{-1})) \circ A \circ (p \cap q)^{-1} \leq C$$

$$(2.5) \stackrel{\text{def}}{=} (p \circ (p \cap q) \circ N \circ (p \cap q)^{-1}) \cap (q \circ D \circ 1) = (C \cap p),$$

where  $D \stackrel{\text{def}}{=} [1 \circ (N - (p \circ 1))] \cap [(N - (p \circ 1)) \circ 1]$

$$(2.6) \stackrel{\text{def}}{=} A \circ 1 = M \circ 1 = (p \circ (p \cap q) \circ N \circ 1) \cap (q \circ (p \cap q) \circ N \circ 1).$$

Now, exactly as in the case of  $\Phi_N$ ,  $\langle \mathfrak{Rc}(U), p, q \rangle \models (e_1 \wedge e_2)[\text{Id} \upharpoonright N, A, M]$  iff  $\mathfrak{A} \models \Phi_N \wedge \Phi_A$ , where  $\mathfrak{A}$  is as in (\*2), and  $(e_1 \wedge e_2)[\text{Id} \upharpoonright N, A, M]$  is an evaluated equation, i.e., variables are replaced by concrete elements of the TPA in question.

Let

$$m(x, y, z) \stackrel{\text{def}}{\iff} (\exists v)(p(v) = x \wedge q(v) = y \wedge M(v) = z).$$

To ensure that  $m$  is real multiplication, we let (analogously to the definition of  $\Phi_A$ )  $\Phi_M$  be the formula  $(M.1) \wedge \dots \wedge (M.5)$ , where

$$(M.1) \stackrel{\text{def}}{=} M : N \times N \longrightarrow N \text{ is onto}$$

$$(M.2) \stackrel{\text{def}}{=} \forall v(\forall x, y, z \in N)(m(x, y, z) \wedge m(x, y, v) \rightarrow z = v)$$

$$(M.3) \stackrel{\text{def}}{=} (\forall x, y \in N)(x \notin \text{Dom}(p) \leftrightarrow m(x, y, x))$$

$$(M.4) \stackrel{\text{def}}{=} (\forall x, y, z \in N)(m(x, y, z) \leftrightarrow m(y, x, z))$$

$$(M.5) \stackrel{\text{def}}{=} (\forall x, y, z \in N)(m(p(x), y, z) \rightarrow (\exists w)(+(z, y, w) \wedge m(x, y, w))).$$

If  $\mathfrak{A}$  is as in (\*2), then

$$(*3) \quad \mathfrak{A} \models \Phi_N \wedge \Phi_A \wedge \Phi_M \text{ implies } \langle N, p \upharpoonright N, +^U, m^U \rangle \cong \langle \omega, \text{pred}, +, \cdot \rangle, \text{ if } \langle U, p, q \rangle \text{ is a standard pairing structure.}$$

(M.4) can be divided into two parts (M.4a) and (M.4b) in the following way.

$$(\forall x, y, z \in N)(m(x, y, z) \leftrightarrow m(y, x, z)) \iff$$

$$\forall v(\forall x, y, z \in N)(m(x, y, z) \wedge m(y, x, v) \rightarrow z = v) \wedge$$

$$\forall v(\forall x, y, z \in N)(m(x, y, z) \leftrightarrow m(y, x, v)).$$

The equational form of  $\Phi_M$  is  $e_3(N, A, M) \stackrel{\text{def}}{=} (3.1) \wedge \dots \wedge (3.4a) \wedge (3.4b) \wedge (3.5)$ , where (3.1), ..., (3.5) correspond to (M.1), ..., (M.5), respectively.

$$(3.1) \stackrel{\text{def}}{=} M^{-1} \circ ((p \circ (p \cap q) \circ N \circ 1) \cap (q \circ (p \cap q) \circ N \circ 1)) = (p \cap q) \circ N \circ 1$$

$$(3.2) \stackrel{\text{def}}{=} (p \cap q) \circ N \circ (p \cap q)^{-1} \circ K \circ K^{-1} \leq 1',$$

where  $K \stackrel{\text{def}}{=} ((p \circ (p \cap q) \circ N \circ 1) \cap (q \circ (p \cap q) \circ N \circ 1)) \cap M$

$$(3.3) \stackrel{\text{def}}{=} (p \circ D) \cap (q \circ (p \cap q) \circ N \circ 1) = p \cap K$$

$$(3.4a) \stackrel{\text{def}}{=} (p \cap q) \circ N \circ (p \cap q)^{-1} \circ K^{-1} \circ ((p \circ q^{-1}) \cap (q \circ p^{-1})) \circ K \leq 1'$$

$$(3.4b) \stackrel{\text{def}}{=} ((p \circ q^{-1}) \cap (q \circ p^{-1})) \circ K \circ (p \cap q) \circ N \circ (p \cap q)^{-1} \leq K$$

$$(3.5) \stackrel{\text{def}}{=} J \leq ((J \circ p^{-1}) \cap (q \circ q^{-1})) \circ [((C \circ q^{-1}) \cap (p \circ p^{-1})) \cap \\ [((p \circ C^{-1}) \cap (q \circ q^{-1})) \circ ((p \circ (p \cap q)^{-1} \circ p^{-1}) \cap (q \circ q^{-1}) \circ M \circ q^{-1})] \circ p, \\ \text{where } J \stackrel{\text{def}}{=} ((p \circ (p \cap q) \circ N \circ p^{-1}) \cap (q \circ (p \cap q) \circ N \circ (p \cap q)^{-1} \circ q^{-1})) \circ \\ M \circ (p \cap q) \circ N \circ (p \cap q)^{-1}, \text{ i.e., } J \text{ denotes those pairs } \langle \langle x, y \rangle, z \rangle \\ \text{for which } x, y, z \in N \text{ and } m(p(x), y, z).$$

Exactly as in the case of  $e_2$ , this equation  $e_3(N, A, M)$  is such that

$$\langle \mathfrak{Re}(U), p, q \rangle \models (e_1 \wedge e_2 \wedge e_3)[\text{Id} \upharpoonright N, A, M] \text{ iff } \mathfrak{A} \models \Phi_N \wedge \Phi_A \wedge \Phi_M.$$

Let  $\mathfrak{B} \subseteq \langle \mathfrak{Re}(U), p, q \rangle$  be any TPA with  $X, A, M \in B$  such that  $\mathfrak{B} \models (e_1 \wedge e_2 \wedge e_3)[X, A, M]$ . Let  $N = \text{Dom}(X)$ . Then  $X = \text{Id} \upharpoonright N$  and hence  $\langle \mathfrak{Re}(U), p, q \rangle \models (e_1 \wedge e_2 \wedge e_3)[\text{Id} \upharpoonright N, A, M]$ , thus  $\langle U, p, q, N, A, M \rangle \models \Phi_N \wedge \Phi_A \wedge \Phi_M$  proving

$$(*4) \langle N, p \upharpoonright N, +^U, m^U \rangle \cong \Omega$$

by (\*3) and the above properties of  $e_1 \wedge e_2 \wedge e_3$ . Summing up, we have (\*5) below, where  $\text{baseof}(\mathfrak{B})$  denotes the set  $U$  for which  $\mathfrak{B} = \langle \mathfrak{Re}(U), \dots \rangle$ .

(\*5) For any TPA  $\mathfrak{B}$  and  $X, A, M \in B$  we have that if  $\mathfrak{B} \models e_1 \wedge e_2 \wedge e_3[X, A, M]$  then  $\langle \text{baseof}(\mathfrak{B}), p^{\mathfrak{B}}, q^{\mathfrak{B}}, \text{Dom}(X), A, M \rangle \models \Phi_N \wedge \Phi_A \wedge \Phi_M$ , so  $\text{Dom}(X), p^{\mathfrak{B}}, A, M$  code an isomorphic copy of the standard model  $\Omega$  of arithmetic as indicated in (\*4).

When applying (\*5), we will need (5.1) below known from the literature. Let  $\langle U, p, q \rangle$  be a standard pairing structure and  $X, A, M, P^+ \in \mathcal{P}(U \times U)$  with  $X \cup P^+ \subseteq \text{Id}$ . Let  $N = \text{Dom}(X)$  and  $P = \text{Dom}(P^+)$  and  $\phi$  be a first order formula in the language of  $\langle U, p, q, N, A, M, P \rangle$ . Now we claim that

(5.1) there is a recursive function associating to every first order formula  $\phi$  of the above kind a TPA-equation  $e_\phi(X, A, M, P^+)$  such that for every  $N, P \subseteq U$  and  $A, M \subseteq U \times U$ ,

$$\langle \mathfrak{Re}(U), p, q \rangle \models e_\phi[\text{Id} \upharpoonright N, A, M, \text{Id} \upharpoonright P] \text{ iff } \langle U, p, q, N, A, M, P \rangle \models \phi.$$

The above (5.1) is proved in [40, §4.5], see the theorem on p.126 (there it is stated in a different form, see also §7.1) as well as in [25, Lemma 3, p.35] (where the result is stated in the present form; our  $e_\phi$  is denoted there as  $g(\phi) = 1$  because  $g(\phi)$  is a TPA-term there).

It is well known that every statement of arithmetic of the form  $\forall P_1 \dots \forall P_n \forall F_1 \dots \forall F_m \phi$  can be equivalently expressed by a formula of the form

$\forall P\psi$ , where both  $\phi$  and  $\psi$  are first order formulas,  $P_1, \dots, P_n, P$  are predicate variables, and  $F_1, \dots, F_m$  are function variables. (If  $F_i : {}^k N \rightarrow N$ , then this  $F_i$  can be replaced by a predicate variable which has  $k + 1$  arguments. See Rogers [29, §16.1 Thm.III] for the rest.) Moreover,  $P$  can be replaced by a one argument predicate variable. After this reduction, for any  $\Pi_1^1$  sentence  $\psi$  (of the form  $\forall P\phi$ , where  $\phi$  is first order) of the language of  $\Omega$  we construct  $\psi^+$  ( $= \forall P\phi^+$ , where  $\phi^+$  is first order again) in the language of  $\langle U, p, q, N, A, M \rangle$  such that

$$(*6) \quad \Omega \models \psi \text{ iff } \langle U, p, q, N, A, M \rangle \models (\Phi_N \wedge \Phi_A \wedge \Phi_M \longrightarrow \psi^+), \text{ for any standard pairing structure } \langle U, p, q \rangle \text{ and any choice of } N, A, M.$$

Indeed, it is easy to see that  $\psi^+$  can be defined by induction on the complexity of  $\psi$ , e.g., let  $(i = j + k)^+$  be  $\exists v(A(v) = i \wedge p(v) = j \wedge q(v) = k) \dots$  and write  $p(v)$  for  $\text{pred}(v)$ . Now apply (\*3).

Then, by (\*5) and (\*6) we have that

$$\Omega \models \psi \quad \text{iff}$$

$$(*7) \quad (\forall \mathfrak{B} \in \text{TPA})(\forall X, A, M \in B)(\mathfrak{B} \models (e_1 \wedge e_2 \wedge e_3)[X, A, M] \Rightarrow \langle \text{baseof}(\mathfrak{B}), p^{\mathfrak{B}}, q^{\mathfrak{B}}, \text{Dom}(X), A, M \rangle \models \psi^+).$$

As each possible value of  $P$  (in the formula  $\forall P\phi$ ) is a subset of  $\omega$ , to  $\psi^+ = \forall P\phi^+$  we associate a quasi-equation  $e_\psi = (P^+ \leq X \wedge X \leq \text{Id}) \rightarrow e_\phi$ , where  $e_\phi$  (or, more precisely,  $e_{\phi^+}$ ) is the TPA-equation corresponding to  $\phi^+$  according to (5.1) above. Let  $\mathfrak{B}, X, A, M$  be as in (\*7). Since  $\psi^+$  is  $\forall P\phi^+$ , we have

$$\langle \text{baseof}(\mathfrak{B}), p^{\mathfrak{B}}, q^{\mathfrak{B}}, \text{Dom}(X), A, M \rangle \models \psi^+ \text{ iff } (\forall P \leq \text{Dom}(X)) \langle \text{baseof}(\mathfrak{B}), p^{\mathfrak{B}}, q^{\mathfrak{B}}, \text{Dom}(X), A, M, P \rangle \models \phi^+.$$

By (5.1), the right hand side holds iff for our equation  $e_\phi$  (associated to  $\phi^+$ )

$$\mathfrak{B} \models (P^+ \leq X \rightarrow e_\phi[X, A, M, \text{Id} \upharpoonright P]),$$

which is equivalent to

$$\mathfrak{B} \models \forall P^+ e_\psi[X, A, M].$$

So we have  $(\forall \mathfrak{B} \in \text{TPA})(\forall X, A, M \in B)$

$$(\langle \text{baseof}(\mathfrak{B}), p^{\mathfrak{B}}, q^{\mathfrak{B}}, \text{Dom}(X), A, M \rangle \models \psi^+ \text{ iff } \mathfrak{B} \models \forall P^+ e_\psi[X, A, M]).$$

Putting this together with (\*7), we get that

$$\begin{aligned} \Omega \models \psi \quad \text{iff} \\ (\forall \mathfrak{B} \in \text{TPA})(\forall X, A, M, P^+ \in B)(\mathfrak{B} \models (e_1 \wedge e_2 \wedge e_3)[X, A, M] \Rightarrow \mathfrak{B} \models e_\psi[X, A, M, P^+]). \end{aligned}$$

But this means exactly that

(\*8)  $\Omega \models \psi$  iff  $\text{TPA} \models (e_1 \wedge e_2 \wedge e_3)(X, A, M) \rightarrow e_\psi(X, A, M, P^+)$ .

Now, by the above result of Jónsson, there is an equation  $e_\psi^+$  such that  $\text{TPA} \models (e_1 \wedge e_2 \wedge e_3) \rightarrow e_\psi$  iff  $\text{TPA} \models e_\psi^+$  and the function  $\{\langle e_\psi, e_\psi^+ \rangle : \psi \text{ is a formula in the language of } \Omega\}$  is computable. So, by (\*8), we have

(\*9)  $\Omega \models \psi$  iff  $\text{TPA} \models e_\psi^+$

and there is a computable function  $f$  with  $e_\psi^+ = f(\psi)$  for any formula  $\psi$  in the  $\Pi_1^1$  language of  $\Omega$ . This  $f$  is computable because Lemma 3 of Némethi [25] explicitly says so (and the rest of our above construction, including Jónsson's, are clearly computable). Thus we have

$$\langle \omega, \text{pred}, +, \cdot \rangle \models \psi \iff \text{TPA} \models e_\psi$$

which completes our proof.  $\square$

## 5 The complexity of $\text{Eq}(\text{TPA})$ is at most $\Pi_1^1$

The following theorem states that  $\text{Eq}(\text{TPA})$  is at most on the  $\Pi_1^1$  level, and so, by Theorem 4.1, it is exactly on the  $\Pi_1^1$  level.

**5.1 THEOREM** *Let  $e$  be an equation in the language of TPA's. Then there is a  $\Pi_1^1$  sentence  $\psi_e$  (effectively calculable from  $e$ ) such that*

(\*10)  $\text{TPA} \models e \iff \Omega \models \psi_e$ .

*Proof.* We will use the well known and convenient  $\lambda$ -notation to form expressions of the language of arithmetic. So let  $r$  be the pairing function  $\lambda x \lambda y. 2^x \cdot 3^y$  on  $\omega$ . From now on, let  $\mathbf{U}_0$ ,  $\mathbf{U}$ ,  $\mathbf{K}$ ,  $\mathbf{O}$ ,  $\mathbf{I}$ ,  $\mathbf{P}$ , and  $\mathbf{Q}$  be predicate (thus second order) variables. We are listing a few formulas that restrict the possible meanings of them so that they can be thought of as representatives of a set of non-pair elements of a TPA, base, (Boolean) identity, (Boolean) zero, multiplicative identity, and projection elements, respectively.

$$\varphi_0 \stackrel{\text{def}}{=} \forall x (\mathbf{U}_0(x) \rightarrow (\neg \exists y \exists z (x = r(y, z))))$$

$$\varphi_1 \stackrel{\text{def}}{=} \forall x (\mathbf{U}(x) \leftrightarrow (\mathbf{U}_0(x) \vee \exists y \exists z (\mathbf{U}(y) \wedge \mathbf{U}(z) \wedge (x = r(y, z))))))$$

$$\varphi_2 \stackrel{\text{def}}{=} \forall x (\mathbf{K}(x) \leftrightarrow \exists y \exists z (\mathbf{U}(y) \wedge \mathbf{U}(z) \wedge (x = r(y, z))))$$

$$\varphi_3 \stackrel{\text{def}}{=} \forall x (\neg \mathbf{O}(x))$$

$$\varphi_4 \stackrel{\text{def}}{=} \forall x (\mathbf{P}(x) \leftrightarrow \exists y \exists z (\mathbf{U}(y) \wedge \mathbf{U}(z) \wedge (x = r(r(y, z), y))))$$



$$\varphi_5 \stackrel{\text{def}}{=} \forall x(\mathbf{Q}(x) \leftrightarrow \exists y \exists z(\mathbf{U}(y) \wedge \mathbf{U}(z) \wedge (x = r(r(y, z), z))))$$

$$\varphi_6 \stackrel{\text{def}}{=} \forall x(\mathbf{I}(x) \leftrightarrow \exists y(\mathbf{U}(y) \wedge (x = r(y, y)))).$$

Assume that  $\mathbf{U}_0, \dots, \mathbf{Q}$  are the first seven predicate variables of the language of arithmetic, the others are enumerated as  $\mathbf{V}_1, \dots, \mathbf{V}_n, \dots$ . If  $e$  is  $\tau_1 = \tau_2$  then let  $k \in \omega$  be such that  $\tau_1, \tau_2 \in TRM_{\text{TPA}}^k$  (TPA terms such that, for  $i > k$ ,  $v_i$  does not occur in them).

Then  $\psi_e$  will be of the form

$$\forall \mathbf{U}_0, \mathbf{U}, \mathbf{K}, \mathbf{O}, \mathbf{I}, \mathbf{P}, \mathbf{Q}, \mathbf{V}_1 \dots \mathbf{V}_k ((\varphi_0 \wedge \dots \wedge \varphi_6 \\ \wedge \forall x[(\mathbf{V}_1(x) \vee \dots \vee \mathbf{V}_k(x)) \rightarrow \mathbf{K}(x)]) \rightarrow (\tau_1^* = \tau_2^*)),$$

where  $\tau_i^*$  is the one-place predicate of arithmetic obtained from  $\tau_i$  by applying to it the translating function  $*$  given below.

DEFINITION 5.1.1. The translating function  $*$  :  $TRM_{\text{TPA}}^k \rightarrow PRED_A^1$  (where  $PRED_A^1$  is the set of one place predicates of the language of arithmetic) is defined by the following clauses:

$$1^* = \mathbf{K}, \quad 0^* = \mathbf{O}, \quad 1'^* = \mathbf{I}, \quad p^* = \mathbf{P}, \quad q^* = \mathbf{Q}$$

$$v_i^* = \mathbf{V}_i$$

$$[\sim \tau]^* = \lambda x(\mathbf{K}(x) \wedge \neg \tau^*(x))$$

$$[\tau \cup \mu]^* = \lambda x(\tau^*(x) \vee \mu^*(x))$$

$$[\tau \cap \mu]^* = \lambda x(\tau^*(x) \wedge \mu^*(x))$$

$$[\tau^{-1}]^* = \lambda x \exists y \exists z ((x = r(z, y)) \wedge \tau^*(r(y, z)))$$

$$[\tau \circ \mu]^* = \lambda x \exists y \exists z \exists w ((x = r(y, w)) \wedge \tau^*(r(y, z)) \wedge \mu^*(r(z, w))).$$

$\Delta$

(Note that for any  $\tau \in TRM_{\text{TPA}}^k$ , if  $\tau^*$  is not simply a predicate variable, then it is of the form  $\lambda x.\chi(x)$ , where  $\chi$  is a second order formula with no  $\lambda$ 's or second order quantifiers in it. Thus  $\tau_1^* = \tau_2^*$  is (logically) equivalent to  $\forall x(\chi_1 \leftrightarrow \chi_2)$ . That is,  $\psi_e$  is indeed  $\Pi_1^1$ .) Clearly, for any  $\tau \in TRM_{\text{TPA}}^k$  we have

$$\Omega \models \forall \mathbf{U}_0, \mathbf{U}, \mathbf{K}, \mathbf{O}, \mathbf{I}, \mathbf{P}, \mathbf{Q}, \mathbf{V}_1 \dots \mathbf{V}_k \\ ((\varphi_0 \wedge \dots \wedge \varphi_6 \wedge \forall x[(\mathbf{V}_1(x) \vee \dots \vee \mathbf{V}_k(x)) \rightarrow \mathbf{K}(x)]) \rightarrow \forall x(\tau^*(x) \rightarrow \mathbf{K}(x))).$$

Now that the construction of  $\psi_e$  is complete, we can start proving (\*10).

( $\Rightarrow$ ) Suppose that  $\Omega \not\models \psi_e$ , that is, there is an expansion

$$\Omega^+ = \langle \Omega, U_0, U, K, O, I, P, Q, V_1, \dots, V_k \rangle$$

of  $\Omega$  such that

$$\Omega^+ \models \varphi_0 \wedge \dots \wedge \varphi_6 \wedge \forall x((\mathbf{V}_1(x) \vee \dots \vee \mathbf{V}_k(x)) \rightarrow \mathbf{K}(x)),$$

but

$$(\tau_1^*)^{\Omega^+} \neq (\tau_2^*)^{\Omega^+}.$$

We will construct a TPA

$$\mathfrak{A} = \langle \mathcal{P}(U' \times U'), \cup, \cap, \sim, 0, 1, \circ, ^{-1}, 1', p, q \rangle$$

in which the equation  $e$  does not hold.

Let  $U'_0$  be any set consisting of elements that are not pairs (in the set-theoretical sense), and such that there is a bijection  $f' : U'_0 \rightarrow U_0$ . Let  $U'$  be the closure of  $U'_0$  under forming ordered pairs (as in the first part of the paper), and let  $f$  be the  $U' \rightarrow U$  function defined by

$$f(u) \stackrel{\text{def}}{=} \begin{cases} f'(u), & \text{if } u \in U_0 \\ r(f(u_1), f(u_2)), & \text{if } u = \langle u_1, u_2 \rangle. \end{cases}$$

Then  $f$  is one-one and onto because  $\varphi_0$  and  $\varphi_1$  are true (in  $\Omega^+$ ). Let  $g : U' \times U' \rightarrow K$  and  $h : \mathcal{P}(U' \times U') \rightarrow \mathcal{P}(K)$  be defined by

$$g(\langle u_1, u_2 \rangle) \stackrel{\text{def}}{=} r(f(u_1), f(u_2)) \quad \text{for all } \langle u_1, u_2 \rangle \in U' \times U',$$

$$h(X) \stackrel{\text{def}}{=} \{g(x) : x \in X\} \quad \text{for all } X \subseteq U' \times U'.$$

Then  $g$  and thus  $h$  are one-one and onto because  $\varphi_2$  is true.

**PROPOSITION 5.1.2.** If the values of the TPA variables  $v_i$  ( $i \leq k$ ) are chosen to be  $h^{-1}(V_i)$  (and this makes sense since  $h^{-1}(V_i) \subseteq U' \times U'$ ), then for all  $\tau \in TRM_{\text{TPA}}^k$ , we have

$$h(\tau^{\mathfrak{A}}) = (\tau^*)^{\Omega^+}.$$

*Proof of Prop.5.1.2.* By induction on  $TRM_{\text{TPA}}^k$  using the bijectivity of  $f$ ,  $g$ , and  $h$  and the assumption that  $\varphi_3, \dots, \varphi_6$  are true in  $\Omega^+$ .

$$h(1^{\mathfrak{A}}) = h(U' \times U') = \{g(x) : x \in U' \times U'\} = (1^*)^{\Omega^+},$$

$$h(0^{\mathfrak{A}}) = h(\emptyset) = \emptyset = (0^*)^{\Omega^+},$$

$$\begin{aligned} h(1'^{\mathfrak{A}}) &= \{g(\langle u, u \rangle) : u \in U'\} = \{r(f(u), f(u)) : u \in U'\} \\ &= \{r(x, x) : x \in U\} = I = (1'^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h(p^{\mathfrak{A}}) &= \{g(\langle \langle u, v \rangle, u \rangle) : u, v \in U'\} \\ &= \{r(r(f(u), f(v)), f(u)) : u, v \in U'\} = \{r(r(y, z), y) : y, z \in U\} \\ &= P = (p^*)^{\Omega^+}. \end{aligned}$$

The proof of  $h(q^{\mathfrak{A}}) = (q^*)^{\Omega^+}$  is similar to that of  $h(p^{\mathfrak{A}}) = (p^*)^{\Omega^+}$ , and  $h(v_i^{\mathfrak{A}}) = V_i = (v_i^*)^{\Omega^+}$  by definition.

Suppose that the statement holds for some  $\tau, \mu \in TRM_{\text{TPA}}^k$ . Then

$$\begin{aligned} h([\sim \tau]^{\mathfrak{A}}) &= h(\{v \in U' \times U' : v \notin \tau^{\mathfrak{A}}\}) \\ &= \{g(v) : v \in U' \times U' \text{ and } g(v) \notin h(\tau^{\mathfrak{A}})\} \\ &= \{x \in K : x \notin (\tau^*)^{\Omega^+}\} = ([\sim \tau]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau \cup \mu]^{\mathfrak{A}}) &= h(\{v \in U' \times U' : v \in \tau^{\mathfrak{A}} \cup \mu^{\mathfrak{A}}\}) \\ &= \{g(v) : g(v) \in (\tau^*)^{\Omega^+} \cup (\mu^*)^{\Omega^+}\} \\ &= \{x : x \in (\tau^*)^{\Omega^+} \cup (\mu^*)^{\Omega^+}\} = ([\tau \cup \mu]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau^{-1}]^{\mathfrak{A}}) &= h(\{\langle v, u \rangle : \langle u, v \rangle \in \tau^{\mathfrak{A}}\}) \\ &= \{r(f(v), f(u)) : r(f(u), f(v)) \in (\tau^*)^{\Omega^+}\} \\ &= \{r(z, y) : r(y, z) \in (\tau^*)^{\Omega^+}\} = ([\tau^{-1}]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau \circ \mu]^{\mathfrak{A}}) &= h(\{\langle u, v \rangle : \exists s(\langle u, s \rangle \in \tau^{\mathfrak{A}} \text{ and } \langle s, v \rangle \in \mu^{\mathfrak{A}})\}) \\ &= \{r(f(u), f(v)) : \exists s(r(f(u), f(s)) \in (\tau^*)^{\Omega^+} \text{ and} \\ &\quad r(f(s), f(v)) \in (\mu^*)^{\Omega^+})\} \\ &= \{r(x, y) : \exists z(r(x, z) \in (\tau^*)^{\Omega^+} \text{ and } r(z, y) \in (\mu^*)^{\Omega^+})\} \\ &= ([\tau \circ \mu]^*)^{\Omega^+}. \end{aligned}$$

This completes the proof of Prop.5.1.2.  $\square$

Now, with the valuation (of the TPA-variables) given in the proposition, we have

$$h(\tau_1^{\mathfrak{A}}) = (\tau_1^*)^{\Omega^+} \neq (\tau_2^*)^{\Omega^+} = h(\tau_2^{\mathfrak{A}}),$$

so that  $\mathfrak{A} \not\models \tau_1 = \tau_2$ , and thus

$$\text{TPA} \not\models \tau_1 = \tau_2.$$

( $\Leftarrow$ ) The proof relies on the fact that by the choice of the pairing function  $r$  there are infinitely many elements that are not “pairs” (i.e., are not elements of  $\text{Rng}(r)$ ).

Suppose that TPA  $\not\models e$ . Then (cf. [35]) there is a TPA

$$\mathfrak{A} = \langle \mathcal{P}(U' \times U'), \cup, \cap, \sim, 0, 1, ^{-1}, \circ, 1', p, q \rangle$$

such that  $U'$  (and thus  $U'_0$ , the non-pair elements of  $U'$ ) is countable and  $e$  fails in it. Let us fix  $\mathfrak{A}$  and define an expansion

$$\Omega^+ = \langle \Omega, U_0, U, K, O, I, P, Q, V_1, \dots, V_k \rangle$$

of  $\Omega$  such that

$$\Omega^+ \models \varphi_0 \wedge \dots \wedge \varphi_6 \wedge \forall x ((\mathbf{V}_1(x) \vee \dots \vee \mathbf{V}_k(x)) \rightarrow \mathbf{K}(x)),$$

but

$$(\tau_1^*)^{\Omega^+} \neq (\tau_2^*)^{\Omega^+}.$$

The pairing function  $r$  was defined in such a way that there are infinitely many “no-pairs” in  $\omega$  (say the powers of 5). Let  $U_0$  be any subset of  $\omega$  whose cardinality equals that of  $U'_0$  and is such that  $U_0 \cap \text{Rng}(r) = \emptyset$ . Then

$$\Omega^+ \models \varphi_0.$$

If  $f' : U'_0 \rightarrow U_0$  is one-one and onto, we define

$$f(u) \stackrel{\text{def}}{=} \begin{cases} f'(u), & \text{if } u \in U'_0 \\ r(f(u_1), f(u_2)), & \text{if } u = \langle u_1, u_2 \rangle, u_1, u_2 \in U' \end{cases}$$

and

$$U \stackrel{\text{def}}{=} \text{Rng}(f).$$

**PROPOSITION 5.1.3.**  $f \in {}^{U'}U$ ,  $f$  is one-one and onto.

*Proof* of Prop.5.1.3. The fact that  $f$  is defined for all  $u \in U'$  can be proved by induction on the construction of  $U'$  (from  $U'_0$ ).  $f$  is clearly onto, so it remains to prove that it is one-one.

Suppose that  $f(u) = f(v)$  for some  $u, v \in U'$ . It is clear that both  $u$  and  $v$  are either in  $U'_0$  or in  $U' \setminus U'_0$ , otherwise the image of, say  $u$ , would be a “pair”, i.e., an element of  $\text{Rng}(r)$ , but not that of  $v$ , contradicting our assumption. If  $u, v \in U'_0$  then  $u = v$  follows from  $f$ 's being one-one. Let  $u = \langle u_1, u_2 \rangle$ ,  $v = \langle v_1, v_2 \rangle$  be such that  $f(u) = f(v)$  is the least  $n \in U$  which has more than one pre-image in  $U'$ . Since  $f(u_1) = f(v_1) < f(u)$  we have  $u_1 = v_1$ , a contradiction.  $\square$

**PROPOSITION 5.1.4.**  $\Omega^+ \models \varphi_1$ .

*Proof* of Prop.5.1.4. ( $\rightarrow$ ) If  $x \in U$  but  $x \notin U_0$ , then  $x \in \text{Rng}(f) \setminus \text{Rng}(f')$ , whence  $x = r(f(u), f(v))$  for some  $u, v \in U'$ , that is  $x = r(y, z)$  for some  $y, z \in U$ .

( $\leftarrow$ ) If  $x \in U_0$  then clearly  $x \in U$ ; if  $x = r(y, z)$  for some  $y, z \in U$  then  $x = f(\langle f^{-1}(y), f^{-1}(z) \rangle) \in U$ .  $\square$

Define  $g$  by

$$g(\langle u_1, u_2 \rangle) \stackrel{\text{def}}{=} r(f(u_1), f(u_2)) \text{ for all } u_1, u_2 \in U'$$

and let

$$K \stackrel{\text{def}}{=} \text{Rng}(g).$$

Then  $g : U' \times U' \longrightarrow K$  is one-one and onto.

PROPOSITION 5.1.5.  $\Omega^+ \models \varphi_2$ .

*Proof* of Prop.5.1.5. ( $\rightarrow$ )  $x \in K$  implies  $x = g(\langle u, v \rangle) = r(f(u), f(v))$  for some  $u, v \in U'$ . But  $f(u), f(v) \in U$ .

( $\leftarrow$ ) If  $x = r(y, z)$  for some  $y, z \in U$ , then  $x = g(\langle f^{-1}(y), f^{-1}(z) \rangle) \in K$ .  $\square$

Let  $h : \mathcal{P}(U' \times U') \rightarrow \mathcal{P}(K)$  be the mapping induced by  $g$ , that is,

$$h(X) \stackrel{\text{def}}{=} \{g(x) : x \in X\} \text{ for all } X \subseteq U' \times U'.$$

Then  $h$  is one-one and onto, and

$$h(1^{\mathfrak{A}}) = h(U' \times U') = K = (1^*)^{\Omega^+}.$$

Define

$$O \stackrel{\text{def}}{=} h(0^{\mathfrak{A}})$$

$$P \stackrel{\text{def}}{=} h(p^{\mathfrak{A}})$$

$$Q \stackrel{\text{def}}{=} h(q^{\mathfrak{A}})$$

$$I \stackrel{\text{def}}{=} h([1']^{\mathfrak{A}})$$

$$V_i \stackrel{\text{def}}{=} h(v_i^{\mathfrak{A}}) \quad \text{if } 1 \leq i \leq k.$$

PROPOSITION 5.1.6.

$$\Omega^+ \models \varphi_3 \wedge \cdots \wedge \varphi_6 \wedge \forall x((\mathbf{V}_1(x) \vee \cdots \vee \mathbf{V}_k(x)) \rightarrow \mathbf{K}(x)).$$

*Proof* of Prop.5.1.6. In the course of the proof we will make use of the fact that  $f$  is onto.  $h(0^{\mathfrak{A}}) = h(\emptyset) = \emptyset$ , thus  $\varphi_3$  is true.

$h(p^{\mathfrak{A}}) = h(\{\langle \langle u, v \rangle, u \rangle : u, v \in U'\}) = \{r(r(f(u), f(v)), f(u)) : u, v \in U'\} = \{r(r(y, z), y) : y, z \in U\}$ , thus  $\varphi_4$  is true, and a similar argument proves  $\varphi_5$ .

$\varphi_6$  is true since  $h([1']^{\mathfrak{A}}) = h(\{\langle u, u \rangle : u \in U'\}) = \{r(f(u), f(u)) : u \in U'\} = \{r(y, y) : y \in U\}$ .

Finally,  $h(v_i^{\mathfrak{A}}) \subseteq K$  for all  $1 \leq i \leq k$ .  $\square$

In order to show that  $(\tau_1^*)^{\Omega^+} \neq (\tau_2^*)^{\Omega^+}$  we need to establish the following

PROPOSITION 5.1.7. If  $\tau \in TRM_{\text{TPA}}^k$ , then

$$h(\tau^{\mathfrak{A}}) = (\tau^*)^{\Omega^+}.$$

*Proof* of Prop.5.1.7. We have already seen that  $h(1^{\mathfrak{A}}) = (1^*)^{\Omega^+}$ . For the other atomic terms  $\tau$  in  $TRM_{\text{TPA}}^k$ , we have  $h(\tau^{\mathfrak{A}}) = (\tau^*)^{\Omega^+}$  by definition (of  $\Omega^+$ ), so suppose that the statement holds for  $\tau, \mu \in TRM_{\text{TPA}}^k$ . Then

$$\begin{aligned} h([\sim \tau]^{\mathfrak{A}}) &= h(\{u \in U' \times U' : u \notin \tau^{\mathfrak{A}}\}) = \{g(u) : u \in U' \times U' \text{ and } u \notin \tau^{\mathfrak{A}}\} \\ &= \{x \in K : x \notin (\tau^*)^{\Omega^+}\} = (\lambda x(\mathbf{K}(x) \wedge \neg \tau^*(x)))^{\Omega^+} = ([\sim \tau]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau \cup \mu]^{\mathfrak{A}}) &= h(\{u \in U' \times U' : u \in \tau^{\mathfrak{A}} \cup \mu^{\mathfrak{A}}\}) \\ &= \{g(u) : g(u) \in (\tau^*)^{\Omega^+} \cup (\mu^*)^{\Omega^+}\} \\ &= \{x : x \in (\tau^*)^{\Omega^+} \cup (\mu^*)^{\Omega^+}\} = (\lambda x(\tau^*(x) \vee \mu^*(x)))^{\Omega^+} \\ &= ([\tau \cup \mu]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau^{-1}]^{\mathfrak{A}}) &= h(\{\langle v, u \rangle : \langle u, v \rangle \in \tau^{\mathfrak{A}}\}) \\ &= \{r(f(v), f(u)) : r(f(u), f(v)) \in (\tau^*)^{\Omega^+}\} \\ &= \{r(z, y) : r(y, z) \in (\tau^*)^{\Omega^+}\} \\ &= (\lambda x \exists y \exists z (x = r(y, z) \wedge \tau^*(r(y, z))))^{\Omega^+} \\ &= ([\tau^{-1}]^*)^{\Omega^+}, \end{aligned}$$

$$\begin{aligned} h([\tau \circ \mu]^{\mathfrak{A}}) &= h(\{\langle u, v \rangle : \exists s (\langle u, s \rangle \in \tau^{\mathfrak{A}} \text{ and } \langle s, v \rangle \in \mu^{\mathfrak{A}})\}) \\ &= \{r(f(u), f(v)) : \exists s (r(f(u), f(s)) \in (\tau^*)^{\Omega^+} \text{ and } \\ &\quad r(f(s), f(v)) \in (\mu^*)^{\Omega^+})\} \\ &= \{r(x, y) : \exists z (r(x, z) \in (\tau^*)^{\Omega^+} \text{ and } r(z, y) \in (\mu^*)^{\Omega^+})\} \\ &= (\lambda w \exists x \exists y \exists z (w = r(x, y) \wedge \tau^*(r(x, z)) \wedge \mu^*(r(z, y))))^{\Omega^+} \\ &= ([\tau \circ \mu]^*)^{\Omega^+}. \quad \square \end{aligned}$$

We conclude that

$$(\tau_1^*)^{\Omega^+} = h(\tau_1^{\mathfrak{A}}) \neq h(\tau_2^{\mathfrak{A}}) = (\tau_2^*)^{\Omega^+},$$

so that

$$\Omega^+ \not\models \psi_e.$$

We have proved Theorem 5.1  $\square$

## 6 Eq( $\text{TPA}^-$ ) is recursively enumerable

Next we show that the RA-reducts of TPA's (called  $\text{TPA}^-$ ) are much simpler, i.e., they admit a recursive axiomatization. This is so because the class  $\text{TPA}^-$  coincides with the class of those QRA's whose base is not a singleton. In more detail:

**6.1 DEFINITION**  $\mathfrak{A} = \langle \mathcal{P}(U \times U), \dots \rangle$  is a  $\text{TPA}^-$  iff  $\langle \mathcal{P}(U \times U), \dots, p, q \rangle$  is a TPA.  $\triangle$

**6.2 LEMMA** Let  $\mathfrak{A}$  and  $\mathfrak{A}'$  be full RRA's (see [15, §5.3.2]) and suppose that their bases have the same cardinality. Then  $\mathfrak{A} \cong \mathfrak{A}'$ .

*Proof.* The mapping induced by the function  $f : U \longrightarrow U'$  is clearly an RA isomorphism.  $\square$

**6.3 LEMMA**  $U$  is empty or  $|U| \geq \omega$  iff there is an  $U'$  with  $|U| = |U'|$  and  $U' \times U' \subseteq U'$ .

**6.4 LEMMA**  $\mathfrak{A}$  is a full RRA with empty or infinite base iff there is a  $\mathfrak{B} \in \text{TPA}^-$  such that  $\mathfrak{A} \cong \mathfrak{B}$ . Thus

$$\mathbf{I}\{\text{full } \text{TPA}^-\} = \mathbf{I}\{\text{full RRA with empty or infinite base}\}.$$

*Proof.* By Lemmas 6.2 and 6.3.  $\square$

Now consider the class of Q-Relation Algebras introduced in [40, 8.4]. Since  $\text{QRA} \subseteq \text{RRA}$  by [40, 8.4 (iii)], it makes sense to speak of full QRA's, and by [40, 8.4 (iv)] we have

$$(11) \quad \{\text{full QRA's with nonsingleton base}\} = \{\text{full RRA's with empty or infinite base}\}.$$

Combined with Lemma 6.4, this gives

$$(12) \quad \text{TPA}^- \models e \quad \text{iff} \quad \{\text{full QRA}\} \models [(0=1) \vee (1 \neq \text{Id})] \rightarrow e.$$

**6.5 LEMMA**  $\text{QRA} = \mathbf{SP}\{\text{full QRA}\}$ .

*Proof.* Let  $\mathfrak{A} \in \text{QRA}$ . Then there are  $I$  and full RRA's  $\mathfrak{B}_i$  ( $i \in I$ ) with

$$\mathfrak{A} \subseteq \mathfrak{A}^+ \cong \prod_{i \in I} \mathfrak{B}_i.$$

Now  $\mathfrak{A}^+$  has conjugated quasiprojections in its universe (since  $\mathfrak{A}$  has), and thus the  $\mathfrak{B}_i$ 's (being homomorphic images of  $\mathfrak{A}$ ) are (full) QRA's. (Recall that the property of being a pair of conjugated quasiprojection is defined by an equation.)  $\square$

Since Eq(QRA) is recursively enumerable, we have

**6.6 THEOREM**  $\text{Eq}(\text{TPA}^-)$  is recursively enumerable.

*Proof.* By (12) and Lemma 6.5,  $\text{TPA}^- \models e$  iff  $\text{QRA} \models (0=1) \vee (1 \neq \text{Id}) \rightarrow e$ .  $\square$

**6.7 THEOREM**  $\text{TPA}^-$  is not finitely based.

*Proof.* By the above Lemma 6.4 and (11) we have

$$\mathbf{I}\{\text{full QRA}\} = \mathbf{I}(\{\text{full TPA}^-\} \cup \{\mathfrak{B}\}),$$

where

$$\mathfrak{B} \stackrel{\text{def}}{=} \langle \{\emptyset, \{\langle a, a \rangle\}\}, \cup, \cap, \sim, 0, 1, \circ, ^{-1}, 1' \rangle,$$

that is,  $\mathfrak{B}$  is the QRA with singleton base. By Lemma 6.5 this means that

$$\text{QRA} \models e \quad \text{iff} \quad (\text{TPA}^- \models e \wedge \mathfrak{B} \models e).$$

Note that  $\mathfrak{B}$  is term-equivalent to the two-element Boolean algebra, so  $\mathfrak{B}$  is axiomatized by the finite set  $\{\text{BA-axioms}, 1 = 1', x \circ y = x \cap y, x^{-1} = x\}$ .

Now suppose that  $\{e_0, \dots, e_{n-1}\}$  is a basis for  $\text{TPA}^-$ . Let  $\theta \stackrel{\text{def}}{=} 0 \neq 1 \wedge 1 = 1'$  (note that  $\theta$  holds in a full RRA iff its base is a singleton). Let

$$E \stackrel{\text{def}}{=} \{-\theta \rightarrow e_i : i < n\}$$

$$E' \stackrel{\text{def}}{=} \{\theta \rightarrow e'_i : i < m\},$$

where  $\{e'_i : i < m\}$  is a basis for the equational theory of  $\mathfrak{B}$ . Then  $E \cup E'$  is a finite equational basis for QRA, contradicting [40, 8.4(vii)]. Indeed,  $\text{QRA} \models E \cup E'$ , since

$$\mathfrak{A} \in \text{full QRA} \text{ and } \mathfrak{A} \models \neg\theta \text{ implies } \mathfrak{A} \in \text{TPA}^-,$$

whence  $\mathfrak{A} \models e_i$  if  $i < n$ , and

$$\mathfrak{A} \in \text{full QRA} \text{ and } \mathfrak{A} \models \theta \text{ implies } \mathfrak{A} \cong \mathfrak{B},$$

thus  $\mathfrak{A} \models e'_i$  if  $i < m$ . On the other hand, suppose that  $\text{QRA} \models e$ . Then  $\text{TPA}^- \models e$  and  $\mathfrak{B} \models e$ , so  $\{e_i : i < n\} \vdash e$  and  $\{e'_i : i < m\} \vdash e$ , and thus  $E \vdash \neg\theta \rightarrow e$  and  $E' \vdash \theta \rightarrow e$ , giving  $E \cup E' \vdash e$ .  $\square$

## 7 Related work

For the interested reader, here are some more remarks concerning work related to our paper.



1) It turned out that complexity issues, and representability of certain classes of algebras depend on the choice of the underlying set theory. *Non-well-founded set theories* were investigated, from this point of view, in e.g. Kurucz-Németi [19], Sain-Németi [36]. One of the referees of this paper pointed out that it would be interesting to investigate whether our Thm.4.1 would change if we replaced ZFC with some non-well-founded set theory (since it is clear that the axiom of foundation has been used in our proof).

2) Both of our referees pointed out that *directed cylindric algebras*  $CA^\uparrow$  have been introduced for providing a cylindric algebraic analogue of quasi-projective relation algebras, see e.g. Németi-Simon [28] and Sági [30]. Therefore it would be a nice future project to find a cylindric algebraic version of what is going on in this paper, implemented via  $CA^\uparrow$ .

3) Connecting to this, Gábor Sági called our attention to the following. According to our Theorems 6.6 and 6.7, the equational theory of  $TPA^-$  is recursively enumerable, but not finitely based. This result is in interesting contrast with the following one. For finite  $n > 3$ , the equational theory of  $CA^\uparrow$  is finitely based, and in some non-well-founded set theories, each  $CA^\uparrow$  is representable as a set algebra (such that each operation is a concrete, set theoretical one, and they are invariant for the automorphisms of the set theoretical universe).

4) As we already mentioned in subsection 2.3 item 2), in [40], relation algebras have been used for formalizing set theory. Some improvements of the results of [40] can be found in Kurucz [18] and in the more recent paper [2] by Andr eka and N emeti. In both papers, weaker theories of relation algebras were established – to strengthen the original results.

5) In his dissertation [11] and in related papers (e.g. [10]), Mikl os Ferenczi presented a representation theory of cylindric-like algebras, based on relativized set algebras (instead of the “square” ones used traditionally). As a side effect of this, he proved a family of new results solving the finitization problem in new cases (truly finite axiomatizations for finite dimensional cases).

6) One of our referees called our attention to the fact that Tarek Sayed-Ahmed [38] approaches the finitization problem of first order logic FOL *with equality* in a new way. Namely, [38] combines the ideas of Mikl os Ferenczi on *representations by relativized set algebras* (cf. the previous item 5)), and the *semigroup approach* adopted by Ildik o Sain in her solution to the finitization problem for FOL *without equality* (cf. e.g. [35]). This way (similarly to the solution in [35]), [38] gives *truly finite axiomatizations even in infinite dimensional cases*.

7) We mention but do not discuss here that Bal azs Bir o [6] gave a noteworthy negative result concerning one particular way of trying to solve the finitization problem (permutation invariance assumed).

## References

- [1] H. Andréka. Complexity of the equations valid in algebras of relations, 1990. Thesis for DSc with Hungar. Acad. Sci., Budapest.
- [2] H. Andréka and I Németi. Reducing first-order logic to  $Df_3$ , free algebras. In *Cylindric-like algebras and algebraic logic (editors: H. Andréka, M. Ferenczi, and I. Németi)*, Bolyai Society Mathematical Studies (series editor: D. Miklós), pages 15–35. Springer Verlag, 2012.
- [3] H. Andréka, I. Németi, and I. Sain. Applying Algebraic Logic to Logic. In T. Rus M. Nivat, C. Rattray and G. Scollo, editors, *Algebraic Methodology and Software Technology (AMAST'93, Proceedings of the Third International Conference on Algebraic Methodology and Software Technology, The Netherlands, 21-25 June 1993)*, Workshops in Computing, pages 5–26. University of Twente, Springer-Verlag, London, 1994.
- [4] H. Andréka, I. Németi, and I. Sain. Algebraic Logic. In D. M. Gabbay and F. Guenther, editors, *Handbook of Philosophical Logic Vol. II, Second edition*, pages 133–247. Kluwer Academic Publishers, 2001. Preprint form available from the home page of the Rényi Institute.
- [5] H. Andréka, I. Németi, I. Sain, and Á. Kurucz. General Algebraic Logic including Algebraic Model Theory: An Overview. In D. M. Gabbay L. Csirmaz and M. de Rijke, editors, *Logic Colloquium'92 (Proc. Veszprém, Hungary, 1992)*, Studies in Logic, Language and Computation, pages 1–60. CSLI Publications, Stanford, 1995.
- [6] B. Biró. Non-finite-axiomatizability Results in Algebraic Logic. *Journal of Symbolic Logic*, 57(3):832–843, 1992.
- [7] W. J. Blok and D. L. Pigozzi. Algebraizable Logics. *Memoirs Amer. Math. Soc.*, 77, 396:vi+78, 1989.
- [8] W. Craig. Boolean notions extended to higher dimensions. In L. Henkin J. W. Addison and A. Tarski, editors, *The Theory of Models*, pages 55–69. North-Holland, Amsterdam, 1965.
- [9] W. Craig and R. L. Vaught. Finite Axiomatizability Using Additional Predicates. *Journal of Symbolic Logic*, 23:289, 1958.
- [10] M. Ferenczi. A new representation theory, representing cylindric like algebras by relativized set algebras. In *Cylindric-like algebras and algebraic logic (editors: H. Andréka, M. Ferenczi, and I. Németi)*, Bolyai Society Mathematical Studies (series editor: D. Miklós), pages 135–162. Springer Verlag, 2012.

- [11] M. Ferenczi. Representation theory based on relativised set algebras originating from logic, 2012. Dissertation for D. Sc. with Hungar. Acad. of Sciences, Budapest. In Hungarian, ix+125 pp.
- [12] A. Formisano and E. G. Omodeo. An equational re-engineering of set theories. In *FTP (LNCS Selection)*, pages 175–190, 1998.
- [13] L. Henkin. L. Henkin’s remark in the discussion following R. D. Maddux’s lecture at the “Asilomar Conference”, July 6–12, 1987.
- [14] L. Henkin and J. D. Monk. Cylindric Algebras and Related Structures. *Proc. Tarski Symp., Amer. Math. Soc.*, 25:105–121, 1974.
- [15] L. Henkin, J. D. Monk, and A. Tarski. *Cylindric Algebras Part II*. North-Holland, Amsterdam, 1985.
- [16] B. Jónsson. On Binary Relations. In G. Hutchinson, editor, *Proc. of NIH Conference on Universal Algebra and Lattice Theory*, pages 2–5. Laboratory of Computer Research and Technology, National Institute of Health. Bethesda, Maryland, 1982.
- [17] B. Jónsson. The theory of binary relations. In H. Andr eka, J. D. Monk, and I. N emeti, editors, *Algebraic Logic (Proc. Conf. Budapest 1988)*, pages 245–292. Colloq. Math. Soc. J. Bolyai, North-Holland, Amsterdam, 1991.
- [18]  . Kurucz. Weakly associative relation algebras with projections. *Mathematical Logic Quarterly*, 55:138–153, 2009.
- [19]  . Kurucz and I. N emeti. Representability of pairing relation algebras depends on your ontology. *Fundamenta Informaticae*, 44(4):397–420, 2000.
- [20] R. D. Maddux. Finitary algebraic logic, 1987. Invited lecture at the conference “Algebras, Lattices and Logic”, Asilomar California, July 6–12.
- [21] R. D. Maddux. Finitary Algebraic Logic. *Zeitschrift f. Math. Logic u. Grundl. Math.*, 35:321–332, 1989.
- [22] Sz. Mikul as, I. Sain, and A. Simon. The Complexity of the Equational Theory of Relational Algebras with Standard Projection Elements. Technical report, Math. Inst. Hungar. Acad. Sci., Budapest, 1991.
- [23] Sz. Mikul as, I. Sain, and A. Simon. Complexity of Equational Theory of Relational Algebras with Projection Elements. *Bulletin of the Section of Logic*, 21(3):103–111, 1992.
- [24] J. D. Monk. On an Algebra of Sets of Finite Sequences. *Journal of Symbolic Logic*, 35:19–28, 1970.

- [25] I. Németi. Free Algebras and Decidability in Algebraic Logic, 1986. Dissertation for D. Sc. with Hungar. Acad. of Sciences, Budapest. In Hungarian, xviii+169 pp.
- [26] I. Németi. On varieties of cylindric algebras with applications to logic. *Annals of Pure and Applied Logic*, 36:235–277, 1987.
- [27] I. Németi. Algebraization of Quantifier Logics, an Introductory Overview. *Studia Logica*, 50, No 3/4 (a special issue devoted to Algebraic Logic, eds.: W. J. Blok and D. L. Pigozzi):485–570, 1991. Strongly updated and expanded [e.g. with proofs] version is electronically available as: <http://circle.math-inst.hu/pub/algebraic-logic/survey.dvi>, [survey.ps](http://circle.math-inst.hu/pub/algebraic-logic/survey.ps).
- [28] I. Németi and A. Simon. Weakly higher order cylindric algebras and finite axiomatization of the representables. *Studia Logica*, 91:53–62, 2009.
- [29] H. Rogers. *Theory of recursive functions and effective computability*. McGraw Hill, 1967.
- [30] G. Sági. A completeness theorem for higher order logics. *Journal of Symbolic Logic*, 65(2):857–884, 2000.
- [31] I. Sain. On the search for a finitizable algebraization of first order logic (shortened version a). Technical report, Math. Inst. Hungar. Acad. Sci., 1988.
- [32] I. Sain. On finitizing first order logic. *Bulletin of the Section of Logic*, 23(2):66–79, 1994.
- [33] I. Sain. On the Problem of Finitizing First Order Logic and its Algebraic Counterpart (A Survey of Results and Methods). In D. M. Gabbay L. Csirmaz and M. de Rijke, editors, *Logic Colloquium'92* (Proc. Veszprém, Hungary, 1992), Studies in Logic, Language and Computation, pages 243–292. CSLI Publications, Stanford, 1995.
- [34] I. Sain. Definability issues in universal logic. In *Cylindric-like algebras and algebraic logic* (editors: H. Andréka, M. Ferenczi, and I. Németi), Bolyai Society Mathematical Studies (series editor: D. Miklós), pages 393–419. Springer Verlag, 2012.
- [35] I. Sain. On the Search for a Finitizable Algebraization of First Order Logic. *Logic Journal of the IGPL*, 8(4):495–589, July 2000. Electronically available as: [http://www3.oup.co.uk/igpl/Volume\\_08/Issue\\_04/](http://www3.oup.co.uk/igpl/Volume_08/Issue_04/).
- [36] I. Sain and I. Németi. Fork algebras in usual and in non-well-founded set theories. (extended abstract) part i and part ii. *Bulletin of the Section of Logic*, 24(3 and 4):158–168 and 182–192, 1995.

- [37] T. Sayed Ahmed. Algebraic logic, where does it stand today? *Bulletin of Symbolic Logic*, 11(4):465–516, 2005.
- [38] T. Sayed Ahmed. On notions of representability for cylindric polyadic algebras, and a solution to the finitizability problem for quantifier logics with equality. *Mathematical Logic Quarterly*, In press, 2015.
- [39] A. Simon. What the Finitization Problem is Not. In C. Rauszer, editor, *Algebraic Methods in Logic and in Computer Science* (Proceedings of the '91 Banach Semester, Banach Center), volume 28 of *Banach Center Publications*, pages 95–116. Institute of Math., Polish Acad. Sci., Warszawa, 1993.
- [40] A. Tarski and S. Givant. *A Formalization of Set Theory Without Variables*, volume 41 of *AMS Colloquium Publications*. Providence, Rhode Island, 1987.
- [41] M. van de Vel. Interpreting first-order theories into a logic of records. *Studia Logica*, 72:411–432, 2002.
- [42] P. A. S. Veloso and A. M. Haeberer. A Finitary Relational Algebra for Classical First-order Logic. *Bulletin of the Section of Logic*, 20(2):52–62, June 1991.
- [43] P. A. S. Veloso and A. M. Haeberer. A New Algebra of First-order Logic. In *Methodology and Philosophy of Science* (Proc. 9th International Congress on Logic, Upsala, Sweden), 1991.