

BIROn - Birkbeck Institutional Research Online

Ameh, E.M. and Tyrrel, S. and Harris, J. and Ignatiou, A. and Orlova, Elena and Nocker, A. (2018) The absence or presence of a lytic coliphage affects the response of Escherichia coli to heat, chlorine, or UV exposure. Folia Microbiologica 63 (5), pp. 599-606. ISSN 0015-5632.

Downloaded from: https://eprints.bbk.ac.uk/id/eprint/22292/

Usage Guidelines: Please refer to usage guidelines at https://eprints.bbk.ac.uk/policies.html or alternatively contact lib-eprints@bbk.ac.uk.

Manuscript Details

Manuscript number	IJHEH_2018_155
Title	Fate of heat-, chlorine- or UV- exposed Escherichia coli in absence and presence of a lytic coliphage
Article type	Full Length Article

Abstract

Disinfection aims at maximal inactivation of target organisms and the sustainable suppression of their regrowth. Whereas many disinfection efforts achieve efficient inactivation when the effect is measured directly after treatment, there are questions about the sustainability of this effect. One aspect is that the treated bacteria might recover and regain the ability to grow. In an environmental context another question is how amenable surviving bacteria are to predation by omnipresent bacteriophages. Provisional data suggested that bacteria when subjected to sublethal heat stress might develop a phage-resistant phenotype. The result made us wonder about the susceptibility to phage-mediated lysis for bacteria exposed to a gradient of chlorine and UV-LED disinfection strengths. Whereas bacteria exposed to low sublethal chlorine doses still underwent phage-mediated lysis, the critical chlorine Ct of 0.5 mg min L-1 eliminated this susceptibility and induced phage resistance in the cells that survived treatment. In the case of UV, even the smallest tested dose of 2.8 mJ cm-2 abolished phage lysis leading to direct regrowth. Results suggest that there is a possibility that bacteria surviving disinfection might have higher environmental survival chances directly after treatment compared to non-treated cells. A reason could potentially lie in their compromised metabolism that is essential for phage replication.

Keywords	disinfection; heat; chlorine; UV-LED; bacteriophages; phage resistance
Corresponding Author	Andreas Nocker
Corresponding Author's Institution	IWW Water Center
Order of Authors	Ekwu Ameh, Sean Tyrrel, Jim Harris, Athanasios Ignatiou, Elena Orlova, Andreas Nocker
Suggested reviewers	Hans-Curt Flemming, Mark van Loosdrecht, Banu Ormeci, Michael Templeton

Submission Files Included in this PDF

File Name [File Type]

Cover_letter_IJHEH.docx [Cover Letter]

Graphical_abstract.pptx [Abstract]

20180225_Ameh_et_al.docx [Manuscript File]

Figure_1.pptx [Figure]

Figure_2.pptx [Figure]

Figure_3.pptx [Figure]

Figure_4.pptx [Figure]

Figure_supplementary_1S.pptx [Figure]

Figure_supplementary_2S.pptx [Figure]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given: Examples of selected raw data has been added as supplementary figures. Processed data is included in the manuscript. February 25, 2018

Editor, International Journal of Hygiene and Environmental Health

Dear Editor,

we would like to submit a manuscript entitled 'Fate of heat-, chlorine- or UV- exposed *Escherichia coli* in absence and presence of a coliphage' for consideration of publication.

The study assessed the sustainability of the effect of heat, chlorine and UV stress on the culturability of *Escherichia coli* in the absence and presence of a lytic coliphage. *E. coli* was exposed to stress gradients of increasing intensity followed by monitoring the effect on colony formation over four consecutive days. Prior to plating bacterial suspensions, a lytic coliphage was added to one set of samples. Results demonstrated (1) differences in the sustainability of the different disinfection methods and, most importantly, (2) that there is likelihood that sublethally stressed bacteria are not susceptible to lysis by certain lytic phages any more. Instead they seem to acquire a phage-resistant phenotype. We assume that the metabolic repercussions of the stress interfere with the production of phage particles. As a consequence, bacteria surviving disinfection can have higher chances of survival in the environment (with bacteriophages being responsible to a large part of bacterial mortality apart from protozoan grazers). Data is in line with other studies reporting that phages 'sense' the viability of the host cells.

We believe that this manuscript represents a comprehensive data set with relevance for different fields of microbiology where disinfection and biocidal treatments are applied. It might contribute knowledge to understand the sometimes unsatisfying outcome of disinfection procedures.

All authors approve of its content and conclusions. If I may answer any questions, please contact me.

Respectfully, Andreas Nocker

Dr. Andreas Nocker IWW Water Centre Applied Microbiology Moritzstrasse 26 Muehlheim an der Ruhr Germany

E: andreas.nocker@gmail.com T: +49-151 15999115

Graphical abstract

Bacteria without disinfection



Amenable to lytic phages

Surviving bacteria after disinfection





No lysis, but direct regrowth

Log change in bacterial CFU after disinfection

Bacteria surviving disinfection can show less susceptibility to lytic bacteriophages leading to direct regrowth compared to bacteria that were not subjected to disinfection.

1	Fate of heat-, chlorine- or UV- exposed <i>Escherichia coli</i> in absence and
2	presence of a lytic coliphage
3	
4	Ekwu M. Ameh ^a , Sean Tyrrel ^a , Jim Harris ^b , Athanasios Ignatiou ^c , Elena Orlova ^c ,
5	Andreas Nocker ^{a,d,*}
6	
7	¹ Cranfield Water Science Institute, School of Water, Environment and Energy, Cranfield
8	University, Cranfield, Bedfordshire, MK43 0AL, United Kingdom
9	² Cranfield Institute for Resilient Futures, Cranfield University, Cranfield, Bedfordshire,
10	MK43 0AL, United Kingdom
11	³ Institute for Structural and Molecular Biology, Department of Biological Sciences,
12	Birkbeck University of London, Malet Street, London WC1E 7HX, United Kingdom
13	⁴ IWW Water Centre; Moritzstraße 26, 45476 Mülheim an der Ruhr, Germany
14	
15	
16	
17	
18	
19	* Corresponding author: IWW Water Centre; Moritzstraße 26, 45476 Mülheim an der Ruhr,
20	Germany. Tel: +49-208 40303 383, Email: andreas.nocker@gmail.com

60		
61 62		
62 63	21	Highlights
64 65	22	
66 67	23	• Study assessed sustainability of disinfection on <i>E. coli</i> in presence of a phage.
68 69	24	• Bacteria exposed to heat, chlorine and UV prior to phage addition.
70 71 70	25	• Stress exposure can interfere with ability of phage to lyse the bacteria.
72 73 74	26	• There is a possibility that stressed bacteria acquire a phage-resistant phenotype.
75 76	27	• Phages can 'sense' the viability of bacteria.
77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107	28	Stress might increase survival chances in an environment where phages are present.
108 109 110		
110 111 112 113 114		
115 116 117		
118		

29 ABSTRACT

Disinfection aims at maximal inactivation of target organisms and the sustainable suppression of their regrowth. Whereas many disinfection efforts achieve efficient inactivation when the effect is measured directly after treatment, there are questions about the sustainability of this effect. One aspect is that the treated bacteria might recover and regain the ability to grow. In an environmental context another question is how amenable surviving bacteria are to predation by omnipresent bacteriophages. Provisional data suggested that bacteria when subjected to sublethal heat stress might develop a phage-resistant phenotype. The result made us wonder about the susceptibility to phage-mediated lysis for bacteria exposed to a gradient of chlorine and UV-LED disinfection strengths. Whereas bacteria exposed to low sublethal chlorine doses still underwent phage-mediated lysis, the critical chlorine Ct of 0.5 mg min L⁻¹ eliminated this susceptibility and induced phage resistance in the cells that survived treatment. In the case of UV, even the smallest tested dose of 2.8 mJ cm⁻² abolished phage lysis leading to direct regrowth. Results suggest that there is a possibility that bacteria surviving disinfection might have higher environmental survival chances directly after treatment compared to non-treated cells. A reason could potentially lie in their compromised metabolism that is essential for phage replication.

47 Keywords: disinfection; heat; chlorine; UV-LED; bacteriophages; phage resistance

- **48**

1. Introduction

Whereas heat is undoubtedly one of human's best weapons for food preservation, treatment with chlorine and ultraviolet light in the UV-C range represent two of the most common disinfection procedures for water treatment (Hijnen et al., 2006; Munakata and Kuo, 2016). Although the aim of disinfection is to maximally reduce counts of live target organisms, there are (in contrast to sterilization that utilizes harsher killing conditions) always surviving microbes (in a biofilm context referred to as 'persisters'). Factors known to reduce the efficiency of chlorination include the presence of chlorine demand (Haas and Karra, 1984), the performance of disinfection at suboptimal pH (White, 1999) or cold temperatures (Le Dantec et al, 2002). Factors reducing the efficiency of UV-C disinfection, on the other hand, include insufficient transmissivity of the treated water (Shin et al. 2001), the presence of substances with UV absorbing properties (Templeton et al., 2006) or shielding effects by particulates (Mamane-Gravetz and Linden, 2004).

Microbiological assessment of the success of disinfection should focus both at measuring (1) the immediate efficiency directly after treatment and (2) the sustainability of the effect. The first can be determined relatively easily and depends on the chosen treatment conditions, the nature of the treated matrix and the target organism. Much more uncertainty is associated with the latter due to the fact that it needs more than a single diagnostic snapshot at a given time point to provide an answer. The sustainability of disinfection depends both on the innate capacity of the specific organism to repair the induced damage and the impact of disinfection on the organism's ability to survive in the surrounding environment. For bacteria, a critical biological factor consists in their susceptibility to bacteriophages that represent (together with protozoan grazers) the most important group of bacterial predators in real-world conditions (Clokie et al., 2011). Numbers of phages in the environment are estimated to exceed the ones of bacteria by up to 10-fold explaining a strong dynamics of bacterial

populations (Labrie et al., 2010; Brüssow and Hendrix, 2002). Bacteria are however not equally susceptible to phage attack in every stage of their life cycle (McGrath and van Sinderen, 2007) and phage replication depends on bacterial vitality (Said et al., 2010). In initial experiments we observed that bacteria exposed to sublethal heat stress were less susceptible to phage-mediated lysis. This led to the question whether a phage-resistant phenotype can also originate from other stresses. Chlorine and UV treatment were chosen as they are commonly applied water disinfection procedures and rely on different inactivation mechanisms.

For all three stresses, bacteria were subjected to stress gradients ranging from very mild to lethal. The research question was how amenable bacteria, which were subjected to sublethal disinfection stress, are to phage-mediated lysis (in comparison with controls without phage). Whereas the experiments with heat were produced by monitoring the optical densities of bacterial suspensions over time, experiments with chlorine and UV were performed in more depth with membrane filtration of samples at different time points after disinfection and subsequent phage challenge. E. coli and a lytic coliphage were chosen as models. The coliphage was isolated from an environmental brook and induced efficient lysis of its bacterial host within approx. 6 hours.

2. Materials and methods

2.1. Bacterial growth conditions

Escherichia coli ATCC 25922 (a reference strain for antimicrobial susceptibility testing) was streaked from glycerol stock onto tryptone soya agar (TSA; CM0131; Oxoid Ltd, Basingstoke, Hampshire, UK) and grown for 24 h at 30°C. Single colonies were subsequently transferred into 10 mL of dilute (strength: 10 %) tryptone soya broth (TSB; CM1016; Oxoid Ltd, Basingstoke, Hampshire, UK) contained in 50 mL Falcon tubes followed by overnight incubation at 20 °C (for heat experiments) or 25°C (for chlorine and UV experiments) at 250 rpm in a temperature-controlled mini shaker (cat. number 980151, VWR, USA). Tubes were shaken in a 45° angle. Cell density was measured in a spectrophotometer (JENWAY 6310, Camlab, England, UK) at 600 nm (OD₆₀₀) and adjusted to an OD₆₀₀ = 1.0 by dilution with 10% TSB (equilibrated to room temperature). Aliquots of 1 mL of the density-adjusted bacterial culture were then transferred aseptically into sterile 1.5 mL microcentrifuge tubes and spun at 5,000 x g for 5 minutes. The supernatants were removed carefully using a tip and the cell pellets were resuspended in equal volumes of 0.1 µm filtered mineral water (Evian, Evian-les-basin, France). This washing step was repeated two times to remove organic traces from the culture broth. Bacteria were finally resuspended in phosphate buffered saline (PBS, pH7) for the heat experiment or in filtered mineral water for the chlorine and UV experiment. Whereas the *E. coli* suspension was used undiluted (approximately 10⁹ CFU mL⁻¹) for the heat experiment, cells were diluted to a final concentration of 10⁵ CFU mL⁻¹ for chlorine and UV disinfection. 2.2. Isolation of a lytic phage

A water sample was collected from Chicheley brook at a location receiving treated effluent discharge from the Cranfield University wastewater treatment plant (Cranfield, Bedforshire, UK). The sample was passed through a 0.22 µm membrane filter (Millex GP, Merck Millipore, Darmstadt, Germany). 10 mL of the filtered flow through was mixed in a 50 mL Falcon tube with an equal volume of double strength TSB and supplemented with 5 mM CaCl₂. The mixture was inoculated with 0.4 mL of an overnight *E. coli* culture (grown at 20 °C) followed by incubation (30°C; 250 rpm) until clearance indicated bacteriophages-mediated cell lysis. Chloroform was added to a final concentration of 2% to eliminate remaining intact bacteria followed by centrifugation at 5,000 x g for 5 min and passage of the supernatant through a 0.22 µm membrane filter (Millex GP, Merck Millipore, Darmstadt, Germany). The phage suspension was serially diluted in sodium magnesium buffer (SM buffer; 100 mM NaCl, 8 mM MgS04*7H20, 50 mM Tris-Cl, adjusted to pH 7.5). Aliquots of 100 µl were mixed with 150 µL of log-phase E. coli followed by incubation for 10 minutes to allow adsorption. The phage-bacterial mixture was added to 3 mL of molten 0.7% TSB agar (maintained at 48°C), mixed immediately by gentle vortexing and then distributed evenly over TSA agar plates supplemented with CaCl₂ to a final concentration of 5 mM. The soft agar was allowed to solidify for 20-30 min and plates were incubated overnight at 30°C (or indicated temperature) to allow for plaque formation. The plaque with the largest diameter was transferred into SM buffer and re-suspended. To ensure purity, the soft agar overlay method to obtain new plaques was repeated three times. To obtain a phage stock, isolated plaques were picked using sterile wooden toothpicks to inoculate 5 mL log-phase E. coli cultures followed by incubation at 30°C with shaking (250 rpm) for 8 hours. Lysate from single plaques were treated with chloroform to a final concentration of 2%, mixed and centrifuged at 5,000 x g for 5 min. The phages were recovered from the upper phase suspension and passed through a 0.22 µm filter (Millex GP, Merck Millipore, Darmstadt, Germany). Phage titers tended to be $> 10^9$ PFU ml⁻¹ and stocks were stored at 4°C.

2.3. Transmission electron microscopy

For phage preparation, 3 µl of phage suspension (approx. 10° PFU ml⁻¹) were pipetted on to a negatively glow discharged 10 µm thick C-FlatTM carbon grid (400-mesh) with no dilution and allowed to sit for 1 minute. The sample droplet was then partially blotted on Whatman[™] quantitative filter paper, Grade 1. Three μ l of 2 % uranyl acetate stain solution was immediately applied. After 1 min of staining, all excess fluid was removed from the grid surface by positioning the filter in an angle and the sample was allowed to air-dry (approx. 3-5 min). The sample on each grid was then imaged in low dose conditions on a Tecnai 10 transmission electron microscope (FEI company, Oregon, USA) operating at 100 keV. Images were taken at 20 K magnification and captured using a Gatan Ultrascan 4000 $4k \times 4k$ CCD camera with an ultra-sensitive phosphor scintillator (Gatan, USA) to produce a final pixel sampling of 11 Angstroms per pixel.

2.4. Heat experiments and monitoring of optical densities

Heat stress was performed by exposing ten aliquots (1 ml each) of E. coli suspensions $(OD_{600} = 1.0, \text{ in PBS})$ to different temperatures (4°C, 20°C, 37°C, 41°C, 44°C or 50°C) for 24 hours. Cells were subsequently harvested by centrifugation (5,000 g for 5 min) and re-suspended in 1 ml TSB supplemented with 5 mM CaCl₂. The ten aliquots exposed to the same temperature were pooled in 50 ml conical centrifuge tubes (Fisherbrand, Fisher Scientific UK Ltd., Loughborough) to obtain 10 ml suspensions for each temperature. Temperature-exposed bacteria were challenged with phage at a multiplicity of infection (MOI) of 1 by adding 10¹⁰ phage particles from the high titer phage stock followed by shaking at 250 rpm at 20°C. Optical densities (600 nm) of 1 ml aliquots of cell suspensions were measured on a TECAN M200 Pro plate reader (Tecan UK Ltd, Reading, UK) in

transparent 48 well tissue culture plates (non-treated, flat bottom, cat. nr. TCP001048; Jet Biofil, Braine I'Alleud, Belgium). Readings were taken directly after addition of phage and after 6, 12, and 24 hours. The experiment was repeated three times.

2.5. Disinfection by chlorination

A chlorine solution was prepared by diluting 95 μ L of sodium hypochlorite stock solution (10-15% available chlorine, Sigma-Aldrich, USA) in 50 mL of ultrapure water of 100 mL volume flask to a concentration of approximately 200 mg L⁻¹ free chlorine. Volumes of 0.5, 1.25, 5, 12.5 and 25 µL of this chlorine solution were added to 50 mL aliquots of bacterial suspensions (10⁵ CFU/mL in 0.22 µm filtered Evian mineral) and stirred at 150 rpm in chlorine demand-free beakers to obtain the following final free chlorine concentrations: 0.002, 0.005, 0.02, 0.05 and 0.1 mg L⁻¹. Cells were exposed to chlorine at room temperature (20 °C) for 10 min translating to a Ct value range between 0.02 and 1 mg min L^{-1} (free chlorine concentration x time). Chlorine disinfection was stopped by addition of 250 µL of 0.1N sodium thiosulphate (Na₂S₂O₃*5H₂O) (Acros Organics, Geel, Belgium). A bacterial suspension without added chlorine served as a control. After letting the samples stand for 10 min, TSB and CaCl₂ were added to final concentrations of 10% TSB and 5 mM CaCl₂, respectively, to provide nutrients for the bacteria and to enable phage adsorption. To one series of samples, a small volume of concentrated phage stock was added to obtain a phage concentration of 10⁵ PFU/mL (corresponding to a multiplicity of infection (MOI) of 1. For time point t = 0, samples were membrane-filtered directly after phage addition, no time for phage adsorption was provided. An identical series of bacterial suspensions was left without phage addition with one aliquot also being membrane-filtered at t=0 for colony enumeration. Suspensions (with and without phage) were shaken on an orbital shaker (250 rpm) at 20°C for 4 days in total and colonies enumerated by membrane filtration after 6, 12, 24, 48, 72 and

96 hours. All experiments were performed with chlorine demand-free glassware prepared according to Charnock and Kjønnø (2000) and carried out as three independent repeats.

2.6. Disinfection by LED-UV

UV disinfection was performed at room temperature (20 °C) using a UV-C-LED system consisting of a UVCLEAN lamp with multi-chip collections of UV LEDs from Sensor Electronic Technology (Columbia, South Carolina, USA). LEDs emitted UV light at 260 nm with a power output of 15 mW. Samples were placed at a distance of 1 cm to the UV source. The delivered dose was verified using a UV-C radiometer (VLX-3W from Vilber Lourmat) and determined to be 20 W/m². Bacterial suspensions (10⁵ CFU mL⁻¹ in filtered Evian water) were transferred in 20 mL aliquots into a 25 mL Pyrex petri dishes (Fisher Scientific, UK). During UV exposure suspensions were stirred constantly using a magnetic stir bar on a stirrer set to 400 rpm. Bacteria were exposed to the following UV doses: 2.8, 4.8, 10.5, 15.4, 20.3, and 39.9 mJ/cm². A bacterial suspension that was treated identically, but not exposed to UV served as a control. Following UV exposure, suspensions were supplemented with TSB and CaCl₂ to final concentrations of 10% and 5 mM CaCl₂, respectively. To one series of samples phages were added to achieve an MOI of 1. For time point t = 0 samples were membrane-filtered directly after phage addition, no time for phage adsorption was provided. An identical series of bacterial suspensions was left without phage addition with one aliquot also being membrane-filtered at t=0 for colony enumeration. Suspensions (with and without phage) were shaken at 250 rpm at 20°C for 4 days in a laboratory incubator with a transparent lid receiving only normal laboratory light during the day. Colonies were enumerated by membrane filtration after 6, 12, 24, 48, 72 and 96 hours. Experiments were carried out in three independent repeats with standard deviations being slightly greater than for the chlorine experiment.

591			
592			
594	222		
595			
596	223	2.7. Bacterial quantification by plate counting and flow cytometry	
597	224		
598	224		
599	225	For enumeration of <i>E. coli</i> , the membrane filtration method was used. Filtered mineral	
600 601			
602	226	water (~ 3 mL) was poured into the filter manifold (Combisart, Sartorius, UK) prior to	
603			
604	227	addition of 100 μ L of sample (10^5 CFU/mL) to ensure homogeneous distribution of bacter	ria
605			
606	228	on the filter. Samples were filtered onto 0.45 μ m cellulose filters (47 mm, white gridded, C	at.
608			
609	229	No. HAWG04/86 Fisher Scientific, UK) and placed on 55 mm Petri dishes with membrane	9
610	220	lactore alucuronide agar (MIGA: Oxoid Fisher Scienitic LIK) MIGA gives rise to green	F
611	230	lactose gluculonide agai (MLOA. Oxold, Fisher Scientife, OK). MLOA gives lise to green	Ľ.
612	231	<i>coli</i> colonies Plates were incubated at 35°C for 24 hours before enumeration Flow	
613			
614 615	232	cytometry was performed as described previously (Nocker et al., 2017). Bacterial were	
616			
617	233	stained with SYBR Green I and propidium iodide with final concentrations of 1 x and 3 μ M	1.
618			
619			
620			
621			
622			
623			
024 625			
626			
627			
628			
629			
630			
631			
032 633			
634			
635			
636			
637			
638			
639			
040 641			
642			
643			
644			
645			
646			
047 678			11
649			
-			

3. Results

E. coli was subjected to gradients of heat, chlorine or UV of increasing intensities with aliquots not undergoing treatment serving as controls. Bacteria were subsequently challenged with a lytic coliphage or not. A phage was isolated for this purpose from a brook receiving treated wastewater discharge. The phage was identified by transmission electron microscopy to belong to the group of Myoviridae, which have double-stranded DNA and contractile tails (Fig. 1A). The kinetics of the phage to lyse *E. coli* at a multiplicity of infection (MOI) of 1.0 was measured by flow cytometry. Time to maximal lysis at 20 °C was around 6 hours under given conditions (Fig. 1B).

3.1. Fate of E. coli after heat stress

In an initial experiment we examined the effect of temperature on fully grown bacteria in regard to their susceptibility to phage-mediated lysis. E. coli was grown at 20°C and subsequently suspended in physiological salt to prevent cell replication. Bacterial suspensions were exposed to either 4 °C, 20 °C, 37 °C, 41 °C, 44 °C or 50 °C for 24 h prior to phage challenge at 20°C (in the presence of nutrients). Bacteria exposed to temperatures $\leq 37^{\circ}$ C were all found to be susceptible to phage lysis reflected by a decrease in OD_{600} (Fig. 2A). Bacteria exposed to heat (41 and 44 $^{\circ}$ C) on the other hand were not lysed and the OD₆₀₀ slightly increased suggesting the presence of viable bacteria (especially when exposed to 41°C). Exposure to 50°C, on the other hand, can be considered lethal as no change in optical density was observed and no colonies were obtained after this treatment.

To unequivocally assign the temperature effect to phage susceptibility, bacterial aliquots with no added phage served as controls (Fig. 2B). Optical densities of bacteria exposed to 4, 20 or 37°C increased rapidly suggesting healthy bacteria. Also optical densities

of bacteria exposed to 41 and 44°C showed an increase (although slow and moderate)
indicating that these aliquots contained viable cells and that bacteria have the ability to
survive exposure to these temperatures. The slow increase was probably caused either by a
longer lag time due to stress exposure or by the reduced number of viable cells in the
bacterial population. Only aliquots exposed to 50°C showed a straight line suggesting that
this temperature was lethal.

The described trends are reflected in a slope analysis indicating the rates of change in optical densities (Fig. 2A and B). In case of bacterial suspensions exposed to 4, 20, 30°C prior phage challenge, previously positive slopes (as seen in the control without phage) are turned negative by phage addition resulting in lysis and consequently a drop in optical densities. For E. coli suspensions treated at 41 and 44°C, on the other hand, slopes were slightly positive in all samples independent of whether a phage was added or not. Obviously exposure to these temperatures prevented overall lysis and a dip in OD_{600} . Although interesting in itself, the question which remained unsolved from the experiment was the distribution of viability states within the bacterial population. Only a small portion of the E. *coli* population might still have remained viable and been amenable to lysis. As measurement of optical density has limited sensitivity and does not reflect changes to a small proportion of bacteria, colony counting was applied for subsequent experiments.

0 278

279 3.2. Fate of *E. coli* after chlorine disinfection

754 280

Bacteria were exposed to increasing free chlorine doses up to 1 mg min L⁻¹ (followed by addition of thiosulphate to neutralize the disinfectant). The increasing damage to the cell membrane inflicted by increasing chlorine doses was visualized by flow cytometry in combination with membrane integrity staining. Bacterial membrane damage started to be visible in dot blots in samples exposed to 0.5 mg min L⁻¹ and bacteria in samples exposed to

1 mg min L⁻¹ were readily stained with propidium iodide indicating substantial membrane damage (data not shown). After chlorine exposure bacteria were either mixed with the lytic bacteriophage or not. The ability of the cells to form colonies was assessed over 4 days at time points 0, 6, 12, 24, 48, 72 and 96 hours. Using selective MLGA agar, E. coli appear as green colonies or, in case of many bacteria, as a green lawn (supplementary Fig. 1S). Directly after disinfection (time point zero), strong growth was detected for bacteria exposed to up to 0.2 mg min L⁻¹ free chlorine meaning that these doses had little effect on culturability. Exposure to 0.5 mg min L⁻¹ chlorine on the other hand led to a substantial decrease in colonies on the filters (although bacterial membrane damage was only moderate as assessed by flow cytometry). Higher chlorine doses completely suppressed growth. This effect at time point zero was independent of whether phage was added or not. At later time points differences between the two sample series became visible.

The change in colony numbers within the first 6 hours (representing the time to maximal lysis) is graphically shown in Fig. 3. In the absence of phage, the number of colonies on filters increased over time for all samples where growth was seen directly after disinfection. This is reflected in a positive log change in colony numbers within the first 6 hours. For bacteria exposed to the highest chlorine dose (1 mg min L⁻¹), the chlorine effect was sustained with no colonies detected even after 96 hours. In case, a phage was added directly after disinfection, the outcome was very different. Bacterial densities of samples exposed to chlorine doses up to 0.2 mg min L⁻¹ were visibly reduced after 6 hours with the log changes in CFU being negative for those samples (Fig. 3). The reduction in bacterial colonies was strongest for aliquots with no chlorine, however lytic activity was observed up to chlorine doses of 0.2 mg min L^{-1} . For bacterial suspensions exposed to 0.5 mg min L^{-1} chlorine, on the other hand, colony numbers did not decrease, but rather an increase in colonies was observed. Bacterial numbers on these filters were comparable to the ones in the corresponding sample where no phage had been added (Fig. 1S) and the log change in CFU

within the first 6 hours was identical for the two experimental series (independent of whether a phage was added or not; Fig. 3). The chlorine dose of to 0.5 mg min L⁻¹ therefore seemed to erase the difference between samples with and without phage, probably because such treated bacteria were not lysed by the phage. In other words, the phage appeared unable to lyse E. *coli* that were exposed to a chlorine dose of 0.5 mg min L⁻¹. No phage effect could be deducted for the highest chlorine dose (1 mg min L⁻¹) as this treatment sustainably eliminated all E. coli growth, independent of the presence of phage.

3.3. Fate of E. coli after UV disinfection

E. coli suspensions were exposed to different UV-C doses up to approx. 40 mJ cm⁻¹ (which is the minimal dose that has be applied for disinfection of drinking water in a number of countries; DVGW, 2006). In contrast to chlorine, the killing by UV left bacterial membranes intact as assessed by flow cytometry (data not shown)). As for the chlorine experiment, bacteria were subsequently either mixed with the lytic bacteriophage or not. Bacterial densities on filters at time point zero (directly after disinfection) progressively decreased with increasing UV doses (both for samples where phage was absent or present; supplementary Fig. 2S). No colonies were observed in the sample experiencing the highest radiation. Like for chlorine disinfection, monitoring was performed over 4 consecutive days with samples being membrane-filtered and cultured after 0, 6, 12, 24, 48, 72 and 96 hours. Also in line with chlorination, the change in colony numbers within the first 6 hours was numerically assessed (Fig. 4).

In the absence of a phage, colony numbers on filters increased over time for all samples. This is reflected in a positive log change in colony numbers within the first 6 hours (Fig. 4). The higher the applied UV dose, the longer regrowth was delayed translating into decreasing log changes in CFU within the first 6 hours. After 12 hours, regrowth was also

887		
888 889	338	observed for the sample exposed to the highest dose and that did not show any growth
890		
891	339	directly after UV disinfection (supplementary Fig. 28). The reappearance of colonies might
892 893 804	340	be due to repair of UV damage over time. Another possibility is the replication of bacteria
895 896	341	that survived UV treatment (e.g. due to shading effects during stirring).
897 898	342	The same outcome applied to all samples where a phage had been added after UV
899 900	343	disinfection with the non-irradiated control being the only exception. Whereas all bacterial
901 902	344	suspensions undergoing UV treatment showed increasing colony numbers and positive log
903 904	345	changes in CFU within the first 6 hours (as in the samples without phage), only the non-
905 906 907	346	irradiated control showed a decrease in bacterial numbers in the first hours after phage
908 909	347	addition by > 3 log units (Fig. 4). It seemed that only those bacteria that were not exposed to
910 911	348	UV were lysed by the phage, whereas even low UV doses efficiently prevented lysis. The
912 913	349	fact that the dose of 2.8 mJ cm ⁻² extinguished the phage effect suggests that UV exposure
914 915	350	inflicted changes to the bacteria that prevented phage-mediated lysis, but did not suppress
916 917	351	growth.
918 919	352	
920 921		
922		
923		
924		
925		
920 927		
928		
929		
930		
931		
932		
933		
935		
936		
937		
938		
939		
940		
941 042		
943		1
944		

353 4. Discussion

We investigated in this study the sustainability of the effect of different disinfection intensities on the culturability of E. coli in the absence and presence of a lytic bacteriophage that exerted maximal lytic activity within several hours. Apart from the universally relevant thermal disinfection, chlorine and UV were chosen not only because they are frequently applied in water treatment, but also because they differ in their mechanisms of action. Whereas wet heat is assumed to kill primarily through inducing enzyme inactivation and protein denaturation (Mackey et al., 1991), chlorine as an unspecific oxidant inflicts damage to any cellular component it reacts with (including the cell envelope, nucleic acids and enzymes (Camper and McFeters, 1979; du Preez et al., 1995). UV on the other hand inactivates microbes by damaging their nucleic acids (Cutler and Zimmerman, 2011). Independent of phage addition, results of the disinfection experiments demonstrated differences in regard to the sustainabilities of the different treatments when comparing conditions that abolished growth directly after disinfection. No regrowth was measured in aliquots subjected to 50 °C or a chlorine dose of 1 mg min L⁻¹, whereas a UV dose of 40 mJ cm⁻² (the minimal UV dose recommended for disinfecting drinking water in a number of countries) did not prevent re-appearance of colonies after 12 hours. Full regrowth was obtained after 72-96 hours. It should be noted in this context that all samples were supplemented with nutrient broth after disinfection meaning that conditions allowed for rapid growth. This observation is in line with a study from Kollu and Örmeci (2014) reporting regrowth of E. coli after UV irradiation at 40 mJ cm⁻². No conclusion was however possible on the origin of the regrowing E. coli. It is not clear to date, whether the majority of regrowth comes from disinfection survivors (e.g. through shading events) or from recovery after repair. Both scenarios have support in the literature (Bohrerova et al., 2014; Hu et al., 2005). It

should be noted however that a relatively large volume (1 ml, representing 5% of the total

volume) was filtered directly after UV disinfection. As no single colony appeared directly after UV disinfection, repair appears to be more likely.

The main focus of this study was however the investigation of the susceptibility of stressed bacteria to lytic phages. A typical reason for the loss in susceptibility to bacteriophage attack is the destruction of surface antigens on the bacterial envelope and the resulting inability of phages to adsorb (Abedon, 2012). This possibility could especially apply to bacteria exposed to heat or chlorine with their effects on surface proteins through denaturation or oxidative damage. In the case of UV exposure, the destruction of surface antigens appears less likely as the integrity of the membrane was not compromised. A more probable hypothesis might be that the biosynthetic and metabolic capabilities of the UV-irradiated bacteria are affected to an extent where phage replication is no longer supported within the host cell. The overall viability and vitality of the bacterial host can be seen as an important determinant for the success of phage propagation. For E. coli 0157:H7, Awais et al. (2006) reported an elegant recombinant phage-based assay that allowed visualization of infected bacteria that supported phage proliferation. The phage was devoid of its lytic capability, but induced the expression of green fluorescence within host cells. It was shown with this assay that only culturable cells allowed phage replication, whereas viable but non-culturable (VBNC; produced by starvation at 4 °C) and dead cells merely allowed phage adsorption, but not biosynthesis. The observation was employed later in an assay to detect VBNC cells by Said et al. (2010). UV-exposed E. coli that had lost culturability were reported to allow adsorption, but did not support lysis. Further support came from subsequent results by Fernandes et al. (2014) for Salmonella enteriditis. When applying a phage-based system to assess the viability of the bacteria, VBNC cells (interestingly produced by treatment with a low chlorine dose) were reported to be recognized by the phage, but no lysis was induced. A different underlying mechanism was suggested for heat-treated cells that

404 were recognized by the phage only to a minor extent, possibly as phage receptors on the cell
 405 envelope had been destroyed by the heat.

In broad consent with these authors, our study corroborates that lytic phages can sense the viability or vitality of their bacterial hosts, which consequently determines the bacteria's fate. It is well known that viral proliferation requires a 'significant commitment of host resources to reproduce viral particles' (Maynard et al., 2011). The lytic success is thus dependent on the existence of a functional and energized metabolic machinery capable of production of nucleic acids, proteins and lipids that represent the building blocks of phage particles. A study using a marine bacterium model estimated that ~75% of carbon and nitrogen resources were redirected into the production of virions (Ankrah et al., 2014). Other researchers reported a strong depletion of bacterial transcripts upon phage infection (Chevallereau et al., 2016) and a strong perturbation to the interconnected network of metabolic pathways within the bacterial host (Birch et al., 2012). It is well conceivable that the metabolism of injured bacteria with compromised cellular functions might not be able to support this metabolic burden and therefore not undergo lysis. This does not answer on the other hand, how these bacteria would recover over time and why they would not be amenable at a later stage to phage attack. Instead of working with bacterial populations with undefined distribution of bacteria in different viability states, future experiments would profit from single cell techniques. It also will need to be shown whether the observations hold true for other lytic phages. Despite many open questions, results presented here suggest that sublethally stressed bacteria that survive mild disinfection conditions might show less susceptibility to lytic phages. For a limited time window, such bacteria might in consequence have an increased survival chance in the environment where phages are one of the major causes of bacterial mortality.

1123		
1124 1125	428	In conclusion, the role of bacteriophages in influencing the bacteria's fate after
1126 1127	429	disinfection is typically an overlooked parameter. We report here in a laboratory-based study
1128 1129	430	the possibility that stresses with different modes of action can modulate the bacteria's
1130 1131	431	susceptibility to phage-mediated lysis. Although more bacteria-phage interactions need to be
1132 1133	432	studies for generalization, sublethal disinfection could potentially rescue bacteria from phage-
1134 1135	433	mediated lysis. The stress intensity threshold where such resistance is observed appears
1136 1137	434	different between different stresses. We hypothesize that injury by mild disinfection might
1138 1139	435	compromise the bacterial metabolic and biosynthetic machinery to an extent that it does not
1140 1141	436	meet the high demand for phage replication
1142 1143	100	neet the high demand for phage representation.
1144		
1145		
1147		
1148		
1149		
1150		
1151		
1152		
1153		
1155		
1156		
1157		
1158		
1159		
1160		
1161		
1162		
1164		
1165		
1166		
1167		
1168		
1169		
1170		
1172		
1173		
1174		
1175		
1176		
1177		
11/8		00
1180		20

1181			
1182			
1183	437	Acknowledgements	
1184	107	Territo Wieugements	
1185			
1186	438	The contents presented here are based on the PhD thesis by the first author Ekwu Mark Am	ieh
1187			
1100	439	(Ameh EM, 2016). The research was supported by the Commonwealth Scholarship scheme	;
1109			
1191	440	of the British government (grant nr. NGCA-2012-326).	
1192			
1193			
1194			
1195			
1196			
1197			
1198			
1199			
1200			
1201			
1202			
1203			
1205			
1206			
1207			
1208			
1209			
1210			
1211			
1212			
1213			
1214			
1215			
1210			
1217			
1219			
1220			
1221			
1222			
1223			
1224			
1225			
1226			
1227			
1220			
1230			
1231			
1232			
1233			
1234			
1235			
1236			
1237			. .
1238			21
1239			

Figure legends

Fig. 1 Electron micrographic (EM) picture of bacteriophage isolated from wastewater effluent discharge and kinetics of bacterial lysis. (A) Phage with prolate head belonging to the group of Myoviridae, which are double-stranded DNA phages with contractile tails. Dimensions of the phage are shown in nm. (B) Kinetics of E. coli lysis as measured by flow cytometry. Changes in intact cell concentrations are shown for a bacterial suspension with the phage added at an MOI=1. A bacterial suspension without phage served as a control. Error bars represent standard deviations from three independent repeats. Fig. 2 Effect of temperature history of fully grown *E. coli* on susceptibility to phage lysis (A) in comparison with control without phage (B) and corresponding rates of change of OD_{600} (slope analyses). Bacteria (grown at 20° C; $OD_{600} = 1$) were suspended in PBS and exposed to different temperatures (4, 20, 37, 41, 44 or 50°C) for 24 h prior to resuspension in TSB and phage challenge (MOI = 1) at 20 $^{\circ}$ C. Changes in optical densities were monitored using a microplate reader. Error bars show standard deviations from three independent experiments. Rates of change in OD₆₀₀ of *E. coli* were determined by slope analysis and refer to 6 hours after addition of phage as well as for the control without phage. Fig. 3 Change in colony forming units (CFUs) of *E.coli* on MLGA agar within 6 h after exposure to different chlorine disinfection strengths and absence of presence of a bacteriophage. Samples without chlorine addition served as controls. Error bars show standard deviations from three independent repeats.

Fig. 4 Change in colony forming units (CFUs) of *E.coli* on MLGA agar within 6 h after
exposure to different UV doses and absence of presence of a bacteriophage. Samples without
UV irradiation served as controls. Error bars show standard deviations from three
independent repeats.

470 Supplementary Fig. S1 Effect of increasing chlorine disinfection strengths on the
471 culturability of *E. coli* in the absence or presence of a lytic bacteriophage over a time course
472 of 96 hours. Phage (if applicable) was added directly after disinfection and dechlorination at
473 an MOI = 1. Samples were taken after indicated times and bacteria were collected by
474 membrane filtration. Filters were placed on MLGA agar, where *E. coli* forms green colonies.
475 Plates were incubated at 35 °C for 24 h. Samples without chlorine addition served as controls.
476 Representative results from three independent repeats are shown.

478 Supplementary Fig. S2 Effect of increasing UV irradiation doses on the culturability of
479 *E.coli* in the absence or presence of a lytic bacteriophage over a time course of 96 hours.
480 Phage (if applicable) was added directly after disinfection at an MOI = 1. Samples were taken
481 after indicated times and bacteria were collected by membrane filtration. Filters were placed
482 on MLGA agar, where *E. coli* forms green colonies. Plates were incubated at 35 °C for 24 h.
483 Samples without UV irradiation served as controls. Representative results from three
484 independent repeats are shown.

1358			
1359			
1360	485	References	
1362			
1363	486		
1364			
1365	487	Abedon S1 (2012) Bacterial 'immunity' against bacteriophages. Bacteriophage 2:50-54.	
1366	400	Amph FM 2016. The use of heaterianhages as natural biogentral against heaterial	
1367	400	Amen EW. 2010. The use of bacteriophages as natural biocontrol agents against bacteriar	
1368	489	pathogens. PhD thesis Cranfield University.	
1370			
1371	490	http://dspace.lib.cranfield.ac.uk/handle/1826/11331 (accessed 19.10.2017).	
1372			
1373	491	Ankrah NYD, May AL, Middleton JL, Jones DR, Hadden MK, Gooding JR, LeCleir GR,	
1374			
1376	492	Wilhelm SW, Camagna SR, Bucan A (2014) Phage infection of an environmentally	
1377	193	relevant marine bacterium alters host metabolism and lysate composition. ISME I	
1378	775	relevant marine bacterium arters nost metaborism and rysate composition. ISIVIL 3	
1379	494	8:1089–1100.	
1380			
1382	495	Awais R, Fukudomi H, Miyanaga K, Unno H, Tanji Y (2006) A recombinant bacteriophage) -
1383			
1384	496	based assay for the discriminative detection of culturable and viable but nonculturable	
1385	407	Each michig cali (157:117 Distochard Prog 22:052, 850	
1386	47/	Escherichia con 0157.H7. Biotecinioi Flog 22.855-859.	
1388	498	Birch EW, Ruggero NA, Covert MW (2012) Determining Host Metabolic Limitations on	
1389			
1390	499	Viral Replication via Integrated Modeling and Experimental Perturbation. PLoS Comp	ut
1391			
1392	500	Biol 8:e1002746.	
1394	504	Debremere 7 Becombly L Linder $K(2014)$ largester as of recovery of E soliting water	
1395	501	Bonrerova Z, Rosenblum J, Linden K (2014) Importance of recovery of <i>E. con</i> in water	
1396	502	following ultraviolet light disinfection J Environ Eng 141.04014094	
1397			
1398	503	Brüssow H, Hendrix RW (2002) Phage genomics: small is beautiful. Cell 108:13-16.	
1400			
1401	504	Camper AK, McFeters GA (1979).Chlorine injury and the enumeration of waterborne	
1402	505		
1403	505	coliform bacteria. Appl Environ Microbiol 37:633–641.	
1404 1405	506	Charnock C. Kiønnø O (2000) Assimilable organic carbon and biodegradable dissolved	
1406	500	Chamber C, Rjonno C (2000) Assimilatio organic carbon and broacgradable dissorved	
1407	507	organic carbon in Norwegian raw and drinking waters. Water Research 34:2629–2642.	
1408			
1409	508	Chevallereau A, Blasdel BG, De Smet J, Monot M, Zimmermann M, Kogadeeva M, Sauer	U,
1410			
1412	509	Jorth P, Whiteley M, Debarbieux L, Lavigne R (2016) Next-generation "-omics"	
1413			
1414			o :
1415			24
1410			

1417			
1418			
1419 1420	510	approaches reveal a massive alteration of host RNA metabolism during bacteriophage	
1421 1422	511	infection of Pseudomonas aeruginosa. PLoS Genet 12:e1006134.	
1423 1424	512	Clokie MRJ, Millard AD, Letarov AV, Heaphy S (2011) Phages in nature. Bacteriophages	
1425 1426 1427	513	1:31-45.	
1428 1429	514	Cutler TD, Zimmerman JJ (2011) Ultraviolet irradiation and the mechanisms underlying its	3
1430 1431	515	inactivation of infectious agents. Anim Health Res Rev 12:15-23.	
1432 1433	516	DVGW (German Technical and Scientific Association for Gas an Water) (2006) UV device	e
1434 1435	517	for the disinfection of the water supply. German standard W294.	
1436 1437	518	Fernandes E, Martins VC, Nóbrega C, Carvalho CM, Cardoso FA, Cardoso S, Dias J, Deng	3
1438 1439 1440	519	D, Kluskens LD, Freitas PP, Azeredo J (2014). A bacteriophage detection tool for	
1440 1441 1442	520	viability assessment for Salmonella cells. Biosensors and Bioelectronics 52:239-246.	
1443 1444	521	Haas CN, Karra SB (1984) Kinetics of microbial inactivation by chlorine - II Kinetics in th	e
1445 1446	522	presence of chlorine demand. Water Res 18:1451–1454.	
1447 1448	523	Hijnen WAM, Beerendonk EF, Medema GJ (2006) Inactivation credit of UV radiation for	
1449 1450	524	viruses, bacteria and protozoa (oo)cysts in water: A review. Water Res 40:3-22.	
1451 1452	525	Hu JY, Chu XN, Quek PH, Feng YY, Tan XL (2005) Repair and regrowth of <i>Escherichia</i>	
1455 1454 1455	526	<i>coli</i> after low- and medium-pressure ultraviolet disinfection. Water Science Technol:	
1456 1457	527	Water Supply 5:101-108.	
1458 1459	528	Kollu K, Örmeci B (2014) Regrowth potential of bacteria after ultraviolet disinfection in	
1460 1461	529	absence of light and dark repair. J Environ Eng 141:04014069.	
1462 1463	530	Labrie J, Samson E, Moineau, S (2010) Bacteriophage resistance mechanisms. Nature	
1464 1465	531	Reviews Microbiol 8:317-327.	
1466 1467	532	Le Dantec C, Duguet JP, Montiel A, Dumoutier N, Dubrou S, Vincent V (2002) Chlorine	
1468 1469 1470	533	disinfection of atypical mycobacteria isolated from a water distribution system. Appl	
1471 1472	534	Environ Microbiol 68:1025-1032.	
1473 1474			25
1475			

1476			
1477			
1478	535	Mackey BM Miles CA Parsons SE Seymour DA (1991) Thermal denaturation of whole	
1479	555	Mackey Divi, which cry, raisons 5E, beymour Dry (1991) Thermal denaturation of whole	
1480	536	cells and cell components of Escherichia coli examined by differential scanning	
1481		······································	
1483	537	calorimetry. J General Microbiology 137:2361-2374.	
1484			
1485	538	Mamane-Gravetz H, Linden KG (2004) Impact of particle aggregated microbes on UV	
1486			
1487	539	disinfection. In: Proceedings of the American Water Works Association Water Quality	
1488			
1489	540	Technology Conference, November 14–18, San Antonio, USA.	
1490			
1492	541	Maynard ND, Gutschow MV, Birch EW, Covert MW (2011) The virus as metabolic	
1493	540	angineer Biotechnol I 5:686, 604	
1494	J42	engineer. Diotechnor J 5.000–094.	
1495	543	McGrath S van Sinderen D (2007) Bacteriophages: Genetics and molecular biology	
1496			
1497	544	Norfolk, UK, Horizon Scientific Press.	
1499			
1500	545	Munakata N, Kuo J (2016) Disinfection Processes. Water Environ Res 88:1192-1229.	
1501			
1502	546	du Preez M., Kfir R, Coubrough P (1995) Investigation of injury of coliforms after	
1503			
1504	547	chlorination. water Sci Tech 31:115-118.	
1505	548	Nocker A. Cheswick R. Dutheil de la Rochere PM. Denis M. Léziart T. Jarvis P. (2017) Wh	en
1507	540	Notker A, Cheswick R, Dunen de la Rochere I W, Denis W, Leziart I, Jarvis I (2017) wi	CII
1508	549	are bacteria dead? A step towards interpreting flow cytometry profiles after chlorine	
1509			
1510	550	disinfection and membrane integrity staining. Environ Technol 38:891-900.	
1512			
1513	551	Said MB, Masahiro O, Hassan A (2010) Detection of viable but noncultiviable <i>Escherichia</i>	
1514	550	. li - Arn UNV inne disting arrive a latic OR altered Arm Mismakial (0.121-127	
1515	552	con after UV irradiation using a lytic Qp phage. Ann Microbiol 60:121-127.	
1516	553	Shin G. Linden K.G. Arrowood MI. Sobsey MD (2001) Low-pressure UV inactivation and	
1517	555	Shin O, Enden KO, Anowood Wis, Sobsey WD (2001) Low pressure O v indervation and	
1510	554	DNA repair potential of Crystosporidium paryum oocysts. Appl Environ Microbiol	
1520			
1521	555	67:3029-3032.	
1522			
1523	556	Templeton M, Andrews RC, Hofmann R (2006) Impact of iron particles in groundwater on	
1524			
1526	557	the UV inactivation of bacteriophages MS2 and 14. J Appl Microbiol 101:732-741.	
1527	558	White GC (1999) The handbook of chlorination and alternative disinfectants. J. Wiley, New	7
1528	220	white Ge (1999) The handbook of emormation and alternative distinctions. J. whey, iver	v
1529	559	York.	
1530			
1537	560		
1533			26
1534			





Figure 1. Electron micrographic (EM) picture of bacteriophage isolated from wastewater effluent discharge and kinetics of bacterial lysis. (**A**) Phage with prolate head belonging to the group of Myoviridae, which are double-stranded DNA phages with contractile tails. Dimensions of the phage are shown in nm. (**B**) Kinetics of *E. coli* lysis as measured by flow cytometry. Changes in intact cell concentrations are shown for a bacterial suspension with the phage added at an MOI=1. A bacterial suspension without phage served as a control. Error bars represent standard deviations from three independent repeats.

Figure 2



Figure 2. Effect of temperature history of fully grown *E. coli* on susceptibility to phage lysis (A) in comparison with control
without phage (B) and corresponding rates of change of OD₆₀₀ (slope analyses). Bacteria (grown at 20°C; OD₆₀₀ = 1) were
suspended in PBS and exposed to different temperatures (4, 20, 37, 41, 44 or 50°C) for 24 h prior to resuspension in TSB and
phage challenge (MOI = 1) at 20°C. Changes in optical densities were monitored using a microplate reader. Error bars show
standard deviations from three independent experiments. Rates of change in OD₆₀₀ of *E. coli* were determined by slope
analysis and refer to 6 hours after addition of phage as well as for the control without phage.



0 0.02 0.05 0.2 0.5 Chlorine concentrations (mg min L⁻1)

Figure 3. Change in colony forming units (CFUs) of *E.coli* on MLGA agar within 6 h after exposure to different chlorine disinfection strengths and absence of presence of a bacteriophage. Samples without chlorine addition served as controls. Error bars show standard deviations from three independent repeats.

 \blacksquare with phage

□ without phage

-4

Figure 4



Figure 4. Change in colony forming units (CFUs) of *E.coli* on MLGA agar within 6 h after exposure to different UV doses and absence of presence of a bacteriophage. Samples without UV irradiation served as controls. Error bars show standard deviations from three independent repeats.





Figure 1S. Effect of increasing chlorine disinfection strengths on the culturability of *E.coli* in the absence or presence of a lytic
 bacteriophage over a time course of 96 hours. Phage (if applicable) was added directly after disinfection and dechlorination at
 an MOI = 1. Samples were taken after indicated times and bacteria were collected by membrane filtration. Filters were placed
 on MLGA agar, where *E. coli* forms green colonies. Plates were incubated at 35 ° C for 24 h. Samples without chlorine
 addition served as controls. Representative results from three independent repeats are shown.





Figure 2S. Effect of increasing UV irradiation doses on the culturability of *E.coli* in the absence or presence of a lytic
 bacteriophage over a time course of 96 hours. Phage (if applicable) was added directly after disinfection at an MOI = 1.
 Samples were taken after indicated times and bacteria were collected by membrane filtration. Filters were placed on MLGA

³⁶ agar, where *E. coli* forms green colonies. Plates were incubated at 35 ° C for 24 h. Samples without UV irradiation served as controls. Representative results from three independent repeats are shown.