



BIROn - Birkbeck Institutional Research Online

Joly-Mascheroni, Ramiro M. and Senju, Atsushi and Shepherd, Alex J. (2008) Dogs catch human yawns. *Biology Letters* 4 (5), pp. 446-448. ISSN 1744-9561.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/2562/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively



BIROn - Birkbeck Institutional Research Online

Enabling open access to Birkbeck's published research output

Dogs catch human yawns

Journal Article

<http://eprints.bbk.ac.uk/2562>

Version: Accepted (Refereed)

Citation:

Joly-Mascheroni, R.M.; Senju, A.; Shepherd, A.J. (2008)
Dogs catch human yawns -
Biology Letters 4 (5), pp.446-448

© 2008 The Royal Society

[Publisher version](#)

All articles available through Birkbeck ePrints are protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

[Deposit Guide](#)

Contact: lib-eprints@bbk.ac.uk

Biology Letters (2008) 4, 446–448. doi:10.1098/rsbl.2008.0333

Dogs catch human yawns

Ramiro M Joly-Mascheroni¹, Atsushi Senju^{1*} and Alex J Shepherd¹

¹ School of Psychology, Birkbeck, University of London, Malet Street, London WC1E 7HX, UK.

* Author for correspondence (a.senju@bbk.ac.uk)

Summary

This study is the first to demonstrate that human yawns are possibly contagious to domestic dogs (*Canis familiaris*). Twenty-nine dogs observed a human yawning or making control mouth movements. Twenty-one dogs yawned when they observed a human yawning, but control mouth movements did not elicit yawning from any of them. The presence of contagious yawning in dogs suggests that this phenomenon is not specific to primate species and may indicate that dogs possess the capacity for a rudimentary form of empathy. Since yawning is known to modulate levels of arousal, yawn contagion may help coordinate dog-human interaction and communication. Understanding the mechanism as well as the function of contagious yawning between humans and dogs requires more detailed investigation.

Key Words:

yawning, contagious yawning, dog, empathy, social cognition

Introduction

Contagious yawning (i.e., yawning triggered by perceiving others yawning) is a well-documented phenomenon (e.g. Moore, 1942; Platek et al., 2003; Provine, 1986; Senju et al., 2007). Although some claim that contagious yawning is a response to an innate releasing mechanism (Provine, 1986), the relationship between the susceptibility to contagious yawning and self-reported scores of empathy (Platek et al., 2003), as well as the absence of contagious yawning in individuals with autism spectrum disorder (Senju et al., 2007), suggest that it may be related to the capacity for empathy (Preston & De Waal, 2002). Even though spontaneous yawning is widespread among vertebrate species (Baenninger, 1987; Heusner, 1946), contagious yawning has been

reported to occur only in humans and in chimpanzees (*Pan troglodytes*, Anderson et al., 2004). Stumptail macaques (*Macaca arctoides*) also yawn in response to observed yawns, although it is unclear whether this behaviour is similar to contagious yawning found in other species or simply reflects stress (Paukner & Anderson, 2006).

The current study examined whether contagious yawning can be observed in domestic dogs (*Canis familiaris*). Dogs are unusually skilled at reading human social and communicative cues. They can follow human gaze and pointing (Hare et al., 2002; Miklósi et al., 2003, 2006), they can show sensitivity to others' knowledge states (e.g. indicating the location of a hidden toy more frequently to someone not involved in hiding it, than to someone who did the hiding, Virányi et al., 2006) and they are even able to match their own actions to observed human actions (Topál et al., 2006). Dogs' unique social skills in interacting with humans may be the result of selection pressures during the process of domestication (Hare & Tomasello, 2005; Miklósi et al., 2003, 2007). Therefore, there is the potential that dogs may also have developed the capacity for empathy towards humans, and may catch human yawns. However, no empirical studies have been reported that systematically investigate contagious yawning in dogs.

In the current study, dogs observed a human experimenter yawning (yawning condition), or demonstrating non-yawning mouth movements (control condition). If dogs have the capacity for contagious yawning, they should yawn more in the yawning condition than in the control condition.

Methods

Dogs older than 14 weeks participated in the study (n = 29, mean age: 6.4 years, 12 females, 17 males, see Supplementary Table for details). Testing was conducted in places familiar to each dog, between 11 am and 4 pm and after they had finished walking and had one of their two daily meals. Informed consent was obtained from their owners. The study was approved by the Research Ethics Committee of the School of Psychology, Birkbeck, University of London.

The testing consisted of two conditions. Each condition lasted 5 minutes, separated by a 5-minute interval. The dogs were given a small edible treat before starting each condition. In the yawning condition, the experimenter, who was a stranger to the dogs, attracted the dog's attention by calling them by name, and then acted a yawning movement

with vocalization when the dog established eye contact with the experimenter (see Supplementary Figure). The experimenter repeated this sequence for 5 minutes. The number of presented yawns varied between subjects (range: 10-19), mainly because of individual differences in the time required to re-establish eye contact with the experimenter after each yawn. No feedback was given to any of the dog's responses. The control condition was exactly the same as the yawning condition, except that the experimenter displayed non-yawning mouth-opening actions without vocalization, instead of yawns (see Supplementary Figure). The order of testing conditions was counterbalanced between subjects (see Supplementary Table for details). During the testing, the owner sat behind the dog quietly.

During the experiment, the experimenter recorded the number of yawns elicited. In addition, the dogs' behaviours were video-recorded and coded by an independent observer, who was not informed of the purpose of the study. The number of yawns recorded by the second coder was identical to that of the experimenter for all of the dogs, which demonstrates a high reliability of the coding.

Results

The yawning condition elicited yawns for 21 out of 29 dogs (one example is shown in Figure 1), and no dogs yawned in the control condition ($p < .001$, McNemar's test, McNemar, 1947). No yawns were observed during the interval between conditions either. On average, dogs yawned 1.9 times in the yawning condition (range: 0-5, see Supplementary Table). On average, it took 1 min 39 s for the dogs to yawn (standard deviation: 1 min 28 s, range: 5 s - 4 min 47 s), after the experimenter had presented 4.5 yawns (standard deviation: 4.4, range: 1 - 17). The number of yawns in the yawning condition did not differ significantly between sexes ($z = -1.30$, $p > .1$, Wilcoxon test) and was not affected by the order of testing (i.e. yawning or control condition first, $z = -1.27$, $p > .1$, Wilcoxon test). Age did not correlate with the number of yawns ($r = .14$, $p > .1$, Spearman's test). The correlation between the number of presented yawns and the number of elicited yawns did not reach significance ($r = -.36$, $p > .05$, Spearman's test).



Figure 1. A yawn response during the presentation of a human yawn. The experimenter's face can be seen at the top of each image, which was recorded in the mirror placed behind the dog. The dog observed the experimenter yawning (left), started yawning as the experimenter finished yawning (middle) and then completed yawning (right).

Discussion

The current study demonstrates that human yawns are possibly contagious to dogs. The presentation of human yawning elicited yawns in 72 % of the dogs tested, which is higher than the rate reported in humans (45-60%) and chimpanzees (33%). This effect cannot be attributed to a general effect of the mere presence of unfamiliar humans, or to the observation of human mouth movements in general, because no dogs yawned in the control condition. This study is the first to demonstrate that the observation of yawning elicits yawning in a non-primate species, as well as the first demonstration of possible contagious yawning between different species. Since yawning is known to modulate the level of arousal (Daquin et al., 2001), such temporally synchronized occurrences of yawning may help coordinate interactions as well as communication between humans and dogs.

The high yawning rate may be due to displaying 'live' acted yawns rather than video-recorded yawns that have been used in some previous studies (Anderson et al., 2004; Paukner & Anderson, 2006; Senju et al., 2007). It is, however, highly unlikely that the present results could be influenced by the experimenter unintentionally reinforcing dog's yawning in the yawning condition. The main analysis is based on dichotomous data (i.e. the dog yawned or did not yawn in each condition). Reinforcement of chance yawns could only account for a difference in absolute number of yawns per dog, and not the presence/absence of yawns for a dog in each condition.

There are at least two possibilities why human yawns elicit yawning in dogs. Firstly, the susceptibility to contagious yawning of dogs may relate to their capacity for empathy. In humans (Platek et al.,

2003; Senju et al., 2007) and in chimpanzees (Anderson et al., 2004), several studies have suggested that contagious yawning relates to the capacity for empathy, although the mechanism underlying this relation is still unclear. As described above, dogs have exceptional capacities to decode social signals from humans, possibly as a result of the domestication process (Hare & Tomasello, 2005; Miklósi et al., 2003, 2007). Therefore, it is also possible that they have the capacity for empathy, such as representing humans' actions and modulating their own behavioural and autonomic responses accordingly (see also Tópal et al., 2006), and that this underlies contagious yawning. Alternatively, dogs may have developed the capacity for contagious yawning during past synchronous yawning experiences with humans (e.g. Heyes & Ray, 2000), either by sharing (e.g. boring) experiences with humans or by observing humans yawn contagiously to dog's yawns.

Secondly, it is also possible that the dogs' yawns may have been induced by mildly heightened tension or stress. For example, macaques are known to exhibit 'emotion yawns' or 'social yawns' during antagonistic social encounters (Deputte, 1994; Paukner & Anderson, 2006; Smith, 1999). Thus, we cannot rule out the possibility that human yawns are perceived as antagonistic to the dogs. Previous studies have demonstrated that dogs may yawn in response to an encounter with novel humans (Beerda et al., 1998; Hennessy et al., 2006). Some have argued that it is a response to acute stress (Beerda et al., 1998), although this argument is not supported by more recent empirical studies (e.g. Hennessy et al., 1998; Rooney et al., 2007). However, as physiological measurements of stress, such as saliva/blood cortisol or heart rate, were not available for the current sample, and because there has not been enough consensus and standardization on how to interpret dogs' behaviours (Diederich & Giffroy, 2006), further studies, ideally with concurrent physiological measurements, are required to test whether contagious yawning between humans and dogs is mediated by a stress- or tension-related response. At least, the current results cannot be attributed to more general factors related to stress, such as an encounter with a strange human or to the recording equipment, as no dogs yawned in the control condition

Although the current study cannot discriminate between these two possible explanations, further studies could explore the mechanism underlying, as well as the evolutionary origin of, contagious yawning and its relationship to the capacity for empathy. For example, the current study demonstrated that dogs appear to catch human yawns.

However, for practical reasons, we did not examine whether they yawn contagiously to conspecifics' yawns. If the dogs' capacity for contagious yawning has evolved with the capacity for reading human communicative signals, it is possible that dogs are more sensitive to human yawns than dogs' yawns. In contrast, if they have acquired the capacity for contagious yawning as an adaptation for within-species social interaction and communication, which has transferred to dog-human interaction, they may be more sensitive to the yawns of conspecifics. Such contrasts may help highlight the functional specialization of contagious yawning in dogs and, possibly, in other species as well.

Acknowledgments

We thank Chad Arlt, Gergely Csibra, Juliane Kaminski, Toshikazu Hasegawa and anonymous reviewers for their inputs. Atsushi Senju was supported by an ESRC/MRC Interdisciplinary Postdoctoral Fellowship.

References

- Anderson, J. R., Myowa-Yamakoshi, M., & Matsuzawa, T. 2004 Contagious yawning in chimpanzees. *Proc. R. Soc. B.* **271**(Suppl. 6), S468-S470. (doi:10.1098/rsbl.2004.0224)
- Baenninger, R 1987 Some comparative aspects of yawning in *Betta splendens*, *Homo sapiens*, *Panthera leo*, and *Papio sphinx*. *J. Comp. Psychol.* **101**, 349-354.
- Beerda, B., Schilder, M. B. H., van Hooff, J. A. R. A. M., de Vries, H. W., & Mol, J. A. 1998 Behavioural, saliva cortisol and heart rate responses to different types of stimuli in dogs. *Appl. Anim. Behav. Sci.*, **58**, 365-381.
- Daquin, G., Micallef, J., & Blin, O. 2001 Yawning. *Sleep Med. Rev.*, **5**, 299-312.
- Deputte, B. L. (1994). Ethological study of yawning in primates. 1. Quantitative analysis and study of causation in 2 species of Old World monkeys (*Cercocebus albigena* and *Macaca fascicularis*). *Ethology*, **98**, 221-245.
- Diederich, C., & Giffroy, J.-M. 2006 Behavioural testing in dogs: A review of methodology in search for standardisation. *Appl. Anim. Behav. Sci.*, **97**, 51-72. (doi:10.1016/j.applanim.2005.11.018)
- Hare, B., Brown, M., Williamson, C., & Tomasello, M. 2002 The domestication of social cognition in dogs. *Science*, **298**, 1634-1636.
- Hare, B., & Tomasello, M. 2005 Human-like social skills in dogs? *Trends*

- Cogn. Sci.*, **9**, 439-444. (doi: 10.1016/j.tics.2005.07.003)
- Heyes, C. M., & Ray, E. D. 2000 What is the significance of imitation in animals? *Adv. Study Behav.* **29**, 215-245.
- Hennessy, M. B., Morris, A., & Linden, F. 2006 Evaluation of the effects of a socialization program in a prison on behavior and pituitary-adrenal hormone levels of shelter dogs. *Appl. Anim. Behav. Sci.*, **99**, 157-171. (doi:10.1016/j.applanim.2005.09.011)
- Heusner, A. P. 1946 Yawning and associated phenomena. *Physiol. Rev.*, **26**, 156-168.
- McNemar, Q. 1947 Note on the sampling error of the difference between correlated proportions or percentages. *Psychometrika*, **12**, 153-157.
- Miklósi, Á., Kubinyi, E., Topál, J., Gácsi, M., Virányi, Z., & Csányi, V. 2003 A simple reason for a big difference: Wolves do not look back at humans, but dogs do. *Curr. Biol.*, **13**, 763-766.
- Miklósi, Á., & Soproni, K. 2006 A comparative analysis of animals' understanding of the human pointing gesture. *Anim. Cogn.* **9**, 81-93. (doi: 10.1007/s10071-005-0008-1)
- Miklósi, Á., Topál, J., & Csányi, V. 2007 Big thoughts in small brains? Dogs as a model for understanding human social cognition. *Neuroreport*, **18**, 467-471.
- Moore, J. E. 1942 Some psychological aspects of yawning. *J. Gen. Psychiatry*, **27**, 289-294.
- Paukner, A., & Anderson, J. R. 2006 Video-induced yawning in stump-tail macaques (*Macaca arctoides*). *Biol. Lett.* **2**, 36-38. (doi:10.1098/rsbl.2005.0411)
- Platek, S. M., Critton, S. R., Myers, T. E., & Gallup, G. G. 2003 The role of self-awareness and mental state attribution, *Cogn. Brain Res.* **17**, 223-227. (doi:10.1016/S0926-6410(03)00109-5)
- Preston, S. D. & de Waal, F. B. 2002 Empathy: Its ultimate and proximate bases. *Behav. Brain Sci.* **25**, 1-20.
- Provine, R. R. 1986 Yawning as a stereotyped action pattern and releasing stimulus. *Ethology*, **72**, 448-455.
- Rooney, N. J., Gaines, S. A., & Bradshaw, J. W. S. 2007 Behavioural and glucocorticoid responses of dogs (*Canis familiaris*) to kennelling: Investigating mitigation of stress by prior habituation. *Physiol. Behav.* **92**, 847-854. (doi:10.1016/j.physbeh.2007.06.011)
- Senju, A., Maeda, M., Kikuchi, Y., Hasegawa, T., Tojo, Y., & Osanai, H. 2007 Absence of contagious yawning in children with autism spectrum disorder. *Biol. Lett.*, **3**, 706-708. (doi: 10.1098/rsbl.2007.0337)
- Smith, E. 1999 Yawning: An evolutionary perspective. *Hum. Evol.*, **14**,

191-198. (10.1007/BF02440156)

- Topál, J., Byrne, R., Miklósi, Á., & Csányi, V. 2006 Reproducing human actions and action sequences: "Do as I Do!" in a dog. *Anim. Cogn.*, **9**, 355-367. (doi:10.1007/s10071-006-0051-6)
- Virányi, Z., Topál, J., Miklósi, Á., & Csányi, V. 2006 A nonverbal test of knowledge attribution: a comparative study on dogs and children. *Anim. Cogn.*, **9**, 13-26. (doi:10.1007/s10071-005-0257-z)