



BIROn - Birkbeck Institutional Research Online

Hart, Sarah and Sbeiti Clarke, Amal (2020) Commuting Involution Graphs in classical affine Weyl Groups. *Communications in Algebra* 48 (7), pp. 2941-2957. ISSN 0092-7872.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/30779/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

Commuting Involution Graphs in Classical Affine Weyl Groups

Sarah Hart and Amal Sbeiti Clarke

Abstract

In this paper we investigate commuting involution graphs in classical affine Weyl groups. Let W be a classical Weyl group of rank n , with \widetilde{W} its corresponding affine Weyl group. Our main result is that if X is a conjugacy class of involutions in \widetilde{W} , then the commuting involution graph $\mathcal{C}(\widetilde{W}, X)$ is either disconnected or has diameter at most $n + 2$. This bound is known to hold for types \widetilde{A}_n and \widetilde{C}_n , so the main work of this paper is to prove the theorem for types \widetilde{B}_n and \widetilde{D}_n .

1 Introduction

Let G be a group, and X a subset of G . The *commuting graph* $\mathcal{C}(G, X)$ is the graph with vertex set X , with an edge joining vertices x and y whenever x and y commute in G . An early use of commuting graphs, in the case where X is the set of all non-identity elements of G , was in Brauer and Fowler's paper on groups of even order [5], where they showed that if G is a group of even order with more than one conjugacy class of involutions, then any two involutions are distance at most 3 apart in $\mathcal{C}(G, X)$.

If X is a set of involutions, then we call $\mathcal{C}(G, X)$ a *commuting involution graph*. Such graphs have been studied in a wide variety of groups, most often in the cases where X is a conjugacy class of involutions, or the set of all involutions, of G . A well-known example of the use of commuting involution graphs is Fischer's work on 3-transposition groups [7], where in that case, X was a conjugacy class of involutions, the product of any pair of which had order at most 3.

There has been a lot of interest in commuting graphs in the past few years, particularly around questions relating to connectedness and diameters. Writing $\mathcal{C}(G)$ for the commuting graph $\mathcal{C}(G, G \setminus Z(G))$, Iranmanesh and Jafarzadeh [12] conjectured that if G is finite, then $\mathcal{C}(G)$ is either disconnected or has diameter bounded above by a constant independent of the group G . This was disproved by Giudici and Parker [8], who showed that for all positive integers d , there is a finite 2-group G such that $\mathcal{C}(G)$ has diameter greater than d . However, Morgan and Parker [13] subsequently showed that if G is a finite group with trivial centre, then either $\mathcal{C}(G)$ is connected, or its diameter is at most 10.

This article is concerned with commuting involution graphs in classical affine groups. These groups, being Coxeter groups, each have an associated Coxeter graph, where vertices correspond to simple reflections, and pairs of vertices are joined by an edge precisely when they do *not* commute. In fact, given a Coxeter group W with generating set R of simple reflections, if we remove the edge labels from the Coxeter graph, then we obtain the complement of the commuting involution graph $\mathcal{C}(W, R)$. The reason for considering

commuting involution graphs in the context of these particular Coxeter groups is that it represents a natural continuation of a series of papers by Bates et al., looking at commuting involution graphs for various groups G , where X is an involution conjugacy class ([1], [2], [3], [4]). In particular, they showed in [1] that in the symmetric group, $\mathcal{C}(G, X)$ is either disconnected or has diameter at most 4. Then, in [2], they extended this to show that if G is any finite Coxeter group, then $\mathcal{C}(G, X)$ is either disconnected or has diameter at most 5. Later, Perkins [14] looked at affine groups of type \tilde{A}_n , where it turns out that when $\mathcal{C}(G, X)$ is connected it has diameter at most 6. Most recently, Hart and Sbeiti Clarke [9], considered the case of \tilde{C}_n . Here, $\mathcal{C}(G, X)$, when connected, has diameter at most $n+2$. The purpose of this paper is to complete the analysis of classical affine groups by dealing with types \tilde{B}_n and \tilde{D}_n . Our main result is as follows.

Theorem 1.1. *Let W be a classical Weyl group of rank n and \tilde{W} its corresponding affine group, with X a conjugacy class of involutions of \tilde{W} . Then either $\mathcal{C}(\tilde{W}, X)$ is disconnected, or it is connected with diameter at most $n+2$.*

This paper is structured as follows. In the remainder of this section, we summarise results for finite classical Weyl groups that we will need. In Section 2, we describe the labelled cycle form for affine involutions, first introduced in [9] for type \tilde{C}_n , which we will use in Section 3 to develop a characterisation of involution conjugacy classes in types \tilde{B}_n and \tilde{D}_n . In Section 4, we state and prove detailed results on connectedness and diameter of commuting involution graphs in types \tilde{B}_n and \tilde{D}_n . In particular we obtain necessary and sufficient conditions under which $\mathcal{C}(\tilde{W}, X)$ is connected, and show that when it is connected, its diameter is at most $n+2$, thus proving Theorem 1.1 in these cases. Finally, in Section 5, we briefly discuss the exceptional affine groups.

In the remainder of this section, we will review facts about involution conjugacy classes in the classical finite Weyl groups. Throughout this paper, we will use the convention that a Coxeter group of type Γ is denoted $W(\Gamma)$, where Γ is the associated Coxeter graph.

Let W be of type A_{n-1} . Then $W \cong \text{Sym}(n)$, and W acts on \mathbb{R}^n by permuting the subscripts of the standard basis $\{e_1, \dots, e_n\}$. We view the elements of W as permutations in the usual way. If W is of type B_n or D_n , then we view the elements of W as signed permutations; they act on \mathbb{R}^n by permuting the subscripts of the standard basis $\{e_1, \dots, e_n\}$ and changing their signs. For example, given $w = (\bar{1} \ 2^+ \ \bar{3}) \in W(B_n)$, we have $w(e_1) = -e_2$, $w(e_2) = e_3$ and $w(e_3) = -e_1$.

Expressing σ as a product of disjoint cycles, we say that a cycle $(i_1 \cdots i_r)$ of σ is *positive* if there is an even number of minus signs above its elements, and *negative* if the number of minus signs is odd. For example, $(\bar{1} \ \bar{3} \ 2)$ is a negative cycle, whereas $(\bar{4} \ \bar{5})$ is positive. The group $W(B_n)$ consists of all signed permutations of n , while $W(D_n)$ is the subgroup of index 2 consisting of signed permutations with an even number of minus signs. We record here for reference the characterisation of conjugacy classes in the groups $W(A_{n-1})$, $W(B_n)$ and $W(D_n)$. The conjugacy classes in these groups have been known for almost a century, but for a modern treatment see Carter [6].

Theorem 1.2.(i) *Elements of $W(A_{n-1})$ are conjugate if and only if they have the same cycle type. Involution conjugacy classes are parameterised by the number m of transpositions in involutions of the class.*

(ii) Elements of $W(B_n)$ are conjugate if and only if they have the same signed cycle type. Involutions contain only positive 2-cycles, negative 1-cycles and positive 1-cycles (fixed points). There is exactly one conjugacy class of involutions for each triple (m, k, l) where m is the number of transpositions, k is the number of negative 1-cycles, l is the number of fixed points and $2m + k + l = n$.

(iii) Conjugacy classes in $W(D_n)$ are parametrised by signed cycle type, with one class for each signed cycle type except in the case where the signed cycle type contains only even length, positive cycles, where there are two classes for each signed cycle type. Involutions contain only positive 2-cycles, negative 1-cycles, and fixed points, with the number of negative 1-cycles always being even. As in $W(B_n)$, there is one conjugacy class for each triple (m, k, l) , except for the case $(m, 0, 0)$ (so that $2m = n$). In this case, elements are conjugate precisely when the number of minus signs in their expressions as products of disjoint signed cycles is congruent modulo 4.

For example, in $W(D_4)$, $(\overset{+}{1}\overset{+}{2})(\overset{+}{3}\overset{+}{4})$ is conjugate to $(\bar{1}\bar{2})(\bar{3}\bar{4})$, but not to $(\bar{1}\bar{2})(\overset{+}{3}\overset{+}{4})$.

We write $\text{Diam } \mathcal{C}(G, X)$ for the diameter of $\mathcal{C}(G, X)$ when it is a connected graph, in other words, the maximum distance $d(x, y)$ between any $x, y \in X$ in the graph. We state here for reference the known results on connectedness and diameters for finite classical Weyl groups.

Theorem 1.3 (Theorems 1.1 and 1.2 of [1]). *Let $W = W(A_{n-1})$ and X be a conjugacy class of involutions having m transpositions and l fixed points (where $2m + l = n \geq 2$).*

- (i) *If $l = 1$, then $\mathcal{C}(W, X)$ is disconnected.*
- (ii) *If $n = 4$ and $m = 1$, then $\mathcal{C}(W, X)$ is disconnected.*
- (iii) *If $n \in \{6, 8, 10\}$ and $l = 2$, then $\text{Diam } \mathcal{C}(W, X) = 4$.*
- (iv) *In all other cases, $\text{Diam } \mathcal{C}(W, X) \leq 3$.*

Theorem 1.4 (Theorem 1.1 of [2]). *Suppose that W is of type B_n ($n \geq 2$) or D_n ($n \geq 4$), and let X be a conjugacy class of involutions with signed cycle type (m, k, l) , where $2m + k + l = n$. Let $t := \max\{k, l\}$. Then the following hold.*

- (i) *If $m = 0$, then $\mathcal{C}(W, X)$ is a complete graph.*
- (ii) *If $t = 0$, then $\text{Diam } \mathcal{C}(W, X) \leq 2$.*
- (iii) *If $t = 1$ and $m > 0$, then $\mathcal{C}(W, X)$ is disconnected.*
- (iv) *If $t \geq 2$ and $n > 5$, then $\text{Diam } \mathcal{C}(W, X) \leq 4$.*
- (v) *If $n = 5$, $m = 1$ and $t = 2$ then $\text{Diam } \mathcal{C}(W, X) = 5$. If $n = 5$, $m = 1$ and $t = 3$ then $\text{Diam } \mathcal{C}(W, X) = 2$. Finally, if $n = 4$, $m = 1$ and $t = 2$, then $\mathcal{C}(W, X)$ is disconnected.*

2 Affine involutions and the labelled cycle form

Our job in this section is to establish what involutions in classical affine Weyl groups look like. Let W be a finite Weyl group, with root system Φ contained in a Euclidean vector space $V \cong \mathbb{R}^n$. For a given root α , recall that the corresponding coroot α^\vee is given by $\alpha^\vee = \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, and we write Φ^\vee for the set of all coroots. The affine Weyl group \tilde{W} is then

the semidirect product of W with the translation group of the coroot lattice $L = L(\Phi^\vee)$. We usually identify the translation group with L . Then, any $w \in \widetilde{W}$ can be written in the form (σ, \mathbf{v}) where $\sigma \in W$ and $\mathbf{v} = (v_1, \dots, v_n) \in L$. See, for example, [11, Chapter 4] for more details.

For $\sigma, \tau \in W$ and $\mathbf{u}, \mathbf{v} \in L$ we have

$$(\sigma, \mathbf{v})(\tau, \mathbf{u}) = (\sigma\tau, \mathbf{v}^\tau + \mathbf{u}).$$

We note that $(\sigma, \mathbf{v})^{-1} = (\sigma^{-1}, -\mathbf{v}^{\sigma^{-1}})$. Moreover, the element (σ, \mathbf{v}) is conjugate to (τ, \mathbf{u}) via some (g, \mathbf{w}) if and only if

$$(\tau, \mathbf{u}) = (\sigma, \mathbf{v})^{(g, \mathbf{w})} = (g^{-1}\sigma g, \mathbf{v}^g + \mathbf{w} - \mathbf{w}^{g^{-1}\sigma g}). \quad (1)$$

The reflections of \widetilde{W} are the affine reflections $s_{\alpha, k}$ ($\alpha \in \Phi$, $k \in \mathbb{Z}$). Recall that, for \mathbf{v} in V ,

$$\begin{aligned} s_\alpha(\mathbf{v}) &= \mathbf{v} - 2\frac{\langle \alpha, \mathbf{v} \rangle}{\langle \alpha, \alpha \rangle} \alpha = \mathbf{v} - \langle \alpha, \mathbf{v} \rangle \alpha^\vee \\ s_{\alpha, k}(\mathbf{v}) &= \mathbf{v} - 2\frac{(\langle \alpha, \mathbf{v} \rangle - k)}{\langle \alpha, \alpha \rangle} \alpha = s_\alpha(\mathbf{v}) + k\alpha^\vee. \end{aligned}$$

If R is a set of simple reflections for W and $\tilde{\alpha}$ is the highest root (that is, the root with the highest coefficient sum when expressed as a linear combination of simple roots), then it can be shown that $R \cup \{s_{\tilde{\alpha}, 1}\}$ is a set of simple reflections for \widetilde{W} .

We will now focus on the classical affine groups. Since it turns out that

$$W(\tilde{A}_n) \leq W(\tilde{D}_n) \leq W(\tilde{B}_n) \leq W(\tilde{C}_n),$$

we assume for the moment that $\widetilde{W} = W(\tilde{C}_n)$. The Coxeter graph \tilde{C}_n is as follows:

$$\tilde{C}_n(n \geq 2) \quad \begin{array}{ccccccc} r_1 & r_2 & r_3 & \dots & r_{n-1} & r_n & r_{n+1} \\ \bullet & \bullet & \bullet & \dots & \bullet & \bullet & \bullet \end{array}$$

The root system Φ of type C_n consists of the long roots $\{2e_i : 1 \leq i \leq n\}$ and the short roots $\{\pm e_i \pm e_j : 1 \leq i < j \leq n\}$. Therefore, the set of coroots contains $\{e_1, \dots, e_n\}$, meaning that the coroot lattice here is just $\{(\lambda_1, \dots, \lambda_n) : \lambda_i \in \mathbb{Z}\} = \mathbb{Z}^n$. So $W(\tilde{C}_n)$ consists of all elements $w = (\sigma, \mathbf{v})$ where σ is a signed permutation and $\mathbf{v} \in \mathbb{Z}^n$. For the simple roots of Φ we can take $2e_1, e_1 - e_2, \dots, e_{n-1} - e_n$, with corresponding reflections $(\bar{1}), (\bar{1} \bar{2}), \dots, (n-1 \ n)$. The highest root is $2e_n$. Consequently, we can set $r_1 = ((\bar{1}), \mathbf{0})$, $r_i = ((i-1 \ i), \mathbf{0})$ for $2 \leq i \leq n$, and $r_{n+1} = ((\bar{n}), e_n)$.

By the definition of group multiplication in \widetilde{W} , we see that the element (σ, \mathbf{v}) of \widetilde{W} is an involution precisely when $(\sigma^2, \mathbf{v}^\sigma + \mathbf{v}) = (1, \mathbf{0})$. This allows us to characterise the involutions in \widetilde{W} .

Lemma 2.1 (Lemma 2.1 of [9]). *A non-identity element (σ, \mathbf{v}) of \widetilde{W} is an involution if and only if σ , when expressed as a product of disjoint signed cycles, has the form*

$$\sigma = (a_1^+ b_1^+) \cdots (a_t^+ b_t^+) (a_{t+1}^- b_{t+1}^-) \cdots (a_m^- b_m^-) (c_{2m+1}^-) \cdots (c_{n-l}^-) (d_{n-l+1}^-) \cdots (d_n^-)$$

for some a_i, b_i, c_i, d_i, t, m and l ; and, writing $\mathbf{v} = (v_1, \dots, v_n)$, we have $v_{b_i} = -v_{a_i}$ when σ contains $(a_i^+ b_i^+)$, $v_{b_i} = v_{a_i}$ when σ contains $(a_i^- b_i^-)$ and $v_{d_i} = 0$ for $n-l < i \leq n$.

As described in [9], Lemma 2.1 allows us to use a shorthand for writing involutions of \widetilde{W} . For an element (w, \mathbf{v}) of \widetilde{W} with $\mathbf{v} = (v_1, \dots, v_n)$, above each signed number i in the expression of w as a product of disjoint signed cycles we write v_i . For example, we write $((1^{\overline{-}} 2^{\overline{+}} 3^{\overline{-}}), 4e_1 + 5e_2 + 6e_3)$ as $(1^{\overline{4}} 2^{\overline{5}} 3^{\overline{6}})$. Suppose (σ, \mathbf{v}) is an involution with the form given in Lemma 2.1. For transpositions $(a_i^{\pm} b_i^{\pm})$ of σ , where the number above a_i determines the number above b_i as described in Lemma 2.1, we write $(a_i^{\overline{\lambda}} b_i^{\overline{\lambda}})$ for $(a_i^{\overline{+}} b_i^{\overline{+}})$ and $(a_i^{\overline{\lambda}} b_i^{\overline{\lambda}})$ for $(a_i^{\overline{-}} b_i^{\overline{-}})$. We call this the *labelled cycle form* of (σ, \mathbf{v}) . Where it is helpful, we adopt the convention that cycles $(d_i^{\overline{0}})$ are omitted, as these fix both d_i and e_{d_i} . In labelled cycle form, the simple reflections of $W(\widetilde{C}_n)$ become $r_1 = (\overline{1}^{\overline{0}})$, $r_i = (i-1^{\overline{+}} i)$ for $2 \leq i \leq n$, and $r_{n+1} = (\overline{n}^{\overline{1}})$.

Notation 2.2. For a vector $\mathbf{v} = (v_1, \dots, v_n)$ we write $\sum \mathbf{v}$ for $\sum_{i=1}^n v_i$, the coordinate sum of \mathbf{v} . For a signed permutation σ , we let $\text{Neg}(\sigma)$ be the number of minus signs in σ . By extension, for an element $x = (\sigma, \mathbf{v})$ of $W(\widetilde{C}_n)$ we write $\sum x$ for the coefficient sum of \mathbf{v} and $\text{Neg}(x)$ for the number of minus signs in σ . Suppose in addition that x is an involution. Then x can be written in labelled cycle form, as described in Lemma 2.1, for suitable a_i, b_i, c_i, d_i, t, m and l . We then define $f(x) = 2 \sum_{i=1}^m v_{a_i} + \sum_{j=1}^{n-2m-l} v_{c_{2m+j}}$. That is, $f(x)$ is equal to twice the sum of the numbers over the 2-cycles of x , plus the sum of the numbers over the negative 1-cycles of x .

We will show in Section 3 that the way in which certain conjugacy classes of $W(\widetilde{C}_n)$ split in $W(\widetilde{B}_n)$ and $W(\widetilde{D}_n)$ is dependent upon the functions Neg and f . We note that we will only ever need to know the value of $f(x)$ up to congruence modulo 4, so any ambiguity about the choice of ordering of a_i and b_i in transpositions will not matter.


Example 2.3. The involution

$$x = \left((1^{\overline{+}} 2^{\overline{+}}) (3^{\overline{-}} 4^{\overline{-}}) (5^{\overline{-}}) (6^{\overline{-}}) (7^{\overline{+}}), (1, -1, 3, 3, 2, 4, 0) \right) = (1^{\overline{1}} 2^{\overline{3}}) (3^{\overline{2}} 4^{\overline{4}}) (7^{\overline{0}})$$

in \widetilde{C}_7 has $m = 2$ disjoint 2-cycles, and $l = 1$ fixed point. Thus $n - 2m - l = 2$, and


$$\begin{aligned} \sum x &= 1 - 1 + 3 + 3 + 2 + 4 + 0 = 12, \\ \text{Neg}(x) &= 4, \\ f(x) &= 2(v_{a_1} + v_{a_2}) + (v_{c_5} + v_{c_6}) = 2(1 + 3) + (2 + 4) = 14. \end{aligned}$$

There is a subgroup of index 2 in $W(\tilde{C}_n)$ which is of type \tilde{B}_n , having the following Coxeter graph.




Here r_1, \dots, r_n are the same as in the \tilde{C}_n graph, and we set $s = (n-1 \frac{1}{n})$. This agrees with the fact that the B_n root system has short roots e_i and long roots $\pm e_i \pm e_j$, meaning that the highest root is $e_{n-1} + e_n$. Note that all the generators of this group are elements (σ, \mathbf{v}) satisfying the property that the coefficient sum of \mathbf{v} is even, and it is not difficult to see that in fact every element with this property lies in $W(\tilde{B}_n)$. That is, $W(\tilde{B}_n) = \{(\sigma, \mathbf{v}) \in W(\tilde{C}_n) : \sum \mathbf{v} \in 2\mathbb{Z}\}$, a subgroup of index 2 in $W(\tilde{C}_n)$.

It is also useful to consider another copy of $W(\tilde{B}_n)$ in $W(\tilde{C}_n)$, which we will denote $\overline{W}(\tilde{B}_n)$, whose Coxeter graph is as shown.



In this case, r_2, \dots, r_{n+1} are as in $W(\tilde{C}_n)$, and we set $t = (1 \frac{1}{2})$. It is not hard to see that $\overline{W}(\tilde{B}_n) = \{w \in W(\tilde{C}_n) : \sum w \equiv \text{Neg}(w) \pmod{2}\}$. Certainly, the set of such elements has index 2 in $W(\tilde{C}_n)$ and contains $\overline{W}(\tilde{B}_n)$; hence it must exactly be $\overline{W}(\tilde{B}_n)$. We observe that $W(\tilde{B}_n)$ and $\overline{W}(\tilde{B}_n)$ are interchanged by the automorphism of $W(\tilde{C}_n)$ induced by the nontrivial graph automorphism ω of its Coxeter graph, which interchanges r_i with r_{n+2-i} for each i .

The intersection of $W(\tilde{B}_n)$ and $\overline{W}(\tilde{B}_n)$ is another affine Coxeter group, this time of type $W(\tilde{D}_n)$, with Coxeter graph



where r_i, s and t are as defined in $W(\tilde{B}_n)$ and $\overline{W}(\tilde{B}_n)$.

Finally, for completeness, we note that we can consider $W(\tilde{A}_{n-1})$ to be the subgroup of $W(\tilde{D}_n)$ whose elements have no minus signs.

3 Conjugacy classes of involutions

In this section we give a characterisation of involution conjugacy classes in terms of *labelled cycle types*.

Definition 3.1. Let a be an involution in $W(\tilde{C}_n)$. The *labelled cycle type* of a is the tuple (m, k_e, k_o, l) , where m is the number of transpositions, k_e is the number of negative 1-cycles with an even number above them, k_o is the number of negative 1-cycles with an odd number above them, and l is the number of positive 1-cycles (fixed points), in the labelled cycle form of a .

For example, the labelled cycle type of $(12)^{\overset{0}{+}}(3)^{\overset{2}{-}}(4)^{\overset{1}{-}}(5)^{\overset{3}{-}}(6)^{\overset{0}{-}}(7)^{\overset{3}{-}}(8)^{\overset{0}{+}}$ is $(1, 2, 3, 1)$.

Theorem 3.2 (Theorem 2.6 of [9]). *Involutions in $W(\tilde{C}_n)$ are conjugate if and only if they have the same labelled cycle type. In particular, every involution is conjugate to exactly one element $a = a_{m, k_e, k_o, l}$ of the form*

$$a = (1\ 2)^{\overset{0}{+}} \cdots (2m-1\ 2m)^{\overset{0}{+}} (2m+1)^{\overset{0}{-}} \cdots (2m+k_e)^{\overset{0}{-}} (2m+k_e+1)^{\overset{1}{-}} \cdots (n-l)^{\overset{1}{-}} (n-l+1)^{\overset{0}{+}} \cdots (n)^{\overset{0}{+}}.$$

For completeness, we state the corresponding result for type \tilde{A}_n . See [14] for details. Since elements of $W(\tilde{A}_{n-1})$ have no minus signs, involutions here contain transpositions $(a\ b)^{++}$ and fixed points (positive 1-cycles) only, and the positive 1-cycles must have zeroes above them. We may therefore omit the signs and 1-cycles and write each involution in the form

$$(a_1\ b_1)^{\lambda_1} (a_2\ b_2)^{\lambda_2} \cdots (a_m\ b_m)^{\lambda_m}.$$

Theorem 3.3 (see [14]). *Involutions in $W(\tilde{A}_{n-1})$ are conjugate if and only if they have the same number of transpositions, except in the case where there are no fixed points. In this case there are two conjugacy classes, and elements $\prod_{i=1}^m (a_i\ b_i)^{\lambda_i}$ and $\prod_{i=1}^m (c_i\ d_i)^{\mu_i}$ are conjugate if and only if $\sum_{i=1}^m \lambda_i \equiv \sum_{i=1}^m \mu_i \pmod{2}$.*

For example, in $W(\tilde{A}_3)$, $(1\ 2)^{\overset{0}{-}}(3\ 4)^{\overset{0}{-}}$ and $(1\ 2)^{\overset{1}{-}}(3\ 4)^{\overset{1}{-}}$ are conjugate to each other, but not to $(1\ 2)^{\overset{1}{-}}(3\ 4)^{\overset{0}{-}}$.

We now move onto $W(\tilde{B}_n)$ and $W(\tilde{D}_n)$. We recall an elementary result from group theory about conjugacy classes in an index-2 subgroup H of a group G . For x in H , equality of the conjugacy classes x^H and x^G is equivalent to the centralizer $C_G(x)$ containing an element outside of H . If $x^H \neq x^G$, then x^G splits as two conjugacy classes in H , and we have $x^G = x^H \cup (x^g)^H$, where g is any element of $G \setminus H$. In particular, this means that if x^G splits in H and $y = x^g$ for some g in $G \setminus H$, then y is not conjugate to x in H .

Theorem 3.4. *The involutions of $W(\tilde{B}_n)$ are those involutions in $W(\tilde{C}_n)$ with labelled cycle type (m, k_e, k_o, l) where k_o is even. Involution conjugacy classes are parameterised by labelled cycle type, with one conjugacy class for each type except in the case where $k_o = 0$ and $l = 0$. In this case there are two conjugacy classes for each type, and two elements x, x' with the same labelled cycle type $(m, k_e, 0, 0)$ are conjugate if and only if $f(x) \equiv f(x') \pmod{4}$.*

Proof. Let x be an involution in $W(\tilde{C}_n)$ with labelled cycle type (m, k_e, k_o, l) . Transpositions of x are of the form $\begin{smallmatrix} \lambda & -\lambda \\ + & + \end{smallmatrix}$ or $\begin{smallmatrix} \lambda & \lambda \\ - & - \end{smallmatrix}$, each of which contributes an even number (zero or 2λ) to $\sum x$. Therefore $\sum x \equiv k_o \pmod{2}$, and so $x \in W(\tilde{B}_n)$ precisely when k_o is even. Suppose this occurs, and that either $l > 0$ or $k_o > 0$. Then x has a 1-cycle of the form either $\begin{smallmatrix} 0 \\ + \end{smallmatrix}$ or $\begin{smallmatrix} \lambda \\ - \end{smallmatrix}$ for some odd λ . But then $\begin{smallmatrix} \lambda \\ - \end{smallmatrix} \in C_{W(\tilde{C}_n)}(x) \setminus W(\tilde{B}_n)$. Therefore, $x^{W(\tilde{B}_n)} = x^{W(\tilde{C}_n)}$, which in particular means that the conjugacy class of x consists of all involutions with the same labelled cycle type as x . It remains to deal with involutions of labelled cycle type $(m, k_e, 0, 0)$. We first show that there are two such classes in $W(\tilde{B}_n)$. Let $\sigma = (1\ 2) \cdots (2m-1\ 2m)(2m+1) \cdots (2n)$ and $a = (\sigma, \mathbf{0})$. Now, $(g, \mathbf{w}) \in C_{W(\tilde{C}_n)}(a)$ if and only if $(\sigma, \mathbf{0})^{(g, \mathbf{w})} = (\sigma, \mathbf{0})$. Using Equation (1) we have

$$\begin{aligned} (\sigma, \mathbf{0}) &= (\sigma, \mathbf{0})^{(g, \mathbf{w})} = (g^{-1}\sigma g, \mathbf{0} + \mathbf{w} - \mathbf{w}^{g^{-1}\sigma g}) \\ &= (\sigma, (w_1, \dots, w_n) - (w_1, \dots, w_n)^\sigma) \\ &= (\sigma, (w_1 - w_2, w_2 - w_1, \dots, w_{2m-1} - w_{2m}, w_{2m} - w_{2m-1}, 2w_{2m+1}, \dots, 2w_n)). \end{aligned}$$

Hence (g, \mathbf{w}) centralises a if and only if $w_{2i-1} = w_{2i}$ for $1 \leq i \leq m$, and $w_{2m+1} = \cdots = w_n = 0$. Thus, $\sum \mathbf{w}$ is even, which means $C_{W(\tilde{C}_n)}(a) \leq W(\tilde{B}_n)$. This means that there are two conjugacy classes in $W(\tilde{B}_n)$ for each labelled cycle type $(m, k_e, 0, 0)$.

Suppose x and x' have the same labelled cycle type $(m, k_e, 0, 0)$. The action of the underlying $W(B_n)$ permutes coordinates and changes signs, but does not change the modulus of coordinates. For example, a λ over a transposition of x could become $-\lambda$ over a different transposition; a μ over a 1-cycle could become $-\mu$, but since μ is even and the calculation of $f(x)$ involves 2λ rather than λ , the value of $f(x^{(g, \mathbf{0})})$ is congruent modulo 4 to $f(x)$ for any $(g, \mathbf{0}) \in W(\tilde{B}_n)$. Therefore, x is conjugate in $W(\tilde{B}_n)$ to some \bar{x} given

by $\bar{x} = \prod_{i=1}^m \begin{smallmatrix} \lambda_i \\ + \end{smallmatrix} (2i-1\ 2i) \prod_{j=1}^{k_e} \begin{smallmatrix} \mu_j \\ - \end{smallmatrix} (2m+j)$, with $f(\bar{x}) \equiv f(x) \pmod{4}$, and x' is conjugate in $W(\tilde{B}_n)$ to some \bar{x}' given by $\bar{x}' = \prod_{i=1}^m \begin{smallmatrix} \lambda'_i \\ + \end{smallmatrix} (2i-1\ 2i) \prod_{j=1}^{k_e} \begin{smallmatrix} \mu'_j \\ - \end{smallmatrix} (2m+j)$ with $f(\bar{x}') \equiv f(x') \pmod{4}$. It is now sufficient to prove that $\bar{x}' \in \bar{x}^{W(\tilde{B}_n)}$ if and only if $f(\bar{x}') \equiv f(\bar{x}) \pmod{4}$.

Consider the vector $\mathbf{w} = (w_1, \dots, w_n)$ given by $w_{2i-1} = \lambda'_i$, $w_{2i} = \lambda_i$, $w_{2m+j} = \frac{1}{2}(\mu'_j - \mu_j)$ for $1 \leq i \leq m$, $1 \leq j \leq k_e$. Putting $\bar{x} = (\bar{\sigma}, \bar{\mathbf{u}})$, we get

$$\begin{aligned} \bar{x}^{(1, \mathbf{w})} &= (\bar{\sigma}, \bar{\mathbf{u}} + \mathbf{w} - \mathbf{w}^{\bar{\sigma}}) \\ &= (\bar{\sigma}, \bar{\mathbf{u}} + (w_1 - w_2, w_2 - w_1, \dots, w_{2m-1} - w_{2m}, w_{2m} - w_{2m-1}, 2w_{2m+1}, \dots, 2w_n)) \\ &= (\bar{\sigma}, (\lambda_1, -\lambda_1, \lambda_2, -\lambda_2, \dots, \lambda_m, -\lambda_m, \mu_1, \dots, \mu_{k_e}) \\ &\quad + (\lambda'_1 - \lambda_1, \lambda_1 - \lambda'_1, \dots, \lambda_m, -\lambda'_m, \mu'_1 - \mu_1, \dots, \mu'_{k_e} - \mu_{k_e})) \\ &= (\bar{\sigma}, (\lambda'_1, -\lambda'_1, \dots, \lambda'_m, -\lambda'_m, \mu'_1, \dots, \mu'_{k_e})) \\ &= \bar{x}'. \end{aligned}$$

Since there are exactly two conjugacy classes with labelled cycle type $(m, k_e, 0, 0)$, two elements x, x' with this labelled cycle type are conjugate in $W(\tilde{B}_n)$ if and only if any element of $W(\tilde{C}_n)$ which conjugates x to x' is contained in $W(\tilde{B}_n)$.

We have that $\bar{x}' \in \bar{x}^{W(\tilde{B}_n)}$ if and only if $(1, \mathbf{w}) \in W(\tilde{B}_n)$, which holds if and only if $\sum \mathbf{w}$ is even. Now,

$$\begin{aligned} \sum_{i=1}^n w_i &= \sum_{i=1}^m (\lambda_i + \lambda'_i) + \sum_{j=1}^{k_e} \frac{1}{2} (\mu'_j - \mu_j) \\ &\equiv - \left(\sum_{i=1}^m \lambda_i + \frac{1}{2} \sum_{j=1}^{k_e} \mu_j \right) + \left(\sum_{i=1}^m \lambda'_i + \frac{1}{2} \sum_{j=1}^{k_e} \mu'_j \right) \pmod{2} \\ &= \frac{1}{2} (f(\bar{x}') - f(\bar{x})) \pmod{2}. \end{aligned}$$

So $\bar{x}' \in \bar{x}^{W(\tilde{B}_n)}$ if and only if $\frac{1}{2} (f(\bar{x}') - f(\bar{x})) \equiv 0 \pmod{2}$, which holds if and only if $f(\bar{x}') \equiv f(\bar{x}) \pmod{4}$. Thus, $x' \in x^{W(\tilde{B}_n)}$ if and only if $f(x') \equiv f(x) \pmod{4}$. \square

Lemma 3.5. *Let $x \in W(\tilde{C}_n)$ be an involution and let ω be the automorphism of $W(\tilde{C}_n)$ corresponding to the non-trivial automorphism of the Coxeter graph for $W(\tilde{C}_n)$.*

If the labelled cycle form of x contains $(\cdots \overset{\lambda}{\uparrow} i \cdots)$, then $\omega(x)$ contains $(\cdots n+1-i \cdots)$.

If the labelled cycle form of x contains $(\cdots \overset{\lambda}{\downarrow} i \cdots)$, then $\omega(x)$ contains $(\cdots n+1-i \cdots)$. In particular, ω maps labelled cycle types (m, k_e, k_o, l) to labelled cycle types (m, k_o, k_e, l) .

Proof. By definition, $\omega(r_i) = r_{n+2-i}$ for $1 \leq i \leq n+1$. Thus, ω interchanges $\overset{0}{\uparrow}(1)$ and $\overset{1}{\downarrow}(n)$, and maps $\overset{0}{\uparrow}(i \ i+1)$ to $\overset{0}{\uparrow}(n+1-i \ n-i)$. We quickly see that ω also maps any $\overset{0}{\uparrow}(i \ j)$ to $\overset{0}{\uparrow}(n+1-i \ n+1-j)$. The following table builds up the various images of involution cycles σ under ω .

σ	Decomposition	Image of Decomposition	$\omega(\sigma)$
$\overset{0}{\uparrow}(n)$	$\overset{0}{\uparrow}(1 \ n) \overset{0}{\uparrow}(1) \overset{0}{\uparrow}(1 \ n)$	$\overset{0}{\uparrow}(n \ 1) \overset{1}{\downarrow}(n) \overset{0}{\uparrow}(n \ 1)$	$\overset{1}{\downarrow}(1)$
$\overset{\lambda}{\uparrow}(i)$	$\overset{0}{\uparrow}(i \ n) [\overset{0}{\uparrow}(n) \overset{1}{\downarrow}(n)]^\lambda \overset{0}{\uparrow}(i \ n)$	$\overset{0}{\uparrow}(n+1-i \ 1) [\overset{1}{\downarrow}(1) \overset{0}{\uparrow}(1)]^\lambda \overset{0}{\uparrow}(n+1-i \ 1)$	$\overset{-\lambda}{\uparrow}(n+1-i)$
$\overset{\lambda}{\downarrow}(i)$	$\overset{0}{\uparrow}(i \ n) [\overset{0}{\uparrow}(n) \overset{\lambda}{\downarrow}(n)] \overset{0}{\uparrow}(i \ n)$	$\overset{0}{\uparrow}(n+1-i \ 1) [\overset{1}{\downarrow}(1) \overset{-\lambda}{\uparrow}(1)] \overset{0}{\uparrow}(n+1-i \ 1)$	$\overset{1-\lambda}{\downarrow}(n+1-i)$
$\overset{\lambda}{\uparrow}(i \ j)$	$\overset{0}{\uparrow}(i \ j) \overset{\lambda}{\downarrow}(i) \overset{-\lambda}{\uparrow}(j)$	$\overset{0}{\uparrow}(n+1-i \ n+1-j) \overset{-\lambda}{\uparrow}(n+1-i) \overset{\lambda}{\downarrow}(n+1-j)$	$\overset{-\lambda}{\uparrow}(n+1-i \ n+1-j)$
$\overset{\lambda}{\downarrow}(i \ j)$	$\overset{0}{\uparrow}(i \ j) \overset{\lambda}{\downarrow}(i) \overset{\lambda}{\downarrow}(j)$	$\overset{0}{\uparrow}(n+1-i \ n+1-j) \overset{1-\lambda}{\downarrow}(n+1-i) \overset{1-\lambda}{\downarrow}(n+1-j)$	$\overset{1-\lambda}{\downarrow}(n+1-i \ n+1-j)$

The result now follows. \square

An immediate consequence of Lemma 3.5 is that for any involution x of $W(\tilde{C}_n)$, then

as long as we do not rearrange the order in which numbers are written in cycles, we have

$$\begin{aligned}\text{Neg}(\omega(x)) &= \text{Neg}(x); \\ \sum \omega(x) &= \text{Neg}(x) - \sum x; \\ f(\omega(x)) &= \text{Neg}(x) - f(x).\end{aligned}\tag{2}$$

Example 3.6. Recall from Example 2.3 that if

$$x = (1 \overset{+}{2})(3 \overset{-}{4})(5 \overset{+}{6})(7 \overset{-}{8}) = \left((1 \overset{+}{2})(3 \overset{-}{4})(5 \overset{+}{6})(7 \overset{-}{8}), (1, -1, 3, 3, 2, 4, 0) \right),$$

then $\text{Neg}(x) = 4$, $\sum x = 12$, and $f(x) = 14$. We have

$$\omega(x) = (7 \overset{-}{6})(5 \overset{+}{4})(3 \overset{-}{2})(1 \overset{+}{8}) = \left((7 \overset{-}{6})(5 \overset{+}{4})(3 \overset{-}{2})(1 \overset{+}{8}), (0, -3, -1, -2, -2, 1, -1) \right).$$

Note that $\text{Neg}(\omega(x)) = 4$ and

$$\begin{aligned}\sum \omega(x) &= 0 - 3 - 1 - 2 - 2 + 1 - 1 = -8, \\ f(\omega(x)) &= 2(-1 - 2) + (-1 - 3) = -10.\end{aligned}$$

Lemma 3.5 also furnishes a verification of the fact that

$$\overline{W}(\tilde{B}_n) = \{y \in W(\tilde{C}_n) : \text{Neg}(y) \equiv \sum y \pmod{2}\}.$$

To see this, we note that $\overline{W}(\tilde{B}_n) = \omega(W(\tilde{B}_n))$. Thus, every y in $\overline{W}(\tilde{B}_n)$ is $\omega(x)$ for some x such that $\sum x \equiv 0 \pmod{2}$. Therefore,

$$\sum y = \text{Neg}(x) - \sum x = \text{Neg}(y) - \sum x \equiv \text{Neg}(y) \pmod{2}.$$

Theorem 3.7. *The involutions of $\overline{W}(\tilde{B}_n)$ are those in $W(\tilde{C}_n)$ with labelled cycle type (m, k_e, k_o, l) where k_e is even. Involution conjugacy classes are parameterised by labelled cycle type, with one conjugacy class for each labelled cycle type (m, k_e, k_o, l) , except in the case where $k_e = 0$ and $l = 0$. In this case, there are two conjugacy classes for each type, and two elements y and y' with the same labelled cycle type $(m, 0, k_o, 0)$ are conjugate if and only if $f(y) + \text{Neg}(y) \equiv f(y') + \text{Neg}(y') \pmod{4}$.*

Proof. Since $\overline{W}(\tilde{B}_n) = \omega(W(\tilde{B}_n))$, the involutions of $\overline{W}(\tilde{B}_n)$ are the elements $y = \omega(x)$ where x is an involution of $W(\tilde{B}_n)$. By Theorem 3.4, the involutions of $W(\tilde{B}_n)$ are precisely the involutions of $W(\tilde{C}_n)$ with cycle type (m, k_e, k_o, l) where k_o is even. Thus, by Lemma 3.5, the involutions of $\overline{W}(\tilde{B}_n)$ are precisely those involutions of $W(\tilde{C}_n)$ with cycle type (m, k_e, k_o, l) where k_e is even. Next we must determine the conjugacy classes. Since ω maps conjugacy classes to conjugacy classes, Theorem 3.4 and Lemma 3.5 immediately show that involution conjugacy classes in $\overline{W}(\tilde{B}_n)$ are parameterised by labelled cycle type, with one conjugacy class for each labelled cycle type except where $k_e = 0$ and $l = 0$, when there are two classes. Suppose y and y' in $\overline{W}(\tilde{B}_n)$ have the same labelled cycle type $(m, 0, k, 0)$. Then $y = \omega(x)$ and $y' = \omega(x')$, for appropriate $x, x' \in W(\tilde{B}_n)$ of labelled cycle type $(m, k, 0, 0)$. Now, x and x' are conjugate in $W(\tilde{B}_n)$ if and only if $f(x) \equiv f(x') \pmod{4}$. By Equation (2), this holds if and only if

$$\text{Neg}(\omega(x)) - f(\omega(x)) \equiv \text{Neg}(\omega(x')) - f(\omega(x')) \pmod{4},$$

which is equivalent to $f(y) - f(y') \equiv \text{Neg}(y) - \text{Neg}(y') \pmod{4}$. Since y and y' have the same labelled cycle type, $\text{Neg}(y) - \text{Neg}(y')$ is even, meaning that

$$\text{Neg}(y) - \text{Neg}(y') \equiv \text{Neg}(y') - \text{Neg}(y) \pmod{4}.$$

Therefore, y is conjugate to y' if and only if $f(y) - f(y') \equiv \text{Neg}(y') - \text{Neg}(y) \pmod{4}$, which holds if and only if $f(y) + \text{Neg}(y) \equiv f(y') + \text{Neg}(y') \pmod{4}$. \square

Theorem 3.8. *In $W(\tilde{D}_n)$, involution conjugacy classes are parameterised by labelled cycle type, with either one, two or four conjugacy classes for each labelled cycle type. Given an element x of $W(\tilde{D}_n)$ with labelled cycle type (m, k_e, k_o, l) , both k_e and k_o must be even, and either $x^{W(\tilde{D}_n)} = x^{W(\tilde{B}_n)}$, or $x^{W(\tilde{D}_n)} = x^{\overline{W(\tilde{B}_n)}}$ (or both) except in the case where $k_e = k_o = l = 0$ and $n = 2m$. Here, there are four conjugacy classes, and if x and y both have cycle type $(m, 0, 0, 0)$, then x is conjugate to y if and only if both $\text{Neg}(x) \equiv \text{Neg}(y) \pmod{4}$ and $f(x) \equiv f(y) \pmod{4}$.*

Proof. Suppose $x \in W(\tilde{D}_n)$ with labelled cycle type (m, k_e, k_o, l) . If any of k_e , k_o or l is nonzero, then x has a cycle $(\overset{0}{a})$ or $(\overset{\lambda}{a})$ for some λ . But then $(\overset{\lambda}{a})$ is contained either in $C_{W(\tilde{B}_n)}(x) \setminus W(\tilde{D}_n)$ or $C_{\overline{W(\tilde{B}_n)}}(x) \setminus W(\tilde{D}_n)$. Therefore, either $x^{W(\tilde{D}_n)} = x^{W(\tilde{B}_n)}$, or $x^{W(\tilde{D}_n)} = x^{\overline{W(\tilde{B}_n)}}$, or both. The only remaining case is where $m = 2n$. If x and y both have cycle type $(m, 0, 0, 0)$, then in order for them to be conjugate in $W(\tilde{D}_n)$ they must be conjugate both in $W(\tilde{B}_n)$ and $\overline{W(\tilde{B}_n)}$, which means that both $f(x) \equiv f(y) \pmod{4}$ and $\text{Neg}(x) \equiv \text{Neg}(y) \pmod{4}$. Hence there are at least four conjugacy classes of this type in $W(\tilde{D}_n)$, but there are also at most four, because $W(\tilde{D}_n)$ has index 2 in both $W(\tilde{B}_n)$ and $\overline{W(\tilde{B}_n)}$. Hence x is conjugate to y if and only if both $\text{Neg}(x) \equiv \text{Neg}(y) \pmod{4}$ and $f(x) \equiv f(y) \pmod{4}$. \square

4 Connectedness and Diameter Results

We begin with an easy observation about connectedness which allows us to infer properties about commuting involution graphs in affine groups from properties of the corresponding graphs in their finite counterparts. For an element $g = (\sigma, \mathbf{v})$ in a conjugacy class X of \widetilde{W} , we define $\hat{g} = \sigma$ and \hat{X} be the conjugacy class of \hat{g} in W . Clearly if $g, h \in X$, then $\hat{g}, \hat{h} \in \hat{X}$.

Lemma 4.1. *Suppose $g, h \in X$. If $d(\hat{g}, \hat{h}) = k$, then $d(g, h) \geq k$. If $\mathcal{C}(W, \hat{X})$ is disconnected, then $\mathcal{C}(\widetilde{W}, X)$ is disconnected.*

Proof. The result follows immediately from the observation that if g commutes with h in \widetilde{W} , then \hat{g} commutes with \hat{h} in W . \square

The aim of this section is to prove the following result.

Theorem 4.2. *Let \widetilde{W} be one of $W(\tilde{C}_n)$, $W(\tilde{B}_n)$ or $W(\tilde{D}_n)$. Let X be a conjugacy class of \widetilde{W} with labelled cycle type (m, k_e, k_o, l) . Then $\mathcal{C}(\widetilde{W}, X)$ is disconnected in each of the following cases.*

(i) $m = 0$ and $l = 0$;

(ii) $m > 0$, $l = 0$ and either $k_e = 1$ or $k_o = 1$;

(iii) $m > 0$ and $\max\{k_e, k_o, l\} = 1$;

(iv) $n = 4$ and $m = 1$;

(v) $n = 6$, $m = 1$, $k_o = k_e = 2$.

In all other cases, $\mathcal{C}(\widetilde{W}, X)$ is connected with diameter at most $n + 2$.

The next three results from [9] give criteria under which labelled 1-cycles, transpositions and double transpositions commute. They will be used repeatedly in the proofs to follow.

Lemma 4.3 (Lemma 2.7 of [9]). *Let $\alpha \in \{1, \dots, n\}$ and $\lambda, \mu \in \mathbb{Z}$. Then $\overset{\lambda}{\alpha}$ commutes with $\overset{0}{\alpha}$ for all λ , whereas $\overset{\lambda}{\alpha}$ commutes with $\overset{\mu}{\alpha}$ if and only if $\lambda = \mu$.*

Lemma 4.4 (Lemma 2.8 of [9]). *Let α, β be distinct elements of $\{1, \dots, n\}$ and let λ, μ and ν be integers.*

(i) $\overset{\lambda}{\alpha\beta}$ and $\overset{\mu}{\alpha\beta}$ commute if and only if $\lambda = \mu$, and $\overset{\lambda}{\alpha}\beta$ and $\overset{\mu}{\alpha}\beta$ commute if and only if $\lambda = \mu$. But $\overset{\lambda}{\alpha}\beta$ and $\overset{\mu}{\alpha}\beta$ commute for all λ and μ .

(ii) $\overset{\lambda}{\alpha\beta}$ and $\overset{0}{\alpha}\overset{0}{\beta}$ commute for all λ , but there is no value of μ or λ for which $\overset{\lambda}{\alpha}\overset{\mu}{\beta}$ and $\overset{0}{\alpha}\overset{0}{\beta}$ commute, and no value of μ or λ for which $\overset{\lambda}{\alpha}\overset{\mu}{\beta}$ and $\overset{\mu}{\alpha}\overset{0}{\beta}$ commute.

(iii) $\overset{\lambda}{\alpha\beta}$ and $\overset{\mu}{\alpha}\overset{\nu}{\beta}$ commute if and only if $\mu - \nu = 2\lambda$, whereas $\overset{\lambda}{\alpha}\overset{\mu}{\beta}$ and $\overset{\mu}{\alpha}\overset{\nu}{\beta}$ commute if and only if $\mu + \nu = 2\lambda$.

Lemma 4.5 (Lemma 2.9 of [9]). *Let $g_1 = \overset{\lambda_1}{\alpha\beta}\overset{\lambda_2}{\gamma\delta}$, $g_2 = \overset{\lambda_1}{\alpha\beta}\overset{\lambda_2}{\gamma\delta}$, $g_3 = \overset{\lambda_1}{\alpha\beta}\overset{\lambda_2}{\gamma\delta}$, $h_1 = \overset{\mu_1}{\alpha\gamma}\overset{\mu_2}{\beta\delta}$, $h_2 = \overset{\mu_1}{\alpha\gamma}\overset{\mu_2}{\beta\delta}$ and $h_3 = \overset{\mu_1}{\alpha\gamma}\overset{\mu_2}{\beta\delta}$, for distinct $\alpha, \beta, \gamma, \delta$ in $\{1, \dots, n\}$ and integers λ_i, μ_i . Then*

(i) g_1 commutes with h_1 if and only if $\mu_1 - \lambda_1 = \mu_2 - \lambda_2$;

(ii) g_1 does not commute with h_2 ;

(iii) g_1 commutes with h_3 if and only if $\mu_1 - \lambda_1 = \mu_2 + \lambda_2$;

(iv) g_2 commutes with h_2 if and only if $\mu_1 - \lambda_1 = \mu_2 - \lambda_2$;

(v) g_2 does not commute with h_3 ;

(vi) g_3 commutes with h_3 if and only if $\mu_1 - \lambda_1 = \lambda_2 - \mu_2$.

Proposition 4.6. *Let X be a conjugacy class of involutions with labelled cycle type $(m, 0, 0, 0)$ in \widetilde{W} , where \widetilde{W} is either $W(\widetilde{B}_n)$, $\overline{W}(\widetilde{B}_n)$ or $W(\widetilde{D}_n)$.*

(i) *If \widetilde{W} is $W(\widetilde{B}_n)$ or $\overline{W}(\widetilde{B}_n)$, then $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq 3$.*

(ii) If \widetilde{W} is $W(\widetilde{D}_n)$, then $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq 4$.

Proof. Let $x \in X$. Then we have $x = \prod_{i=1}^m (a_i b_i)^{\lambda_i}$ for suitable λ, a_i and b_i .

The two conjugacy classes of labelled cycle type $(m, 0, 0, 0)$ in $W(\widetilde{B}_n)$ are interchanged by conjugation with $(\overline{n})^1$; this induces an isomorphism of their commuting involution graphs. Similarly, the corresponding classes of $\overline{W}(\widetilde{B}_n)$ are mapped to those of $W(\widetilde{B}_n)$ by the action of ω , again inducing isomorphisms between the commuting involution graphs. Therefore, it is sufficient to consider the case where $x \in W(\widetilde{B}_n)$ has labelled cycle type

$(m, 0, 0, 0)$ and $f(x) \equiv 0 \pmod{4}$. That is, $X = a^{W(\widetilde{B}_n)}$, where $a = (1\ 2)^{\overset{0}{+}} (3\ 4)^{\overset{0}{+}} \cdots (n-1\ n)^{\overset{0}{+}}$.

We can write $x = \prod_{i=1}^m (a_i b_i)^{\lambda_i}$, where $\sum_{i=1}^m \lambda_i \equiv 0 \pmod{2}$. Now, let $y = \prod_{i=1}^m (a_i b_i)^{\overset{0}{+}}$. By Lemma 4.4, x commutes with y , and the fact that $f(y) = 0$ implies that $y \in X$. Now, y and a lie in a subgroup isomorphic to $W(B_n)$ and so $d(y, a) \leq 2$ by Theorem 1.4. Hence, $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq 3$.

Now, suppose $\widetilde{W} = W(\widetilde{D}_n)$. In this case, by Theorem 3.8, x is conjugate to exactly one of the following:

$$\begin{aligned} w_1 &= (1\ 2)^{\overset{0}{+}} (3\ 4)^{\overset{0}{+}} \cdots (n-3\ n-2)^{\overset{0}{+}} (n-1\ n)^{\overset{0}{+}} = tr_4 r_6 \cdots r_{n-2} r_n, \\ w_2 &= (1\ 2)^{\overset{0}{+}} (3\ 4)^{\overset{0}{+}} \cdots (n-3\ n-2)^{\overset{0}{+}} (n-1\ n)^{\frac{1}{-}} = tr_4 r_6 \cdots r_{n-2} s, \\ w_3 &= (1\ 2)^{\overset{0}{-}} (3\ 4)^{\overset{0}{+}} \cdots (n-3\ n-2)^{\overset{0}{+}} (n-1\ n)^{\overset{0}{+}} = r_2 r_4 r_6 \cdots r_{n-2} r_n, \\ w_4 &= (1\ 2)^{\overset{0}{-}} (3\ 4)^{\overset{0}{+}} \cdots (n-3\ n-2)^{\overset{0}{+}} (n-1\ n)^{\frac{1}{-}} = r_2 r_4 r_6 \cdots r_{n-2} s. \end{aligned}$$

There are automorphisms of the Coxeter graph of $W(\widetilde{D}_n)$ mapping each w_i to any other w_j . Therefore, the four conjugacy classes have isomorphic commuting involution graphs. Without loss of generality, assume x is conjugate to w_1 . This occurs if and only if $\text{Neg}(x) \equiv 0 \pmod{4}$ and $f(x) \equiv 0 \pmod{4}$, the latter being equivalent to $\sum_{i=1}^m \lambda_i \equiv 0 \pmod{2}$.

If m is even, let $y = \prod_{i=1}^m (a_i b_i)^{\overset{0}{+}}$. We have

$$\text{Neg}(y) \equiv 2m - \text{Neg}(x) \equiv \text{Neg}(x) \equiv 0 \pmod{4}$$

and $f(y) = 0$. Therefore $y \in X$. Now, x commutes with y , and since y lies in a subgroup isomorphic to $W(D_n)$ we have $d(x, w_1) \leq 2$ by Theorem 1.4. Thus, $d(x, w_1) \leq 3$. If m is odd, then as $\sum_{i=1}^m \lambda_i$ is even, there must be at least one even λ_i . Therefore, without

loss of generality, we can suppose λ_1 is even. Now, let $x' = (a_1 b_1)^{\lambda_1} (a_2 b_2)^{\overset{0}{+}} \prod_{i=3}^m (a_i b_i)^{\overset{0}{+}}$ and $y' = (a_1 b_1)^{\overset{0}{+}} (a_2 b_2)^{\overset{0}{+}} \prod_{i=3}^m (a_i b_i)^{\overset{0}{+}}$. We have that

$$\text{Neg}(x') \equiv 2m - 2 - \text{Neg}(x) \equiv \text{Neg}(x) \equiv 0 \pmod{4}$$

and $f(x') = 2\lambda_1 \equiv 0 \pmod{4}$. Also, note that $\text{Neg}(y') = \text{Neg}(x')$ and $f(y') = 0$. Thus, $x', y' \in X$ and x' commutes with both x and y' . Moreover $d(y, w_1) \leq 2$. Therefore $d(x, w_1) \leq 4$. Hence, $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq 4$. \square

In the next proof, and subsequently, we will work inductively on the rank of the group. To do this, we use a shorthand notation for lower rank subgroups. Let $\widetilde{W} = W(\widetilde{C}_n)$. For any subset $\{i_1, \dots, i_t\}$ of $\{1, \dots, n\}$ there is a subgroup of \widetilde{W} isomorphic to $W(\widetilde{C}_t)$, consisting of the labelled signed permutations of $\{i_1, \dots, i_t\}$. We write $\widetilde{W}_{\{i_1, \dots, i_t\}}$ for this subgroup, and \widetilde{W}_t for $\widetilde{W}_{\{1, \dots, t\}}$. We use similar notation for $W(\widetilde{B}_n)$ and $W(\widetilde{D}_n)$.

Proposition 4.7. *Let x be an involution of $\widetilde{W} = W(\widetilde{B}_n)$ with labelled cycle type $(1, k_e, 0, 0)$, where $k_e > 0$. If $k_e \leq 2$, then $\mathcal{C}(\widetilde{W}, X)$ is disconnected. Otherwise,*

$$\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq n + 1.$$

Proof. Let $x \in X$. By Theorem 3.4, x is conjugate either to $w_1 = (1\ 2)(3) \cdots (n)$ or to $w_2 = (1\ 2)(3) \cdots (n)$. Suppose first that x is conjugate to w_1 . Then $f(x) \equiv 2 \pmod{4}$.

If $k_e \leq 2$, then $\mathcal{C}(W(B_n), \hat{X})$ is disconnected, so $\mathcal{C}(\widetilde{W}, X)$ must be as well. So assume $k_e \geq 3$, meaning $n \geq 5$. We will show that if the transposition of x is $(1\ 2)$ for some λ , then $d(x, a) \leq n + 1$. Otherwise $d(x, a) \leq n$.

We proceed by induction on n . The first case to consider is $n = 5$, where we have $x = (a\ b)(c)(d)(e)$ for some integers λ, p, q, r . Since $f(x) \equiv 2 \pmod{4}$, we must have $\lambda + p + q + r \equiv 1 \pmod{2}$. Conjugation by elements of the centraliser of w_1 , which includes the subgroup $\langle (1\ 2), (3\ 4), (4\ 5), (5) \rangle$, does not affect $d(x, w_1)$. Hence we can assume without loss of generality that x contains one of the transpositions $(1\ 5)$, $(3\ 4)$ or $(1\ 2)$. Suppose first that x contains $(1\ 5)$. Then $x = (1\ 5)(2)(3)(4)$ for some integers p, q, r such that $\lambda + p + q + r \equiv 1 \pmod{2}$. Define

$$x_1 = (1\ 5)(2)(3)(4), \quad x_2 = (2\ 3)(1)(4)(5), \quad x_3 = (4\ 5)(1)(2)(3).$$

Note that for each i we have $f(x_i) = f(w_1) = 2$, and hence $x_i \in X$. Moreover, by Lemma 4.4, x, x_1, x_2, x_3, w_1 is a path in $\mathcal{C}(\widetilde{W}, X)$. Hence, $d(x, w_1) \leq 4$.

If x contains $(3\ 4)$, then $x = (3\ 4)(1)(2)(5)$ where again $\lambda + p + q + r \equiv 1 \pmod{2}$, and x commutes with $x' = (1\ 5)(2)(3)(4)$. Now,

$$f(x') = 2(p - r + q + \lambda) \equiv 2(\lambda + p + q + r) = f(x) \pmod{4}.$$

Thus, $x' \in X$, and we have seen above that $d(x', w_1) \leq 4$. Consequently, $d(x, w_1) \leq 5$.

Finally, if $x = (1\ 2)(3)(4)(5)$, then x commutes with $x'' = (3\ 4)(1)(2)(5)$ and $f(x'') \equiv f(x) \pmod{4}$, meaning $x'' \in X$. We have seen above that $d(x'', w_1) \leq 5$. Hence,

$d(x, w_1) \leq 6$. Therefore the inductive hypothesis holds for $n = 5$.

We now assume $n \geq 6$ and proceed inductively. Suppose the transposition of x contains some a with $a > 2$. Then x contains $\overset{\lambda}{*} \overset{\mu}{\underline{a}} \overset{\mu'}{\underline{b}} \overset{\mu'}{\underline{c}} \overset{\mu'}{\underline{d}}$ for some integer λ and even integers μ and μ' . Then, by Lemma 4.4, x commutes with the element y containing $\overset{(\mu+\mu')/2}{\underline{c}} \overset{0}{\underline{a}} \overset{2\lambda}{\underline{b}}$, with all its other cycles the same as x . Now

$$f(y) - f(x) = 2|\lambda| + |\mu + \mu'| - (2|\lambda| + |\mu| + |\mu'|),$$

which is congruent modulo 4 to $f(x)$ because μ and μ' are both even. Thus, $y \in X$.

Ignoring the cycle $\overset{0}{\underline{a}}$, we can work within $\widetilde{W}_{\{1, \dots, a-1, a+1, \dots, n\}}$ to see that inductively $d(y, w_1) \leq n - 1$. Hence $d(x, w_1) \leq n$. If the transposition of x is $\overset{\lambda}{*} (1 \ 2)$ then, again by Lemma 4.4, x certainly commutes with an element of X which does not have this transposition. So $d(x, w_1) \leq n + 1$, as required, and $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq n + 1$.

Finally, the case where x is conjugate to w_2 follows from the w_1 case. This is because the conjugacy class X of w_1 in $W(\widetilde{B}_n)$ is mapped by any element g of $W(\widetilde{C}_n) \setminus W(\widetilde{B}_n)$ to the class Y of w_2 in $W(\widetilde{B}_n)$. This map induces an isomorphism of their respective commuting involution graphs. Therefore the graphs have the same diameters. \square

Proposition 4.8. *Let $\widetilde{W} = W(\widetilde{B}_n)$. If X is a conjugacy class whose elements have labelled cycle type $(m, k_e, 0, 0)$ where both $m > 1$ and $k_e \geq 2$, then $\mathcal{C}(\widetilde{W}, X)$ is connected with diameter at most $n - 1$.*

Proof. There are two conjugacy classes for each labelled cycle type. Let

$$x = \prod_{i=1}^m \overset{\lambda_i}{*} (a_i \ b_i) \prod_{j=1}^{n-2m} \overset{\mu_j}{\underline{c}_j},$$

where the μ_j are all even. Then x is conjugate to exactly one of the following:

$$\begin{aligned} w_1 &= \overset{1}{\underline{1}} \overset{0}{\underline{2}} \overset{0}{\underline{3}} \overset{0}{\underline{4}} \cdots \overset{0}{\underline{(2m-1)}} \overset{0}{\underline{2m}} \overset{0}{\underline{(2m+1)}} \cdots \overset{0}{\underline{n}}; \\ w_2 &= \overset{0}{\underline{1}} \overset{0}{\underline{2}} \overset{0}{\underline{3}} \overset{0}{\underline{4}} \cdots \overset{0}{\underline{(2m-1)}} \overset{0}{\underline{2m}} \overset{0}{\underline{(2m+1)}} \cdots \overset{0}{\underline{n}}. \end{aligned}$$

In particular, x is conjugate to w_2 if and only if $f(x) \equiv 0 \pmod{4}$ which is equivalent to $2 \sum_{i=1}^m \lambda_i + \sum_{j=1}^m \mu_j \equiv 0 \pmod{4}$.

By assumption, $m \geq 1$ and $k_e \geq 2$. We proceed by induction on k_e to show that $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq n - 1$. Suppose $k_e = 2$. We claim that x is distance at most 2 from an element y of X which has the transposition $\overset{0}{\underline{\pm}} (n-1 \ n)$. To see why, note that if both $n-1$ and n appear in transpositions of x , or if both appear in 1-cycles of x , then Lemma 4.4 or Lemma 4.5, as appropriate, implies that x commutes with some x' in X which contains a transposition of the form $\overset{\lambda}{\underline{\mp}} (n-1 \ n)$ for some λ . If $n-1$ and n appear in transpositions of

x , then we note that for each pair of double transpositions that commute in Lemma 4.5, the numbers above the first double transposition are λ_1 and λ_2 , the numbers above the second pair are μ_1 and μ_2 , and, in every case, $\lambda_1 + \lambda_2 \equiv \mu_1 + \mu_2 \pmod{2}$, which means x' is guaranteed to be conjugate to X . If $n - 1$ and n are in 1-cycles of x , then x contains

cycles of the form $(\alpha \beta)(\gamma \delta)(n - 1)(n)$; so, writing $\mu = (\mu_1 - \mu_2)/2$, we may choose x'

to be x with those cycles replaced with $(n - 1 \ n) (\gamma \delta) (\alpha)(\beta)$. In either case, x' is an element of X that commutes with a suitable y . The remaining case is when x contains

(for example) the 1-cycle (n) and $n - 1$ appears in a transposition $(\varepsilon \ n - 1)$ for some

$\varepsilon < n - 1$. Then x commutes with x'' in X containing the transpositions $(\varepsilon \ n - 1)$ and

$(\varepsilon' \ n)$ for some λ, λ' and ε' , where we can choose λ to ensure that $x'' \in X$. Lemma 4.5 now implies that x' commutes with an appropriate y – in particular, one containing the

transpositions $(\varepsilon \ \varepsilon')$ and $(n - 1 \ n)$. We have now shown that, in all cases, x is distance at

most 2 from an element y of X which has the transposition $(n - 1 \ n)$. Hence, the claim

holds. Now, y in turn commutes with some z in X with the 1-cycles $(n - 1)$ and (n) . If

we ignore these cycles and work in \widetilde{W}_{n-2} , then, when $n - 2 = 4$, a quick check confirms that $d(z, a) \leq 2$, whereas when $n - 2 > 4$, Proposition 4.6 tells us that $d(z, a) \leq 3$.

Therefore $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq n - 1$.

Finally, suppose $m \geq 2$ and $k_e > 2$. Suppose there is some transposition of x containing an element α with $\alpha > 2m$. Then by Lemma 4.4(iii) x commutes with some $y \in X$

such that y has the 1-cycle (α) . By induction, $d(y, a) \leq n - 2$. Hence, $d(x, a) \leq n - 1$. The final possibility is that the elements of the transpositions of x are $\{1, 2, \dots, 2m\}$. Since $m > 1$, we can use Lemma 4.5 to show that x commutes with some y in X containing

the transposition $(1 \ 2)$. Working in $\widetilde{W}_{\{3, \dots, n\}}$ (using the case $m = 1$ and induction on m) we see that $d(y, a) \leq n - 2$. Hence $d(x, a) \leq n - 1$, which completes the proof of Proposition 4.8. \square

We note the following.

Theorem 4.9 (Theorem 1.2 of [9]). *Theorem 4.2 holds for $W(\widetilde{C}_n)$.*

We next verify the cases stated to be disconnected in Theorem 4.2.

Proposition 4.10. *Let \widetilde{W} be one of $W(\widetilde{B}_n)$, $\overline{W}(\widetilde{B}_n)$ or $W(\widetilde{D}_n)$. Let X be a conjugacy class of involutions in \widetilde{W} whose elements have labelled cycle type (m, k_e, k_o, l) . Then $\mathcal{C}(\widetilde{W}, X)$ is disconnected in each of the following cases.*

- (i) $m = 0$ and $l = 0$;
- (ii) $m > 0$, $l = 0$ and either $k_e = 1$ or $k_o = 1$ (or both);
- (iii) $m > 0$ and $\max\{k_e, k_o, l\} = 1$;

(iv) $n = 4$ and $m = 1$;

(v) $n = 6$, $m = 1$, $k_o = k_e = 2$.

Proof. (i) If $m = 0$ and $l = 0$, then elements of X consist entirely of negative 1-cycles. So the graph is completely disconnected by Lemma 4.3.

(ii) Suppose $m > 0$, $l = 0$ and either $k_e = 1$ or $k_o = 1$. We cannot have both $k_e = 1$ and $k_o = 1$ because involutions of $W(\tilde{B}_n)$ and $W(\tilde{D}_n)$ require k_o to be even, and involutions of $\overline{W}(\tilde{B}_n)$ and $W(\tilde{D}_n)$ require k_e to be even. Now, Lemma 4.4(ii) guarantees that any x in X having the 1-cycle $\overset{\lambda}{(i)}$ can only commute with elements having the same 1-cycle. So $\mathcal{C}(\widetilde{W}, X)$ is disconnected.

(iii) Suppose $m > 0$ and $\max\{k_e, k_o, l\} = 1$. If $l = 0$, then $\mathcal{C}(\widetilde{W}, X)$ is disconnected, by (ii). If $l > 0$, then $X = X^{W(\tilde{C}_n)}$ by Theorems 3.4, 3.7 and 3.8, and therefore $\mathcal{C}(\widetilde{W}, X)$ is disconnected by Theorem 4.9.

(iv) If $n = 4$ and $m = 1$, then $\mathcal{C}(W, \hat{X})$ is disconnected, so $\mathcal{C}(\widetilde{W}, X)$ is disconnected.

(v) If $n = 6$, $m = 1$, $k_o = k_e = 2$, then $X = X^{W(\tilde{C}_n)}$ by Theorems 3.4, 3.7 and 3.8, and therefore $\mathcal{C}(\widetilde{W}, X)$ is disconnected by Theorem 4.9. \square

Proof of Theorem 4.2 By Theorem 4.9, it only remains to prove Theorem 4.2 for $W(\tilde{B}_n)$ and $W(\tilde{D}_n)$. First, let $\widetilde{W} = W(\tilde{B}_n)$ and let X be a conjugacy class of involutions with labelled cycle type (m, k_e, k_o, l) . By Theorem 3.4, k_o must be even. If $l > 0$ or $k_o > 0$, then $X = X^{W(\tilde{C}_n)}$, and so the results from $W(\tilde{C}_n)$ on connectedness and diameter apply. In particular, in all cases not covered by Proposition 4.10, $\mathcal{C}(\widetilde{W}, X)$ is connected with diameter at most $n + 2$. It remains to deal with the case where $l = 0$ and $k_o = 0$. Here, if $k_e = 0$, then $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq 3$ by Lemma 4.6. If $k_e = 1$, then $\mathcal{C}(\widetilde{W}, X)$ is disconnected by Proposition 4.10. So suppose $k_e \geq 2$. If $m = 0$, then $\mathcal{C}(\widetilde{W}, X)$ is completely disconnected. If $m = 1$, then by Proposition 4.7 either $k_e \leq 2$ and $\mathcal{C}(\widetilde{W}, X)$ is disconnected, or $k_e \geq 2$ and $\text{Diam } \mathcal{C}(\widetilde{W}, X) \leq n + 1$. If $m > 1$, then Proposition 4.8 guarantees that $\mathcal{C}(\widetilde{W}, X)$ is connected with diameter at most $n - 1$. Therefore, Theorem 4.2 holds for $W(\tilde{B}_n)$. Since ω maps classes of $W(\tilde{B}_n)$ to classes of $\overline{W}(\tilde{B}_n)$, it follows immediately that Theorem 4.2 also holds for $\overline{W}(\tilde{B}_n)$. Now suppose $\widetilde{W} = W(\tilde{D}_n)$. By Theorem 3.8, if any of k_o, k_e or l are nonzero, then either $X = X^{W(\tilde{B}_n)}$, or $X = X^{\overline{W}(\tilde{B}_n)}$, or both. Therefore, the results from $W(\tilde{B}_n)$ and $\overline{W}(\tilde{B}_n)$ apply. The only remaining case is when X has labelled cycle type $(m, 0, 0, 0)$. Here, $\mathcal{C}(\widetilde{W}, X)$ is connected with diameter at most 4, by Proposition 4.6. Therefore, Theorem 4.2 also holds for $W(\tilde{D}_n)$. \square

5 Concluding Remarks

The results proved in this paper, together with the results of [2], mean that bounds on the diameters of connected commuting involution graphs have been obtained for all finite Coxeter groups, and all classical affine Weyl groups. Specifically, Theorem 1.1 states that if W is a classical Weyl group of rank n , and X a conjugacy class of involutions of its corresponding affine Weyl group \tilde{W} , then either $\mathcal{C}(\widetilde{W}, X)$ is disconnected, or it

is connected with diameter at most $n + 2$. In addition, Theorem 4.2 gives necessary and sufficient criteria for the connectedness of $\mathcal{C}(\widetilde{W}, X)$. A natural question is whether the $n + 2$ bound continues to hold for the exceptional affine groups. There are some preliminary results in [10] and [16]. In particular, complete results are known for types \widetilde{F}_4 and \widetilde{G}_2 , along with results for some classes in \widetilde{E}_6 , \widetilde{E}_7 and \widetilde{E}_8 . Full information for types \widetilde{E}_6 , \widetilde{E}_7 and \widetilde{E}_8 requires more work and is the subject of a forthcoming paper by the current authors.

References

- [1] C. Bates, D. Bundy, P. Rowley and S. Perkins. *Commuting involution graphs for symmetric groups*, J. Algebra, 266(1) (2003), 133–153.
- [2] C. Bates, D. Bundy, P. Rowley and S. Perkins. *Commuting Involution Graphs for Finite Coxeter Groups*, J. Group Theory 6 (2003), 461–476.
- [3] C. Bates, D. Bundy, P. Rowley and S. Perkins. *Commuting involution graphs in special linear groups*. Comm. in Algebra, 32(11) (2004), 4179–4196.
- [4] C. Bates, D. Bundy, P. Rowley and S. Hart. *Commuting involution graphs for sporadic simple groups*. J. Algebra, 316(2) (2007), 849–868.
- [5] R. Brauer, K.A. Fowler. *On groups of even order*, Ann. of Math. (2) 62 (1955), 565–583.
- [6] R. W. Carter. *Conjugacy classes in the Weyl group*, Compositio Math. 25 (1972), 1–59.
- [7] B. Fischer. *Finite groups generated by 3-transpositions*, I. Invent. Math. 13 (1971), 232–246.
- [8] M. Giudici and C. Parker. *There is no upper bound for the diameter of the commuting graph of a finite group*, J. Combin. Theory Ser. A 120 (2013), no. 7, 1600–1603.
- [9] S. Hart and A. Sbeiti Clarke. *Commuting Involution Graphs for \widetilde{C}_n* , Comm. in Algebra 46 (9) (2018), 3965–3985.
- [10] S. Hart and A. Sbeiti Clarke. *A Note on Commuting Involution Graphs in Affine Coxeter Groups*, Birkbeck Mathematical Sciences Preprint Series No. 43 (2018). <http://www.bbk.ac.uk/ems/research/pure/preprints>
- [11] J.E. Humphreys. *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, 29 (1990).
- [12] A. Iranmanesh and A. Jafarzadeh. *On the commuting graph associated with the symmetric and alternating groups*, J. Algebra Appl. 7 (2008) 129–146.
- [13] G. L. Morgan and C. W. Parker. *The diameter of the commuting graph of a finite group with trivial centre*, J. Algebra 393 (2013), 41–59.
- [14] S. Perkins. *Commuting Involution Graphs in the affine Weyl group \widetilde{A}_n* , Arch. Math. 86 (2006), no. 1, 16–25.

- [15] R. W. Richardson. *Conjugacy Classes of Involutions in Coxeter Groups*, Bull. Austral. Math. Soc. 26 (1982), 1–15.
- [16] A. Sbeiti Clarke. *Affine Coxeter Groups, Conjugacy Classes and Commuting Involution Graphs*, Ph.D. Thesis, Birkbeck (University of London), 2018.