




## BIROn - Birkbeck Institutional Research Online

Hart, Sarah and Anabanti, Chimere (2020) A question of Mazurov on groups of exponent dividing 12. *Communications in Algebra* 48 (12), pp. 5372-5373. ISSN 0092-7872.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/40588/>

*Usage Guidelines:*

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html> or alternatively contact [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk).



ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/lagb20>

## A question of Mazurov on groups of exponent dividing 12

Chimere Stanley Anabanti , Sarah Beatrice Hart & Michael C. Slattery

To cite this article: Chimere Stanley Anabanti , Sarah Beatrice Hart & Michael C. Slattery (2020): A question of Mazurov on groups of exponent dividing 12, Communications in Algebra, DOI: 10.1080/00927872.2020.1788569

To link to this article: <https://doi.org/10.1080/00927872.2020.1788569>



© 2020 Austrian Science Fund (FWF).  
Published with license by Taylor & Francis  
Group, LLC.



Published online: 08 Jul 2020.



Submit your article to this journal [↗](#)



Article views: 156



View related articles [↗](#)



View Crossmark data [↗](#)

## A question of Mazurov on groups of exponent dividing 12

Chimere Stanley Anabanti<sup>a\*</sup> , Sarah Beatrice Hart<sup>b</sup> , and Michael C. Slattery<sup>c</sup> 

<sup>a</sup>Institut für Analysis und Zahlentheorie, Technische Universität Graz (TU Graz), Graz, Austria; <sup>b</sup>Department of Economics, Mathematics and Statistics, Birkbeck, University of London, London, UK; <sup>c</sup>Department of Mathematical and Statistical Sciences, Marquette University, Wisconsin, USA

### ABSTRACT

Mazurov asked whether a group of exponent dividing 12, which is generated by  $x$ ,  $y$  and  $z$  subject to the relations  $x^3 = y^2 = z^2 = (xy)^3 = (yz)^3 = 1$ , has order at most 12. We show that if such a group is finite, then the answer is yes.

### ARTICLE HISTORY

Received 27 December 2019  
Revised 12 June 2020  
Communicated by Mark L. Lewis

### KEYWORDS

Exponent; finite presentation; groups

### 2010 MATHEMATICS

**SUBJECT CLASSIFICATION**  
20F05; 20D60

The following question of Mazurov is listed as Question 19.53 in the collection of open problems in the Kourovka Notebook [2].

**Question 1 (Mazurov).** *Let  $G$  be a group of exponent 12 generated by elements  $x$ ,  $y$ ,  $z$  such that  $x^3 = y^2 = z^2 = (xy)^3 = (yz)^3 = 1$ . Is it true that  $|G| \leq 12$ ?*

Recall that the **exponent** of a group  $G$  is the smallest positive integer  $n$  such that  $g^n = 1$  for all  $g \in G$ ; meanwhile,  $G$  has **period**  $n$  whenever the exponent of  $G$  divides  $n$ . In fact, the question as stated in [2] requires “exponent 12” rather than “exponent dividing 12”, but Mazurov has confirmed to the authors that “exponent dividing 12” (that is, period 12) was intended. If the answer to the question is yes, then one consequence would be that groups of period 12 are locally finite (see [3]).

As a step in this direction, we have the following. Here,  $C_3$  is the cyclic group of order 3, and  $A_4$  is the alternating group of degree 4.

**Lemma 2.** *Let  $G$  be a group of exponent dividing 12, which is generated by  $x$ ,  $y$  and  $z$  subject to the relations  $x^3 = y^2 = z^2 = (xy)^3 = (yz)^3 = 1$ . If  $G$  is finite, then  $G$  is either trivial or isomorphic to either  $C_3$  or  $A_4$ .*

**Proof.** Let  $G$  be a group of exponent dividing 12 with the given presentation. Then  $G$  certainly satisfies the additional relations  $(xz)^{12} = 1$  and  $(xyz)^{12} = 1$ . Therefore  $G$  is a quotient of the

**CONTACT** Chimere S. Anabanti  [anabanti@math.tugraz.at](mailto:anabanti@math.tugraz.at)  Institut für Analysis und Zahlentheorie, Technische Universität Graz (TU Graz), Graz 8010 Austria.

\*He is also at the Department of Mathematics, University of Nigeria, Nsukka (UNN), and he uses  [chimere.anabanti@unn.edu.ng](mailto:chimere.anabanti@unn.edu.ng).

This article has been republished with minor changes. These changes do not impact the academic content of the article.

© 2020 Austrian Science Fund (FWF). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

group  $U$  given by

$$U := \langle x, y, z \mid x^3 = y^2 = z^2 = (xy)^3 = (yz)^3 = (xz)^{12} = (xyz)^{12} = 1 \rangle.$$

We observe that a finite group  $G$  of exponent dividing 12 must have order  $2^a 3^b$  for some  $a$  and  $b$ . Therefore, by Burnside's Theorem,  $G$  is solvable. Hence,  $G$  is a solvable quotient of  $U$ . We may therefore employ the command

```
SolvableQuotient(U);
```

in MAGMA [1]. This function returns the largest solvable quotient of a given finitely presented group. The outcome is as follows.

```
>U:= Group<x,y,z|x^3, y^2, z^2, (x*y)^3, (y*z)^3, (x*z)^12, (x*y*z)^12>;
>SolvableQuotient(U);
GrpPC of order 12 = 2^2 * 3
PC-Relations:
$.1^3 = Id($),
$.2^2 = Id($),
$.3^2 = Id($),
$.2$.1 = $.3,
$.3$.1 = $.2 * $.3
```

Therefore,  $|G| \leq 12$ . It is now quick to check by hand that the only possibilities for  $G$ , apart from the trivial group, are  $C_3$  and  $A_4$ .

We note that, in terms of the original Question 1 above, Lemma 2 shows that if a group of exponent exactly 12 with the given relations exists, then it must be infinite.

## Funding

Chimere S. Anabanti is supported by the Austrian Science Fund (FWF): P30934-N35.

## ORCID

Chimere Stanley Anabanti  <http://orcid.org/0000-0003-4564-4179>

Sarah Beatrice Hart  <http://orcid.org/0000-0003-3612-0736>

Michael C. Slattery  <http://orcid.org/0000-0001-8178-3534>

## References

- [1] Bosma, W., Cannon, J., Playoust, C. (2019). The Magma algebra system, Version 2.24–5. <http://magma.maths.usyd.edu.au/calc/>.
- [2] Khukhro, E. I., Mazurov, V. D. Unsolved problems in group theory. The Kourovka Notebook. No 19. arXiv:1401.0300v17. <https://arxiv.org/pdf/1401.0300.pdf>.
- [3] Lytkina, D. V., Mazurov, V. D. (2015). On groups of period 12. *Sib. Math. J.* 56(3):471–475. DOI: 10.1134/S0037446615030106.