



BIROn - Birkbeck Institutional Research Online

Longo, Matthew R. and Betti, V. and Aglioti, S.M. and Haggard, P. (2009) Visually induced analgesia: seeing the body reduces pain. *Journal of Neuroscience* 29 (39), pp. 12125-12130. ISSN 0270-6474.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/5409/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html> or alternatively contact lib-eprints@bbk.ac.uk.



BIROn - Birkbeck Institutional Research Online

Enabling open access to Birkbeck's published research output

Visually induced analgesia: seeing the body reduces pain

Journal Article

<http://eprints.bbk.ac.uk/5409>

Version: Accepted (Refereed)

Citation:

Longo, M.R., Betti, V., Aglioti, S.M. and Haggard, P. (2009)
Visually induced analgesia: seeing the body reduces pain –
Journal of Neuroscience 29(39)

© 2009 Society for Neuroscience

[Publisher version](#)

All articles available through Birkbeck ePrints are protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

[Deposit Guide](#)

Contact: lib-eprints@bbk.ac.uk

Section and Senior Editor: Behavioral/Systems/Cognitive Neuroscience

Title: Visually-Induced Analgesia: Seeing the Body Reduces Pain

Abbreviated title: Pain and the Body

Authors: Matthew R. Longo¹, Viviana Betti^{2,3}, Salvatore M. Aglioti^{2,3}, Patrick Haggard¹

¹Institute of Cognitive Neuroscience, University College London, United Kingdom

²Dipartimento di Psicologia, Università degli studi di Roma 'La Sapienza', Rome, Italy

³IRCCS Fondazione Santa Lucia, Rome, Italy

Corresponding author: Matthew R. Longo, Email: m.longo@ucl.ac.uk

Institute of Cognitive Neuroscience, University College London

17 Queen Square, London, WC1N 3AR, UK

Number of figures and tables: 5

Supplemental material (if applicable): YES

Number of pages: 23

Number of words for Abstract/Introduction/Discussion/Total: 150/500/1,184/5,316

Keywords: Pain, Laser-evoked potentials, Body Representation, Analgesia, Mirror Box, Multisensory

Acknowledgements: This work was supported by Biotechnology and Biological Sciences Research Council (BBSRC) [grant number BB/D009529/1]; a BBSRC ISIS collaboration grant awarded to PH and SMA, a Leverhulme Trust Research Fellowship to PH; a grant by the MIUR and Ministero della Sanità, Italy to SMA; and by a Short-Term Scientific Mission of European Science Foundation COST action BM0605 to VB. We wish to thank Dr Laura Cerone for her help with data collection.

Abstract

Given previous reports of strong interactions between vision and somatic senses, we investigated whether vision of the body modulates pain perception. Participants looked into a mirror aligned with their body midline either at the reflection of their own left hand (creating the illusion that they were looking directly at their own right hand), or at the reflection of a neutral object. We induced pain using an infrared laser and recorded nociceptive laser-evoked potentials (LEPs). We also collected subjective ratings of pain intensity and unpleasantness. Vision of the body produced clear analgesic effects both on subjective ratings of pain and on the N2/P2 complex of LEPs. Similar results were found during direct vision of the hand, without the mirror. Further, these effects were specific to vision of one's own hand, and were absent when viewing another person's hand. These results demonstrate a novel analgesic effect of non-informative vision of the body.

Introduction

Several sensory systems carry information about the body to the brain; the body is a multisensory object, par excellence. Most previous multisensory research has focused on fusion of *events* across modalities (Ernst and Banks, 2002) or on how events in one modality bias other modalities (Jousmäki and Hari, 1998). A less-studied type of interaction involves a continuous *context* in one modality shaping perception of events in another modality. One such context is simply viewing one's own body, which increases tactile spatial acuity (Kennett et al., 2001), accelerates tactile reactions (Tipper et al., 1998), and modulates somatosensory evoked-potentials (SEPs) (Taylor-Clarke et al., 2002). These effects require the visual content of one's own body, but not any visual event. The generality of such effects, however, is unknown, having been investigated exclusively for touch. Here, we investigated the effects of seeing the body on the perception and neural processing of pain.

How might seeing the body affect pain? Contextual modulation has long been understood as an organizing principle of neural pain pathways (cf. Melzack and Wall, 1965). However, modulation from seeing one's body could involve either enhancement or inhibition. If enhancing effects observed previously for touch apply across somatic modalities, increases in perceived pain and laser-evoked potentials (LEPs) might be expected. However, many intermodal interactions involving pain are inhibitory. Touch reduces both pain levels (Wall and Sweet, 1967; Higgins et al., 1971) and cortical pain processing (Lundeberg, 1985; Inui et al., 2006). Chronic pain is associated with reduced tactile sensitivity on the affected region (Moriwaki and Yuge, 1999), and reduced size of the corresponding SI tactile representation (Maihöfner et al., 2003; Pleger et al., 2006). Conversely, tactile discrimination training reduces chronic central pain (Flor et al., 2001; Moseley et al., 2008b), an

effect enhanced by seeing the body (Moseley and Wiech, 2009). In each case, touch and pain appear antagonistic, suggesting that vision of the body might have different effects on touch and pain. Furthermore, seeing *someone else's* hand in pain reduces early cortical pain processing (Valeriani et al., 2008). However, the effects of seeing one's own body on nociception are unknown.

We investigated how vision of one's hand affects perception of infrared laser-induced pain and associated cortical processing. Laser stimulation selectively activates thin nociceptive A δ and C-fibers without activating mechanoreceptive afferents (Treede et al., 2003), creating a pure pain sensation, without touch. We used the mirror box technique (Ramachandran et al., 1995) to create the illusion that the participant's left hand reflected in a mirror aligned with their sagittal plane was actually a direct view of their stimulated right hand. In a control condition participants saw the mirror image of a non-hand object. The mirror box provides an elegant experimental means of manipulating vision of the body while keeping vision non-informative about stimulation, and has been used for this reason before (Harris et al., 2007; Longo et al., 2008a; Moseley & Wiech, 2009). We measured subjective ratings of pain intensity and unpleasantness, using a visual-analogue scale (VAS) and cortical processing using LEPs.

Materials and Methods

Three experiments were conducted. Experiment 1 used the mirror box to investigate the effects of non-informative vision of the body on pain perception. Participants in Experiment 1 experienced pain on their hand, but did not see any visual stimulus appearing to cause it, potentially generating perceptual conflict. Experiment 2 addressed whether this conflict was responsible for reduced pain. Now

participants looked directly at either their hand or the object. The laser stimulator wand, and the red visual laser spot on the hand during actual stimulation, was now visible in the view hand condition, eliminating any intersensory conflict. Experiment 3 investigated whether visually-induced analgesia was specific to viewing one's own hand, or would be similarly elicited by viewing another person's hand.

Participants. Thirty healthy volunteers (18 female) between 18 and 34 years old ($M = 24.6$ years; $SD = 3.7$ years) participated ($n = 14$ for Experiment 1; $n = 16$ for Experiment 2, 12 of whom also participated in Experiment 3) for payment. Participants were predominantly right-handed (Edinburgh Inventory, $M = 74.3$, $SD = 33.1$). Procedures were approved by the local ethics committee.

Procedure. In Experiment 1, participants looked into a mirror aligned with their parasagittal plane. Their right hand was behind the mirror (index finger 20 cm from mirror). A baffle extending above the mirror and a black smock worn by the participant prevented vision of the stimulated right hand and arm. In the 'view hand' condition, the participant's left hand was placed in front of the mirror (index finger 20 cm from mirror). Thus, participants saw what appeared to be their right hand (but was in fact the reflected left hand), at the location where their right hand felt to be (see Figure 1). In the 'view object' condition, a small brown book (12.50 x 18.50 x 2.40 cm) was placed in approximately the same position, and the non-stimulated left hand rested on the left leg below the table. In both conditions, participants were instructed to gaze into the mirror and fixate the hand/object. The laser stimulation was applied to the right hand behind the mirror.

There were four experimental blocks, two of each condition, with ABBA counterbalancing. The first condition was counterbalanced across participants. Each block began with a 60 second induction period of passive looking at the hand or

object. Then 30 laser pulses were applied to the dorsum of the right hand. To avoid nociceptor sensitization, the laser was moved randomly between pulses. At least seven seconds elapsed between pulses. Participants rated pain intensity and unpleasantness after each pulse, using a 101-point visual analogue scale (VAS) where 0 corresponded to no pain, and 100 to the worst pain imaginable.

A questionnaire was administered after each block. The items (translated from Italian) were: (1) “It felt like I was looking directly at my hand, rather than at a mirror image.”; (2) “It felt like the hand I was looking at was *my* hand.”; (3) Did it seem like the hand you saw was a right hand or a left hand?”. Item 1 was given after both conditions, items 2 and 3 only in the hand condition. For items 1 and 2, participants rated their agreement using a 7-point Likert scale, +3 indicating “strongly agree”, -3 “strongly disagree”, and 0 “neither agree nor disagree,” though any intermediate value could be used. Values above 0 indicate agreement, while values below 0 indicate disagreement. Agreement or disagreement was tested by comparing the mean score to 0 using t-tests. Item 3 required dichotomous responses, after which participants indicated the strength of the feeling that the hand was a right/left hand using the VAS. Right hand responses were coded positively, left hand responses negatively, yielding scores between -100 (strong left hand) to +100 (strong right hand).

Experiment 2 was similar to Experiment 1 except that the mirror was removed and participants looked either directly at their stimulated right hand or at the object. In the object condition, the object was placed next to the hand and a cardboard baffle was placed between the hand and the object, occluding direct view of the hand. As the laser stimulator projected a red laser dot onto the hand, a laser pointer was held above the object, projecting a similar red laser light. In the hand condition, the object was removed, and the baffle was moved behind the hand. The questionnaire regarding

visual experience was omitted. To reduce habituation effects, Experiments 2 and 3 were preceded by 10 laser pulses.

Experiment 3 was similar to Experiment 1 with the addition of a condition in which participants saw the mirror reflection of an experimenter's hand. Thus, participants viewed the mirror reflection of (a) their own hand, (b) the object, or (c) somebody else's hand (an experimenter's). The experimenter was seated next to the participant with their left hand positioned in the location that the subject's hand had occupied in experiment 1. Thus, participants gazing into the mirror had the visual experience of another person's right hand appearing in first-person perspective at the location where their own right hand felt to be. The participant's left hand remained in their lap. No attempt was made to disguise the presence of the experimenter, or to make their hand physically resemble the participant's hand. There was a single block of each condition, the order of which was counterbalanced across participants.

Pain stimuli were delivered with an infrared neodymium yttrium aluminium perovskite (Nd:Yap) laser (EL. EN. Group, Florence, Italy. Pulse duration and intensity were adjusted for each participant to the minimum values eliciting clear sensations of pain (laser pulses: intensity 1.5-3.25 J; duration 2-4 ms; diameter 4-5 mm, wavelength of 1.34 μm).

EEG recording. A SynAmp amplifiers system and Scan 4.3 software (Neuroscan, El Paso, TX) were used to record electroencephalographic (EEG) data. Recordings were obtained from 54 scalp electrodes in Experiment 1 and 60 in Experiments 2 and 3. Horizontal electrooculogram was recorded bipolarly from electrodes placed on the outer canthi of each eye, and vertical electrooculogram was recorded from an electrode below the right eye. Reference was at the nose and

ground at AFz. Electrode impedances were kept below 5 K Ω . EEG signals were amplified and digitized at 1000 Hz.

Data analysis. EEG data were analyzed with EEGLAB (Delorme and Makeig 2004) in Matlab 7.3 (Mathworks, Natick, MA). Data were digitally filtered with a bandpass of 0.3 – 30 Hz and segmented into epochs timelocked to laser pulses (-500 to 1000 ms). Baseline was calculated from the 200 ms preceding the pulse. Visual inspection of epoched data was used to remove trials with obvious contamination by ocular movements. Further correction of ocular artifacts was performed using blind source separation with independent components analysis (Jung et al., 2000) on epoched data. To reduce effects of pain habituation (Ernst et al., 1986; Milne et al 1991; Valeriani et al 2003) the first 10 trials of the experiment were eliminated and counterbalance order was included as a between-subjects factor in ANOVAs to exclude variance attributable to habituation (see Supplemental Materials).

Two LEP components were investigated. First, the early, lateralized N1 potential, maximal over contralateral temporal electrodes, and originating from operculoinsular cortex (Garcia-Larrea et al., 2003), possibly including SII (Frot et al 1999; Spiegel et al 1996). Second, the bipolar vertex N2/P2 complex. Intracranial recordings have linked the N2/P2 to parallel activations in at least three brain areas: SI, parasyllian, and cingulate cortices (Ohara et al., 2004). N1 peaks were computed as the minimum value at T7 re-referenced to Fz between 130 and 210 ms. N2 and P2 peaks were computed, respectively, as the minimum value at Cz between 170 and 240 ms and the maximum value at Cz between 300 and 450 ms. N2/P2 amplitude was computed as the difference between P2 and N2 peaks.

Results

Experiment 1

Illusion questionnaire. Participants agreed that “it felt like I was looking directly at my hand, rather than at a mirror image” after seeing their hand (i.e., the mean score was significantly greater than 0), $t(13) = 3.26, p < .01$, but disagreed after seeing the object, $t(13) = -37.39, p < .0001$ (see Figure 3). In the hand condition, participants agreed that “It felt like the hand I was looking at was mine”, $t(13) = 11.59, p < .0001$, and felt like they were looking at a right (rather than a left) hand, $t(13) = 3.21, p < .01$. Thus, the mirror box effectively created the illusion of looking directly at the right hand in the hand – but not the object – condition.

Subjective ratings of pain. Seeing the hand reduced pain intensity, $F(1, 12) = 8.07, p < .02$ (see Figure 4a), and unpleasantness, $F(1, 12) = 5.58, p < .05$. These reductions were strongly correlated across individuals, $r(13) = .92, p < .0001$.

Laser-evoked potentials. Seeing the hand also reduced N2/P2 amplitude, $F(1, 12) = 28.83, p < .001$ (see Figure 5a). No significant modulation of the N1 component was observed (hand: $M = -7.4 \mu\text{V}, SD = 4.3$; object: $M = -6.7 \mu\text{V}, SD = 2.5$), $F(1, 12) = .88$.

There were significant correlations between the analgesic effects of seeing the hand on N2/P2 amplitude and subjective reports of both pain intensity, $r(13) = .72, p < .005$, and unpleasantness, $r(13) = .61, p < .02$.

Experiment 2

Subjective ratings of pain¹. Seeing the hand significantly reduced pain intensity, $F(1, 14) = 9.60, p < .01$, and unpleasantness, $F(1, 14) = 7.56, p < .02$ (see Figure 4b). The two measures were again strongly correlated across participants, $r(15) = .73, p < .001$.

Laser-evoked potentials. Seeing the hand again significantly reduced N2/P2 amplitude, $F(1, 14) = 5.82, p < .05$ (see Figure 5b). No significant modulation of the N1 was observed (hand: $M = -9.6 \mu\text{V}, SD = 4.8$; object: $M = -9.3 \mu\text{V}, SD = 4.6$), $F(1, 14) = .21$.

Correlations between the analgesic effects on subjective ratings of pain and N2/P2 amplitude were in the same direction as in Experiment 1, but did not reach statistical significance for either pain intensity, $r(15) = .32, p = .21$, or unpleasantness, $r(15) = .46, p = .06$.

Experiment 3

One participant showed highly unstable subjective pain ratings, with a progressive escalation of reported pain intensity from 17 in block 1 to 72 in block 3 and was excluded from analyses.

Illusion questionnaire. Participants on average agreed that “it felt like I was looking directly at my hand, rather than at a mirror image” after seeing their hand, $t(10) = 5.19, p < .0005$, but not the object, $t(10) = -23.12, p < .0001$, or the experimenter’s hand, $t(10) = .94, ns$ (see Figure 3). In the own hand condition, participants agreed that “It felt like the hand I was looking at was mine”, $t(10) = 9.93, p < .0001$, but not when seeing someone else’s hand, $t(10) = .90, ns$. Participants reported that it felt like they were looking at a right (rather than a left) hand both when seeing their own hand, $t(10) = 3.63, p < .005$, and someone else’s hand, $t(10) = 2.17, p = .055$. Participants disagreed that “it felt like I was looking at somebody else’s hand” when looking at their own hand, $t(10) = -6.50, p < .0001$, but not when seeing the other person’s hand, $t(10) = -.82, ns$. Therefore, the mirror box produced

the illusion of looking at one's own right hand in the 'view hand' condition, but not when viewing the experimenter's hand.

Subjective ratings of pain. There was a significant main effect of condition both on pain intensity, $F(2, 20) = 5.00, p < .05$, and unpleasantness, $F(2, 20) = 4.79, p < .05$ (see Figure 4c). Seeing one's own hand was analgesic relative to seeing the object, both for pain intensity, $t(10) = -3.07, p < .02$, and unpleasantness, $t(10) = -2.70, p < .05$. Similar reductions were also observed when seeing one's own hand relative to seeing someone else's hand: intensity, $t(10) = -2.53, p < .05$; unpleasantness, $t(10) = -2.55, p < .05$. The object and other hand conditions did not differ significantly, either for intensity, $t(10) = .83$, or unpleasantness, $t(10) = 1.34$. These results suggest that the analgesic effects of seeing a hand are specific to viewing one's own hand.

Laser-evoked potentials. There was a significant effect of condition on N2/P2 amplitude, $F(2, 20) = 4.74, p < .05$ (see Figure 5c). Amplitude was reduced when seeing one's own hand compared to the object, $t(10) = -2.63, p < .05$, and someone else's hand, $t(10) = -2.62, p < .05$. There was no difference between the object and other hand conditions, $t(10) = .05$. There was no effect of condition on N1 amplitude (own hand: $M = -7.31 \mu\text{V}, SD = 4.4$; other hand: $M = -8.0 \mu\text{V}, SD = 3.5$; object: $M = -6.95 \mu\text{V}, SD = 3.8$), $F(2, 10) = .48$.

Discussion

These results demonstrate a novel form of visually-induced analgesia. Looking at one's hand produced significant reductions of subjective intensity and unpleasantness of laser pain, and in the amplitude of the N2/P2 complex of LEPs. These effects were observed both when the illusion of looking directly at the hand was induced with a mirror box (Experiments 1 and 3) as well as when participants

viewed their hand directly during stimulation (Experiment 2). The reduction was specific to seeing one's own hand (Experiment 3). Previous studies have suggested that vision of the body may reduce chronic phantom limb pain (Ramachandran and Rogers-Ramachandran, 1996; Chan et al., 2007), and also that seeing someone else's body in pain induces a specific reduction of early LEPs (Valeriani et al., 2008). The present data show an analgesic effect of seeing the body, not tied to the specific case of phantom limbs, and operating on acute stimulus-generated pain. Thus, we show that viewing the body not only induces plasticity in the cortical representations within which pain signals are processed, but also modulates perceptual processing of individual pain events.

Valeriani and colleagues (2008) recently found that empathy for others' pain attenuates the early, lateralized N1/P1 complex of LEPs, without affecting subjective pain ratings. In contrast, the present results demonstrate that seeing one's own hand attenuates the later, vertex N2/P2 complex *with* corresponding subjective attenuation. This divergence between early and late LEPs suggests that our visual effect and their empathic effect may have qualitatively different neuronal bases. The absence of modulation of the N1 in the present study raises the possibility that the effects of seeing one's hand operate via modulation of evaluative and affective, rather than purely sensory, components of the pain matrix. However, sensory and affective components of pain tend to be strongly correlated, so the relation between Valeriani et al's result and ours remains unclear.

Our findings suggest a reliable attenuation of pain systems for viewing the body. These findings suggest several possible theoretical interpretations. We first considered the theory that conflict between visual and proprioceptive representations could induce visual analgesia. However, Experiment 2 showed similar results when

conflict-inducing mirrors were not used, contrary to this theory. An alternative theory links the analgesic effect to a sense of body-ownership. The experience of viewing one's own body contains multiple dissociable elements (Longo et al., 2008b), including the senses of ownership (i.e., that it is *my* body) and agency (i.e., that I am in control of my body). It is currently unclear which aspect of seeing the hand is responsible for the present effects. One previous study found that the enhancing effect of seeing one's body on touch was linked primarily to sense of ownership (Longo et al., 2008a). In contrast, other effects, such as acceleration of visual reaction times (Longo and Haggard, 2009), and proprioceptive integration (Tsakiris et al., 2006) appeared to be related to agency. It is well established that perceived control reduces both the subjective experience of (Weisenberg et al., 1985) and neural responses to (Salomons et al., 2004) pain, via modulation of anterolateral prefrontal regions involved in pain appraisal (Wiech et al., 2006). It is therefore possible that our effects are mediated by an increased sense of bodily control when viewing the hand. This could be investigated in future studies in which participants actively move the viewed hand. Alternatively, however, viewing the hand in pain without being able to withdraw it could also very well produce a perceived reduction of control.

Crossmodal inhibition. Seeing the body has wide-ranging influences on somatosensation. Interestingly, however, the nature of this influence is quite different between touch and pain. Whereas non-informative vision of the hand *increases* the acuity of touch (Kennett et al., 2001) and the amplitude of SEPs (Taylor-Clarke et al., 2002), the present results demonstrate conversely that seeing the hand *decreases* the subjective experience of pain and the amplitude of LEPs. These opposite effects of seeing the body on touch and pain are consistent with the well-established inhibitory interactions between the two modalities themselves. Given the modality segregation

of somatosensory cortex generally (Mountcastle 1957; Friedman et al 2004), and of touch and pain specifically (Ploner et al., 2000; Ohara et al., 2004; Chen et al., 2009), tactile and nociceptive representations may stand in competitive relations.

Competition between *modalities* might mimic the well-established competitive relations between adjacent *skin regions* within a single modality map due to lateral inhibition within the cortex (e.g., Merzenich et al., 1984).

Could a common mechanism produce such divergent effects on touch and pain? One possibility would be a visually-induced crossmodal activation of GABAergic interneurons. Injection of GABA antagonists increases the size of SI tactile RFs (Dykes et al., 1984; Alloway et al., 1989), suggesting that GABAergic interneurons function to sharpen tactile RFs, increasing tactile acuity. Conversely, GABA agonists are effective treatments for chronic central pain (Canavero and Bonicalzi, 1998), suggesting that reduced GABAergic inhibition may be a major cause of chronic pain. Consistent with this interpretation, patients with complex regional pain syndrome (CRPS) show reduced intra-cortical inhibition measured with paired-pulse TMS (Schwenkreis et al., 2003), while repetitive TMS of motor cortex – known to increase intra-cortical inhibition – alleviates chronic pain (Lefaucheur et al., 2006).

Both the present results showing analgesic effects of seeing the body and previous findings showing tactile enhancement (Kennett et al., 2001) could therefore be explained by visual modulation of somatosensory GABAergic interneurons. This speculation is supported by multisensory influences on cortical inhibition in other physiological systems. For example, viewing a hand extends the TMS-evoked silent period compared to viewing a fixation cross (Schütz-Bosbach et al., 2009).

Furthermore, Dehner and colleagues (2004) described a GABA-dependent auditory-somatosensory inhibitory effect in cats.

Therapeutic implications. The present results also have possible therapeutic implications. Several authors have reported that a mirror box similar to that used in our experiment can reduce phantom limb pain (Ramachandran and Rogers-Ramachandran, 1996; Chan et al., 2007) and chronic central pain (McCabe et al., 2003), though replication attempts have been mixed (e.g., Brodie et al 2007; see Moseley et al., 2008a). The present results extend these findings in three ways, suggesting that vision of the body (whether with a mirror box or not) may have a broader range of analgesic effects than previously suspected.

First, the present results show an analgesic effect of vision of the body for acute, rather than chronic, pain. Second, several authors have suggested that mirror therapy may operate by promoting plastic reorganization within somatosensory map, or by correcting a distorted body image through visual recalibration of proprioception (Ramachandran and Rogers-Ramachandran, 1996; Harris, 1999; Ramachandran and Altschuler, 2009). Yet our results demonstrate analgesic effects of seeing the body in healthy participants without body image distortion. Third, previous studies have generally involved voluntary movement of the unaffected limb, inducing the illusion of control over the affected hand. Therapeutic effects are typically attributed to the mirror-induced match between visually-perceived movement and efferent commands specifying movement (e.g., Ramachandran and Rogers-Ramachandran, 1996; McCabe et al 2003). The present results, however, suggest that qualitatively similar analgesic effects may result from simply seeing the hand, independent of movement or match between efferent and afferent signals.

References

- Alloway KD, Rosenthal P, Burton H (1989) Quantitative measurements of receptive field changes during antagonism of GABAergic transmission in primary somatosensory cortex of cats. *Exp Brain Res* 78: 514-532.
- Brodie EE, Whyte A, Niven CA (2007) Analgesia through the looking-glass? A randomized controlled trial investigating the effect of viewing a 'virtual' limb upon phantom limb pain, sensation and movement. *Euro J Pain* 11: 428-436.
- Canavero S, Bonicalzi V (1998) The neurochemistry of central pain: Evidence from clinical studies, hypothesis and therapeutic implications. *Pain* 74: 109-114.
- Chan BL, Witt R, Charrow AP, Magee A, Howard R, Pasquina PF, Heilman KM, Tsao JW (2007) Mirror therapy for phantom limb pain. *New Engl J Med* 357: 2206-2207.
- Chen LM, Friedman RM, Roe AW (2009) Area-specific representation of mechanical nociceptive stimuli within SI cortex of squirrel monkeys. *Pain* 141: 258-268.
- Dehner LR, Keniston LP, Clemo HR, Meredith MA (2004) Cross-modal circuitry between auditory and somatosensory areas of the cat anterior extosylvian sulcal cortex: A 'new' inhibitory form of multisensory convergence. *Cereb Cortex* 14: 387-403.
- Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Meth* 134: 9-21.
- Dykes RW, Landry P, Metherate R, Hicks TP (1984) Functional role of GABA in cat primary somatosensory cortex: Shaping receptive fields of cortical neurons. *J Neurophysio* 52: 1066-1093.
- Ernst M, Lee MHM, Dworkin B, Zaretzky HH (1986) Pain perception decrement

- produced through repeated stimulation. *Pain* 26: 221-231.
- Ernst MO, Banks MS (2002) Humans integrate visual and haptic information in a statistically optimal fashion. *Nature* 415: 429-433.
- Flor H, Denke C, Schaefer M, Grüsser S (2001) Effect of sensory discrimination training on cortical reorganization and phantom limb pain. *Lancet* 357: 1763-1764.
- Friedman RM, Chen LM, Roe AW (2004) Modality maps within primate somatosensory cortex. *Proc Natl Acad Sci USA* 101: 12724-12729.
- Frot M, Rambaud L, Guénot M, Mauguière F (1999) Intracortical recordings of early pain-related CO₂-laser evoked potentials in the human second somatosensory (SII) area. *Clin Neurophysiol* 110: 133-145.
- Garcia-Larrea L, Peyron R, Laurent B, Mauguière F (1997) Association and dissociation between laser-evoked potentials and pain perception. *NeuroReport* 8: 3785-3789.
- Garcia-Larrea L, Frot M, Valeriani M (2003) Brain generators of laser-evoked potentials: From dipoles to functional significance. *Neurophysiol Clin* 33: 279–292.
- Harris AJ (1999) Cortical origin of pathological pain. *Lancet* 354: 1464-1466.
- Harris JA, Arabzadeh E, Moore CA, Clifford CWG (2007) Noninformative vision causes adaptive changes in tactile sensitivity. *J Neurosci* 27: 7136-7140.
- Higgins JD, Tursky B, Schwarz GE (1971) Shock-elicited pain and its reduction by concurrent tactile stimulation. *Science* 172: 866-867.
- Inui K, Tsuji T, Kakigi R (2006) Temporal analysis of cortical mechanisms for pain relief by tactile stimuli in humans. *Cereb Cortex* 16: 355-365.
- Jousmäki V, Hari R (1998) Parchment-skin illusion: Sound-biased touch. *Curr Bio* 8.

R190-R191.

Jung T-P, Makeig S, Humphries C, Lee T-W, McKeown MJ, Iragui V, Sejnowski TJ
(2000) Removing electroencephalographic artifacts by blind source separation.

Psychophysio 37: 163-178.

Kennett S, Taylor-Clarke M, Haggard P (2001) Noninformative vision improves the
spatial resolution of touch in humans. *Curr Bio* 11: 1188-1191.

Lefaucheur JP, Drouot X, Ménard-Lefaucheur I, Keravel Y, Nguyen JP (2006)
Motor cortex rTMS restores defective intracortical inhibition in chronic
neuropathic pain. *Neurology* 67: 1568-1574.

Longo MR, Haggard P (2009) Sense of agency primes manual motor responses.
Perception 38: 69-78.

Longo MR, Cardozo S, Haggard P (2008a) Visual enhancement of touch and the
bodily self. *Conscious Cogn* 17: 1181-1191.

Longo MR, Schüür F, Kammers MPM, Tsakiris M, Haggard P (2008b) What is
embodiment? A psychometric approach. *Cognition* 107: 978-998.

Lundeberg T (1985) Relief of pain from a phantom limb by peripheral stimulation. *J*
Neurol 232: 79-82.

Maihöfner C, Handwerker H O, Neundörfer B, Birklein F (2003) Patterns of cortical
reorganization in complex regional pain syndrome. *Neurology* 61: 1707-1715.

McCabe CS, Haigh RC, Ring EFJ, Halligan PW, Wall PD, Blake DR (2003) A
controlled pilot study of the utility of mirror visual feedback in the treatment
of complex regional pain syndrome (type 1). *Rheumatol* 42: 97-101.

Melzack R, Wall PD (1965) Pain mechanisms: A new theory. *Science* 150: 971-979.

Merzenich MM, Nelson RJ, Stryker MP, Cynader MS, Schoppmann A, Zook JM

- (1984) Somatosensory cortical map changes following digit amputation in adult monkeys. *J Comp Neurol* 224: 591-605.
- Milne RJ, Kay NE, Irwin RJ (1991) Habituation to repeated painful and non-painful cutaneous stimuli: A quantitative psychophysical study. *Exp Brain Res* 87: 438-444.
- Moriwaki K, Yuge O (1999) Topographical features of cutaneous tactile hypoesthetic and hyperesthetic abnormalities in chronic pain. *Pain* 81: 1-6.
- Moseley GL, Wiech K (2009) The effect of tactile discrimination training is enhanced when patients watch the reflected image of their unaffected limb during training. *Pain* 144: 314-319.
- Moseley GL, Gallace A, Spence C (2008a) Is mirror therapy all it is cracked up to be? Current evidence and future directions. *Pain* 138: 7-10.
- Moseley GL, Zalucki NM, Wiech K (2008b) Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain. *Pain* 137: 600-608.
- Mountcastle VB (1957) Modality and topographic properties of single neurons in cat's somatic sensory cortex. *J Neurophysiol* 20: 408-434.
- Ohara S, Crone NE, Weiss N, Treede R-D, Lenz FA (2004) Cutaneous painful laser stimuli evoke responses recorded directly from primary somatosensory cortex in awake humans. *J Neurophysio* 91: 2734-2746.
- Pleger B, Ragert P, Schwenkreis P, Förster A-F, Wilimzig C, Dinse H, Nicolas V, Maier C, Tegenthoff M (2006) Patterns of cortical reorganization parallel tactile discrimination and pain intensity in complex regional pain syndrome. *Neuroimage* 32: 503-510.
- Ploner M, Schmitz F, Freund H-J, Schnitzler A (2000) Differential organization of

touch and pain in human primary somatosensory cortex. *J Neurophysiol* 83: 1770-1776.

Ramachandran VS, Altschuler AL (2009) The use of visual feedback, in particular mirror visual feedback, in restoring brain function. *Brain* 132: 1693-1710.

Ramachandran VS, Rogers-Ramachandran D (1996) Synaesthesia in phantom limbs induced with mirrors. *Proc R Soc Lond B* 263: 377-386.

Ramachandran VS, Rogers-Ramachandran D, Cobb S (1995) Touching the phantom limb. *Nature* 377: 489-490.

Salomons TV, Johnstone T, Backonja MM, Davidson RJ (2004) Perceived controllability modulates the neural response to pain. *J Neurosci* 24: 7199-7203.

Schütz-Bosbach S, Avenanti A, Aglioti SM, Haggard P (2009) Don't do it! Cortical inhibition and self-attribution during action observation. *J Cogn Neurosci* 21: 1215-1227.

Schwenkreis P, Janssen F, Rommel O, Pleger B, Völker B, Hosbach I, Dertwinkel R, Maier C, Tegenthoff M (2003) Bilateral motor cortex disinhibition in complex regional pain syndrome (CRPS) type I of the hand. *Neurology* 61: 515-519.

Spiegel J, Hansen C, Treede R-D (1996) Laser-evoked potentials after painful hand and foot stimulation in humans: Evidence for generation of the middle-latency component in the secondary somatosensory cortex. *Neurosci Lett* 216: 179-182.

Taylor-Clarke M, Kennett S, Haggard P (2002) Vision modulates somatosensory cortical processing. *Curr Bio* 12: 233-236.

- Tipper SP, Lloyd D, Shorland B, Dancer C, Howard LA, McGlone F (1998) Vision influences tactile perception without proprioceptive orienting. *Neuroreport* 9: 1741-1744.
- Treede RD, Lorenz J, Baumgärtner U (2003) Clinical usefulness of laser-evoked potentials. *Neurophysiol Clin* 33: 303-314.
- Tsakiris M, Prabhu G, Haggard P (2006) Having a body *versus* moving your body: How agency structures body-ownership. *Conscious Cogn* 15: 423-432.
- Valeriani M, Betti V, Le Pera D, De Armas L, Miliucci R, Restuccia D, Avenanti A, Aglioti SM (2008) Seeing the pain of others while being in pain: A laser-evoked potentials study. *Neuroimage* 40: 1419-1428.
- Valeriani, M, de Tomasso M, Restuccia D, Le Pera D, Guido M, Iannetti GD, Libro G, Truini A, Di Trapini G, Puca F, Tonali P, Cruccu G (2003) Reduced habituation to experimental pain in migraine patients: A CO2 laser evoked potential study. *Pain* 105: 57-64.
- Wall PD, Sweet WH (1967) Temporary abolition of pain in man. *Science* 155: 108-109.
- Weisenberg M, Wolf Y, Mittwoch T, Mikulincer M, Aviram O (1985) Subject versus experimenter control in the reaction to pain. *Pain* 23: 187-200.
- Wiech K, Kalisch R, Weiskopf N, Pleger B, Stephan KE, Dolan RJ (2006) Anterolateral prefrontal cortex mediates the analgesic effect of expected and perceived control over pain. *J Neurosci* 26: 11501-11509.

Footnotes

1) The first four participants in Experiment 2 made VAS ratings after each 10 trials, rather than every trial.

Figure Captions

Figure 1: The mirror-box technique in which the subject has the experience of viewing their right hand, while in fact seeing their left hand reflected in a mirror.

Figure 2: Schematic depiction of experimental setups. Apparent line of sight depicted with dashed line, actual line of sight with solid line.

Figure 3: Subjective report data from Experiment 1 (left) and Experiment 3 (right). Error bars are one SEM.

Figure 4: Subjective ratings of pain intensity and unpleasantness in the three experiments. Error bars are one SEM.

Figure 5: Grand mean LEPs recorded from electrode Cz in the three experiments (left panels) and N2/P2 peak-to-peak amplitudes at Cz (right panels). Error bars are one SEM.

Figure 1



Figure 2




Figure 3




Figure 4




Figure 5

